

Palmer, A.R. & Repina, L.N. in
1997

SYSTEMATIC DESCRIPTIONS OF THE CLASS TRILOBITA

H. B. WHITTINGTON, W. T. CHANG, W. T. DEAN, R. A. FORTEY, P. A. JELL,
J. R. LAURIE, A. R. PALMER, L. N. REPINA, A. W. A. RUSHTON, and J. H.
SHERGOLD

[Diagnoses of ordinal and higher taxa by R. A. FORTEY and H. B. WHITTINGTON; authorship
of other diagnoses indicated within.]

Class TRILOBITA Walch, 1771

[*nom. correct. ex* Trilobiten WALCH, 1771, p. 120] [This name, as
"Trilobiten," was published by WALCH (1771, p. 120) in his volume III of
KNORR and WALCH (1768–1774). WALCH's work was rejected for nomen-
clatorial purposes by the International Commission on Zoological Nomen-
clature (Opinion 529, 1958). However, the Commission's remit extends
only to the family-group level (Article 1b(4), *International Code of Zoolo-
gical Nomenclature*, 1985), hence the name, emended to "Trilobita," remains
available. The designation of the genus *Dalmanites* BARRANDE as type
(HENNSMOEN & MOORE in MOORE, 1959, p. 172) is invalid.]

Extinct marine arthropods, body divided
by furrows longitudinally into median (axial)
and lateral (pleural) regions, transversely into
cephalon of fused segments, thorax of articu-
lated segments, and pygidium of few to
many fused segments. Length of mature in-
dividuals ranges from a little over 1 mm to
about 700 mm.

Dorsal exoskeleton and periphery of ven-
ter (doublure) calcified (but see comments
on Naraoiidae); calcified plate (hypostome)
anterior to mouth and below anterior axial
region. Anterior wing of hypostome extend-
ing dorsally to a fossula in the axial furrow.
Facial suture characteristic, absent in
Olenellina, but may be secondarily lost or
modified. Rostral plate or median suture
present, or cheeks conjoined. Cephalon with
one pair of eyes (which may be lost), each eye
lobe connected to axial region by raised
ridge; eye lenses of crystalline calcite. Slightly
oblique-transverse pleural furrows at poste-
rior of cephalon, on each thoracic segment,
and on pygidial segments. Growth of exo-
skeleton proceeded by molting from calcified
protaspis about 1 mm long to adult, and by
sequential release of segments into thorax.
Protaspides of Olenellina and Agnostina not
known. Limbs (where known) include one
pair of antennae, followed postorally by se-
ries of generally similar biramous limbs, of
which three (possibly four in one species) are

cephalic. *Lower Cambrian (Atdabanian)*—
Upper Permian.

Order AGNOSTIDA Salter, 1864

[*nom. correct. SHERGOLD, LAURIE, & SUN, 1990, p. 32, pro Agnostini
SALTER, 1864a, p. 2*] [=Miomera JAEKEL, 1909, p. 394; Agnostida
KOBAYASHI, 1935, p. 81; see discussion in ROBISON & CAMPBELL, 1974,
p. 281]

Diminutive, isopygous, 2 or 3 fulcate
thoracic segments. Cephalic shield with
deeply parabolic outline and maximum
width (tr.) usually anterior to genal angle;
border convex; glabella commonly fusiform,
widest at base (except in Condylopygidae).
Hypostome natant; rostral plate lacking or
uncalcified. Outline of pygidium closely
matching that of cephalon. *Lower Cam-
brian–Upper Ordovician.*

Suborder AGNOSTINA Salter, 1864

[*nom. correct. SHERGOLD, LAURIE, & SUN, 1990, p. 32, pro Agnostini
SALTER, 1864a, p. 2*]

No eye or facial suture; 2 thoracic seg-
ments; cephalothoracic aperture present;
pygidium generally with wide axis, margin
commonly bearing posterolateral spine. Cu-
ticle thin. Calcified protaspis not known.
Lower Cambrian–Upper Ordovician.

INTRODUCTION TO THE SUBORDER AGNOSTINA

JOHN H. SHERGOLD and JOHN R. LAURIE

The historical background to the classifi-
cation of this suborder has recently been dis-
cussed in detail (SHERGOLD, LAURIE, & SUN,
1990, p. 1–3). The classification presented
here is as developed in that paper and essen-
tially follows ideas on relationships that have

short (sag.) or absent (Fig. 30.1). Thorax nonfulcate. Pygidium narrow (tr.), with few segments. Calcified protaspis unknown; earliest meraspid with segmented interocular area. Lower Cambrian.

INTRODUCTION TO SUBORDER OLENELLINA

A. R. PALMER and L. N. REPINA

Olenellina are a morphologically varied and highly diverse group of generally micropygous trilobites that share a primary absence of facial sutures, the presence of a well-developed ocular lobe at all developmental stages, and an ontogeny in which the first mineralized stages are already early meraspids (PALMER, 1957). They are restricted to and characteristic of rocks of later Early Cambrian age and constitute a major suborder within the order Redlichiida (MOORE, 1959). More than 50 genera or subgenera and their associated higher taxa are recognizable (PALMER & REPINA, 1993). These taxa form the principal basis for biostratigraphic subdivisions (Fig. 254) of the later Lower Cambrian rocks of Laurentia (North America exclusive of the eastern seaboard from Newfoundland to Florida and including Spitsbergen and northwestern Scotland) and are major indices for the later Lower Cambrian biostratigraphy of Baltica (northern Europe exclusive of the British Isles), Avalonia (England, Wales, eastern Newfoundland, Nova Scotia, eastern New England, USA), Siberia, and the Moroccan sector of Gondwana (including Spain).

Ancestry of the Olenellina can only be speculative. They first appear as fully developed and morphologically diverse trilobites in the early, but not earliest, part of the shelly fossil record and are the oldest trilobites known. The principal phylogenetic trend within the Olenellina involves the relationship between the ocular lobe and the frontal (anterior) lobe (LA) of the glabella (REPINA, 1990a). In all of the earliest Olenellina, which include forms from Siberia, Laurentia, and the Moroccan sector of Gondwana, the

glabella has parallel sides or tapers forward, LA is short, and the ocular lobe is attached along the entire margin of LA (Fallotaspidinae, Daguinaspidinae). In later genera, LA first becomes elongate, so that the ocular lobes connect only to its posterior part (Judomiidae, Nevadiidae), and then generally expands laterally, and the glabella as a whole expands anteriorly from the level of S1 (Olenellidae, Holmiidae). LA in these genera is also commonly inflated. Accompanying this modification, the distal parts of L3 extend laterally and posterolaterally and encroach on L2, often isolating the S2 furrows (Fig. 255), and L3 takes on a broad M-shape.

This phylogenetic trend underlies the proposed classification of the superfamilies. Within the Fallotaspiidoidea, all Fallotaspididae, the earliest family of the Olenellina, have the ocular lobe attached along the entire margin of LA and an unmodified L3, and the anterior end of LA does not project forward of a line tangent to the anterolateral margin of the ocular lobe. In the remaining, generally younger families of the Fallotaspiidoidea (Archaeaspidoidea, Judomiidae, Neltneriidae, Nevadiidae), the anterior end of LA projects forward of the junction with the anterior margin of the ocular lobe, but L3 remains unmodified. All Olenelloidea, which includes the youngest Olenellina, have the ocular lobe attached to the posterior part of LA and have a modified L3.

Within the superfamilies of the Olenellina, morphological changes of taxonomic value at the family level follow different patterns. The principal character distinguishing the two families in the Olenelloidea is the relationship between the extraocular area and the interocular area. The Holmiidae all have a narrow extraocular area that is less than twice the width of the interocular area. With minor exceptions (*Olenelloides* and some undescribed Laurentian forms), the Olenellidae have a wide extraocular area that is more than twice the width of the interocular area. In the Fallotaspiidoidea, the relationship between the ocular lobe and LA, the shape of the glabella (parallel sided versus tapered),

Mummaspis FRITZ, 1992, p. 17 [**Wanneria occidens* WALCOTT, 1913b, p. 314; OD; holotype (WALCOTT, 1913b, pl. 53, fig. 2), 60080, USNM, Washington, D.C.]. Parts of external surface reticulate. Posterior margin of cephalon nearly straight; intergenital swelling slightly distal to midlength of posterior margin. Preglabellar field absent or length (sag.) less than that of border; S3 deep, continuous across glabella; occipital spine may be present. Ocular furrow deep; outer band of ocular lobe narrower than inner band. Third thoracic segment generally only weakly macropleural. Lower Cambrian: Canada (southern Rocky Mountains), lower *Olenellus* Zone.——FIG. 256, 1a, b. **M. occidens* (WALCOTT); a, complete individual, topotype, USNM 443745, X4 (Fritz, 1992, pl. 9, fig. 2); b, cephalon and partial thorax, topotype, USNM 433750, X1.7 (Fritz, 1992, pl. 10, fig. 2).

Subfamily BICERATOPSINAE Pack & Gayle, 1971

[Biceratopsinae PACK & GAYLE, 1971, p. 895] [=Olenellidae BERGSTROM, 1973b, p. 312, partim; AHLBERG, BERGSTROM, & JOHANSSON, 1986, p. 40, partim; Biceratopsinae REPINA, 1979, p. 22; PALMER & REPINA, 1993, p. 23]

Glabella narrowest at midlength; glabellar furrows very poorly developed; width (tr.) of L1 less than width (tr.) of occipital ring; ocular lobe close to glabella, its posterior tip opposite or anterior to L1. Genal spine absent or opposite midlength of ocular lobe; intergenital spine absent; intergenital angle may be developed. Thorax divided into prothorax and opisthothorax; third segment strongly expanded distally; pleural spine extremely long, with tips posterior to body; fifteenth segment with long axial spine. Lower Cambrian.

Biceratops PACK & GAYLE, 1971, p. 895 [**B. nevadensis*; OD; holotype (PACK & GAYLE, 1971, pl. 102, fig. 2–3), 168225, USNM, Washington, D.C.]. Posterior margin of cephalon nearly straight; genal angle broadly rounded; genal and intergenital spines absent. Length (sag.) of preglabellar field equal to or less than that of anterior border; ocular lobe prominent, close to glabella, with posterior tip opposite S1 and elevated above top of glabella. Opisthothorax of at least 11 segments. Lower Cambrian: USA (Arizona, Nevada), upper *Olenellus* Zone.——FIG. 257, 1. **B. nevadensis*, Nevada; complete individual, holotype, USNM 168225, X3 (Pack & Gayle, 1971, pl. 102, fig. 2).

Peachella WALCOTT, 1910, p. 342 [**Olenellus iddingsi* WALCOTT, 1884b, p. 28; OD; holotype (WALCOTT, 1884b, pl. 9, fig. 12), 15407a, USNM, Washington, D.C.]. Posterior margin of cephalon nearly straight. Glabella extended to or nearly to border furrow; posterior tip of ocular lobe approximately

opposite S1; genal spine at posterolateral corner or slightly advanced, strongly inflated. Opisthothorax of at least 10 segments. Lower Cambrian: USA (California, Nevada), upper *Olenellus* Zone.——FIG. 257, 2. **P. iddingsi* (WALCOTT), California; nearly complete individual, LACMIP 11627, X2 (Palmer & Repina, 1993, fig. 4.2). 2

Subfamily BRISTOLIINAE Palmer & Repina, 1993

[Bristoliniinae PALMER & REPINA, 1993, p. 23] [=Olenellinae POULSEN in MOORE, 1959, p. 162, partim; Olenellidae BERGSTROM, 1973b, p. 312, partim; AHLBERG, BERGSTROM, & JOHANSSON, 1986, p. 40, partim; Fremontiinae REPINA, 1979, p. 22, partim]

Glabella usually strongly constricted at S1 or L2; width (tr.) of anterior part of L1 usually distinctly narrower than occipital ring; glabellar furrows generally well developed. Preglabellar field shorter (sag.) than anterior border or absent. Posterior tip of ocular lobe opposite or anterior to L1. Third thoracic segment with inner part of pleural region strongly expanded distally; pleural spine may be extended posterior to end of thorax. Opisthothorax well developed. Pygidium not known. Lower Cambrian.

Bristolia HARRINGTON, 1956, p. 59 [**Mesonacis bristolensis* RESSER, 1928, p. 7; OD; lectotype (RESSER, 1928, pl. 2, fig. 5–6; SD HARRINGTON, 1956, p. 59), 78390, USNM, Washington, D.C.]. Cephalon subpentagonal to subquadrate in outline; genal spine originating opposite or anterior to L2; intergenital spine absent. Preglabellar field absent; posterior tip of ocular lobe opposite or anterior to L1. Fifteenth thoracic segment with long axial spine; opisthothorax of at least 17 segments. Lower Cambrian: USA (California, Nevada), Greenland (Inglefield Land), *Olenellus* Zone.——FIG. 258, 1. **B. bristolensis*, California; nearly complete individual, UCR 10/7, X1 (Palmer & Repina, 1993, fig. 4.5).

Arcuolenellus PALMER & REPINA, 1993, p. 24 [**Olenellus arcuatus* PALMER in PALMER & HALLEY, 1979, p. 67; OD; holotype (PALMER & HALLEY, 1979, pl. 2, fig. 12), 177200, USNM, Washington, D.C.]. Posterior margin of cephalon strongly and evenly curved back distally; genal spine short, situated at posterolateral corner well behind level of occipital ring; intergenital spine absent. Preglabellar field short (sag.); width of interocular area about half width of glabella at L2; posterior tip of ocular lobe opposite L1. Lower Cambrian: USA (California), upper *Olenellus* Zone; Argentina (San Juan area), *Olenellus* Zone.——FIG. 258, 2. **A. arcuatus* (PALMER), California; cephalon, holotype, USNM 177200, X6 (Palmer & Halley, 1979, pl. 2, fig. 12). **Bolbolenellus** PALMER & REPINA, 1993, p. 24 [**Olenellus euryparia* PALMER in PALMER & HALLEY,




FIG. 257. Olenellidae (p. 409)

posterior margin. Genal spine originating opposite or anterior to S1. Glabella bullet shaped. Ocular lobe close to glabella, its tip opposite or posterior to occipital furrow. Thorax with at least 17 segments; each segment with width (tr.) of thoracic axis greater than width of inner parts of pleural region; third thoracic segment not modified; fifteenth segment with long axial spine.

Opisthothorax not clearly differentiated. Pygidium elongate; sides convergent posteriorly; end bluntly pointed. *Lower Cambrian*.

Gabriellus FRITZ, 1992, p. 20 [**G. lanceatus*; OD; holotype (FRITZ, 1992, pl. 17, fig. 6), 443792, USNM, Washington, D.C.] Characters as for subfamily. *Lower Cambrian*: USA (Nevada), Canada (Cordilleran region), lower *Olenellus* Zone.—FIG. 259,3. *Gabriellus* sp.; complete individual, Canada, GSC 104195, X1.7 (Palmer & Repina, 1993, fig. 4.9).


Subfamily LAUDONIINAE Palmer & Repina, 1993

[Laudoniinae PALMER & REPINA, 1993, p. 24]

Cephalon subquadrate to subhexagonal in outline. Genal spine strongly advanced, originating anterior to S1. Intergenal spine strongly developed in adult at posterolateral corner of cephalon. Width (tr.) of anterior part of L1 nearly equal to that of occipital ring. Thorax with third segment weakly macropleural; 15th segment lacking axial spine. Opisthothorax not clearly differentiated. Pygidium small, bilobate. *Lower Cambrian*.

Laudonia HARRINGTON, 1956, p. 60 [**L. bispinata*; OD; holotype (HARRINGTON, 1956, pl. 15, fig. 4), 9465T1, KUMIP, Lawrence]. Cephalon subquadrate to subpentagonal in outline; genal spine originating opposite or anterior to L3; procranial spine not developed in adult; intergenal ridge distinct. Preglabellar field absent or shorter (sag.) than border; posterior tip of ocular lobe opposite or anterior to L1. Thorax of about 20 segments. External surface reticulate. *Lower Cambrian*: Canada (southern Rocky Mountains), USA (Nevada), Mexico (Caborca), lower *Olenellus* Zone.—FIG. 259,2. *L. amputata* FRITZ; cephalon and partial thorax, Canada, USNM 443754, X2 (Fritz, 1992, pl. 11, fig. 4).

?*Olenelloides* PEACH, 1894, p. 668 [**Olenellus (Olenelloides) armatus* PEACH, 1894, p. 669; OD; lectotype (PEACH, 1894, pl. 32, fig. 4; SD McNAMARA, 1978, p. 637), 472, GSE, Edinburgh]. Cephalon subhexagonal in outline; prominent procranial, genal, or intergenal spine at each angle of the hexagon; intergenal spine close to glabella; LA subglobular. Preglabellar field absent; width (tr.) of narrow interocular area about equal to width of equally narrow extraocular area opposite midlength of ocular lobe; posterior tip of ocular lobe opposite L2. Thorax with 9 segments; axis wider than inner part of pleural region exclusive of spines; third and sixth thoracic segments macropleural. *Lower Cambrian*: northwestern Scotland, middle *Olenellus* Zone.—FIG. 259,1a,b. **O. armatus* (PEACH); a,



1

Neltneria



2

Bondonella

FIG. 268. Neltneriidae (p. 426)

Posterior margin of cephalon deflected slightly forward distally. Glabella parallel sided, bluntly rounded anteriorly. LA elongate. Ocular lobe connected only to posterior part of LA. Ocular lobe close to glabella; interocular area poorly differentiated. Thorax with 17 segments. Lower Cambrian.

Neltneria HUPÉ, 1953a, p. 133 [**Wanneria Jacqueti* NELTNER & POCTEY, 1949, p. 74; OD; lectotype (NELTNER & POCTEY, 1949, pl. 6, fig. 5; SD PALMER & REPINA, 1993, p. 30), R50864, MNHN, Paris]. Preglabellar field absent; interocular area poorly defined, its width (tr.) less than half that of extraocular area; posterior tip of ocular lobe opposite L1. Thorax has 17 segments; each segment with short (exsag.) pleural spine; width (tr.) of inner pleural region about same as that of axis; 11th segment macropleural; 12th to 17th segments decreasing in width (tr.) rapidly and progressively backward. Pygidium small, subquadrate, possibly of 2 or 3 segments. Lower Cambrian: Morocco (Anti-Atlas), *Antatlasia gutta-pluviae* Zone.—FIG. 268, 1. **N. jacqueti*; complete individual, ICS replica 122, $\times 2$ (Palmer & Repina, 1993, fig. 11.1).

Bondonella HUPÉ, 1953a, p. 135 [**B. typica* HUPÉ, 1953a, p. 136; OD; holotype (NELTNER & POCTEY, 1949, pl. 6, fig. 1), R50865, MNHN, Paris]. Length (sag.) of preglabellar field about equal to that of anterior border; interocular area poorly defined, its width (tr.) more than half width of extraocular area; posterior tip of ocular lobe opposite occipital furrow. Thorax with 17 segments; width (tr.) of inner pleural region of each segment narrower than axis; no macropleural segments. Lower Cambrian: Morocco (Anti-Atlas), *Antatlasia gutta-pluviae* Zone.—FIG. 268, 2. **B. typica*; complete individual, holotype, MNHN R50865, $\times 3$ (Palmer & Repina, 1993, fig. 11.2).

Family NEVADIIDAE Hupé, 1953

[nom. transl. REPINA, 1979, p. 19, ex *Nevadiinae* HUPÉ, 1953a, p. 124] [=Nevadiinae POULSEN in MOORE, 1959, p. 196; SÜVOROVA in TCHERNYSHEVA, 1960, p. 62; BERGSTROM, 1973b, p. 309, partim; Nevadiidae REPINA, 1979, p. 19, partim; AHLBERG, BERGSTROM, & JOHANSSON, 1986, p. 40, partim; PALMER & REPINA, 1993, p. 31]

Glabella weakly to strongly tapered forward in dorsal view. LA elongate. Ocular lobe connected only to posterior part of LA. Interocular area narrow. Posterior tips of ocular lobes opposite or anterior to occipital furrow. Width of interocular area variable. Thorax with 15 to 27 segments, without development of macropleurae. Lower Cambrian.

Nevadia WALCOTT, 1910, p. 256 [**N. weeksii*; OD; lectotype (WALCOTT, 1910, pl. 23, fig. 2; SD PALMER




FIG. 269. Nevadiidae (p. 426–428)



Poletaevella

FIG. 270. Superfamily Uncertain (p. 428)

& REPINA, 1993, p. 31), 56792b, USNM, Washington, D.C.). Width (tr.) of interocular area less than one-fourth width of extraocular area; preglabellar field present. Thorax with 27 segments; pleural spines long; width (tr.) of pleurae at least twice width of axis; transition to opisthothorax between segments 15 and 18. Pygidium small, subquadrate. Lower Cambrian: USA (California, White-Inyo region; Nevada; northeastern Washington), Canada (northern Rocky Mountains), "Nevadella" Zone; Russia (northern Siberian Platform), "Nevadella" Zone. —FIG. 269,1. **N. eucharis*; complete individual, Canada, holotype, USNM 60079, X2 (Palmer & Repina, 1993, fig. 13.2).

Buenellus BLAKER, 1988, p. 34 [*B. higginsi* BLAKER, 1988, p. 36; OD; holotype (BLAKER, 1988, fig. 3b), 18.287, MGUH, Copenhagen]. Glabella slightly tapered forward; ocular lobe small, posterior tip about opposite S1 or anterior part of L1; width (tr.) of interocular area equal to or greater than that of extraocular area. Thorax with 17 or 18 segments, maintaining width or widening slightly backward to 8th segment, then tapering posteriorly; posterior segment may be fused with anterior part of simple pygidium; pleural spines short (exsag.); inner pleural regions only slightly wider (tr.) than axis. Lower Cambrian: Greenland (Peary Land), "Nevadella" Zone; Russia, Novaya Zemlya, ?"Nevadella" Zone. —FIG. 269,4. **B. higginsi*, Greenland; complete individual, paratype, MGUH 17.589, X3 (Blaker, 1988, fig. 5a).

Cirquella FRITZ, 1993, p. 858 [*C. nummularia*; OD; holotype (FRITZ, 1993, fig. 7.2-7.4), 102330, GSC, Ottawa]. Glabella moderately tapered forward; glabellar furrows weakly developed; posterior tip of ocular lobe opposite L1. Genal spine absent or indicated only by slight angularity of margin opposite L1. Lower Cambrian: USA (White-Inyo and western Death Valley regions), Canada (southern Rocky Mountains), lower "Nevadella" Zone. —FIG. 269,5. **C. nummularia*; cephalon, Canada, paratype, USNM 458891, X7 (Fritz, 1993, fig. 7.5).

Nevadella RAW, 1936, p. 250 [**Callavia eucharis* WALCOTT, 1913b, p. 315; SD WHITEHOUSE, 1939, p. 191; holotype (WALCOTT, 1913b, pl. 53, fig. 1), 60079, USNM, Washington, D.C.]. Glabella distinctly tapered forward, sides slightly concave, narrowest opposite L2 or S2; preglabellar field short or absent; basal glabellar width (tr.) about equal to width of extraocular area opposite midlength of ocular lobe; posterior tip of ocular lobe opposite L1; width (tr.) of interocular area more than one-fourth width of extraocular area opposite midlength of ocular lobe. Thorax with 17 to 23 segments. Pleural spines long, falcate; opisthothorax not clearly differentiated. Pygidium small, subquadrate. Lower Cambrian: USA (California, Nevada), Canada (Cordilleran region), "Nevadella" Zone. —FIG. 269,1. **N. eucharis*; complete individual, Canada, holotype, USNM 60079, X2 (Palmer & Repina, 1993, fig. 13.2).

Pseudojudomia EGOROVA in GORYANSKII, EGOROVA, & SAVITSKII, 1964, p. 22 [**P. egregia*; OD; holotype (GORYANSKII, EGOROVA, & SAVITSKII, 1964, pl. 5, fig. 1a,b), 8363/15, CNIGR, St. Petersburg]. Border furrows weakly developed; cephalon strongly convex (tr.); posterior margin strongly curved backward. Glabella slightly tapered forward; basal glabellar width (tr.) about equal to width of extraocular area. Lateral parts of poorly differentiated border nearly vertical. Ocular lobe poorly differentiated from interocular area; posterior tip about opposite occipital furrow. Lower Cambrian: Russia (northern Siberian Platform), *Pseudojudomia* Zone. —FIG. 269,3. **P. egregia*; cephalon, holotype, CNIGR 8363/15, X1.5 (Egorova & Savitskii, 1969, pl. 10, fig. 1).

Superfamily Uncertain

Poletaevella DALMATOV & REPINA, 1971, p. 125 [**P. baljutica*; OD; holotype (DALMATOV & REPINA, 1971, fig. 2b-d, 3), 0213/11, BGU, Ulan-Ude]. Olenellinid? with posterior margin of cephalon nearly straight. Glabella parallel sided, bluntly rounded anteriorly; occipital ring unusually short (sag.); palpebral lobe short (exsag.), prominent, located about midway between axial and lateral border furrows, and connected to posterior part of LA by narrow eye ridge, with its posterior tip opposite L2; preglabellar field longer (sag.) than anterior border. Genal spine well developed; intergenal spine not indicated. [The structure of the palpebral lobe is unlike the ocular lobe of all olenellinid trilobites. Assignment of this form even to the Olenellina is questionable.] Lower Cambrian, Russia (eastern Sayan region), zone unknown. —FIG. 270. **P. baljutica*; cephalon, holotype, X13 (Dalmatov & Repina, 1971, fig. 2b).

Postfallotaspis ORLOWSKI, 1985, p. 234 [**P. spinatus*; OD; holotype (ORLOWSKI, 1985, pl. 3, fig. 5a-d), 1.496, IGUW, Warsaw]. The type and only specimen of this genus is too incomplete and poorly illustrated for generic or higher level taxonomic

TREATISE ON INVERTEBRATE PALEONTOLOGY

Part O ARTHROPODA 1 Trilobita, Revised

Volume 1: Introduction,
Order Agnostida, Order Redlichiida

H. B. WHITTINGTON, B. D. E. CHATTERTON, S. E. SPEYER, R. A. FORTEY,
R. M. OWENS, W. T. CHANG, W. T. DEAN, R. A. FORTEY, P. A. JELL,
J. R. LAURIE, A. R. PALMER, L. N. REPINA, A. W. A. RUSHTON,
J. H. SHERGOLD, E. N. K. CLARKSON, N. V. WILMOT, AND S. R. A. KELLY

*Prepared under Sponsorship of
The Geological Society of America, Inc.*

*The Paleontological Society
The Palaeontographical Society*

*SEPM (Society for Sedimentary Geology)
The Palaeontological Association*

RAYMOND C. MOORE
Founder

ROGER L. KAESLER
Editor

ELIZABETH BROSIUS, JANE KERNIS,
JILL HARDESTY, JACK KEIM, AND KAREN RENTERIA
Assistant Editors and Editorial Staff

THE GEOLOGICAL SOCIETY OF AMERICA, INC.
and
THE UNIVERSITY OF KANSAS
BOULDER, COLORADO, and LAWRENCE, KANSAS
1997