

Technical Series
Volume 8
Number 1
Part 1
November 1986




Joel W. Martin

State of Florida
Department of
Environmental Regulation

An Illustrated Guide to the Marine Decapod Crustaceans of Florida

Lawrence G. Abele
and
Won Kim


Illustrated by
Elizabeth Woodsmall



Technical Series
Volume 8
Number 1
Part 2
November 1986

Joel W. Martin

State of Florida
**Department of
Environmental Regulation**

An outline map of the state of Florida, including the panhandle and the Keys, serves as a background for the title. A shrimp is illustrated in the upper right corner, and a crab is in the middle left. A large crayfish is at the bottom left.

An Illustrated Guide to the Marine Decapod Crustaceans of Florida

**Lawrence G. Abele
and
Won Kim**

**Illustrated by
Elizabeth Woodsmall**

An Illustrated Guide to the Marine Decapod Crustaceans of Florida

Part 1

**Lawrence G. Abele
and
Won Kim**

**Florida State University
Department of Biological Science
Tallahassee, Florida 32306**

**Illustrated by
Elizabeth Woodsmall**

**Copyright © 1986 The Florida State University.
All rights reserved.**

**An Illustrated Guide
to the
Marine Decapod Crustaceans
of Florida**

Part 2

**Lawrence G. Abele
and
Won Kim**

**Florida State University
Department of Biological Science
Tallahassee, Florida 32306**

**Illustrated by
Elizabeth Woodsmall**

**Copyright © 1986 The Florida State University.
All rights reserved.**

Contents

List of Species.....	ii
Introduction.....	1
Brief Review of the Literature.....	1
Methods and Materials.....	2
Classification and Arrangement of Taxa.....	2
Acknowledgments.....	3
Diagrammatic Illustration of a Shrimp.....	4
Diagrammatic Illustration of a Crab.....	5
Annotated Checklist of the Decapod Crustaceans of Florida.....	7
Addendum to Checklist.....	69
Key to Families of Florida Decapods.....	70
Keys to Species of Florida Decapods.....	79
Literature Cited.....	731
Taxonomic Index.....	748

List of Species

[See also Addendum to Checklist]

Suborder Dendrobranchiata

Family Aristeidae

1. *Aristaeomorpha foliacea* (Risso, 1827)..... 7, 79, 81
2. *Plesiopenaeus edwardsianus* (Johnson, 1867)..... 7, 79, 81

Family Benthescymidae

3. *Bentheogennema intermedia* (Bate, 1888)..... 7, 79, 81

Family Penaeidae

4. *Funchalia villosa* (Bouvier, 1905)..... 7, 82, 95
5. *Metapenaeopsis gerardoi* Pérez Farfante, 1971..... 7, 83, 87
6. *Metapenaeopsis goodei* (Smith, 1885)..... 7, 83, 87
7. *Metapenaeopsis smithi* (Schmitt, 1924)..... 7, 83, 87
8. *Parapenaeus americanus* (Rathbun, 1901)..... 7, 83, 89
9. *Parapenaeus politus* Smith, 1881..... 7, 83, 89
10. *Penaeopsis serrata* Bate, 1881..... 8, 82, 95
11. *Penaeus aztecus* Ives, 1891..... 8, 84, 85, 91
12. *Penaeus brasiliensis* Latreille, 1817..... 8, 84, 91
13. *Penaeus duorarum* Burkenroad, 1939..... 8, 84, 85, 91
14. *Penaeus setiferus* (Linnaeus, 1767)..... 8, 84, 91
15. *Trachypenaeus constrictus* (Stimpson, 1874)..... 8, 85, 93
16. *Trachypenaeus similis* (Smith, 1885)..... 8, 85, 93
17. *Trachypenaeopsis mobilispinis* (Rathbun, 1920)..... 8, 82, 95
18. *Xiphopenaeus kroyeri* (Heller, 1862)..... 8, 82, 95

Family Solenoceridae

19. *Hadropenaeus affinis* (Bouvier, 1906)..... 8, 98, 101
20. *Hadropenaeus modestus* (Smith, 1885)..... 8, 98, 101
21. *Hymenopenaeus aphoticus* Burkenroad, 1936..... 9, 98, 103
22. *Hymenopenaeus debilis* Smith, 1882..... 9, 98, 103
23. *Mesopenaeus tropicalis* (Bouvier, 1905)..... 9, 97, 107
24. *Pleoticus robustus* (Smith, 1885)..... 9, 97, 107
25. *Solenocera atlantis* Burkenroad, 1939..... 9, 99, 105
26. *Solenocera necopina* Burkenroad, 1939..... 9, 99, 105
27. *Solenocera vioscai* Burkenroad, 1939..... 9, 99, 105

Family Sicyoniidae

28. *Sicyonia brevirostris* Stimpson, 1871..... 9, 109, 111
29. *Sicyonia burkenroadi* Cobb, 1971..... 9, 109, 113
30. *Sicyonia dorsalis* Kingsley, 1878..... 10, 109, 113
31. *Sicyonia laevigata* Stimpson, 1871..... 10, 109, 111
32. *Sicyonia parri* (Burkenroad, 1934)..... 10, 109, 111
33. *Sicyonia stimpsoni* Bouvier, 1905..... 10, 109, 113
34. *Sicyonia typica* (Boeck, 1864)..... 10, 109, 111

Family Sergestidae

35. *Acetes americanus carolinae* Hansen, 1933..... 10, 115, 123
36. *Sergestes armatus* Kroyer, 1855..... 10, 116, 121
37. *Sergestes atlanticus* H. Milne Edwards, 1830..... 10, 116, 119
38. *Sergestes edwardsii* Kroyer, 1855..... 10, 116, 121
39. *Sergestes henseni* (Ortmann, 1893)..... 10, 116, 119
40. *Sergestes paraseminudus* Crosnier and Forest, 1973..... 10, 116, 119
41. *Sergestes pectinatus* Sund, 1920..... 10, 116, 119
42. *Sergestes sargassi* Ortmann, 1893..... 10, 116, 121

43. *Sergestes vigilax* Stimpson, 1860..... 11, 116, 121
 44. *Sergia extenuatus* Burkenroad, 1940..... 11, 117, 123
 45. *Sergia splendens* Sund, 1920..... 11, 117, 123

Family Luciferidae

46. *Lucifer faxoni* Borradaile, 1915..... 11, 125, 126a
 47. *Lucifer typus* H. Milne Edwards, 1837..... 11, 125, 126a

Suborder Pleocyemata

Infraorder Stenopodidea

Family Stenopodidae

48. *Microprosthema semilaeve* (Von Martens, 1872)..... 11, 281, 283
 49. *Odontozona libertae* Gore, 1981..... 11, 281, 283
 50. *Stenopus hispidus* (Olivier, 1811)..... 11, 281, 283
 51. *Stenopus scutellatus* Rankin, 1898..... 11, 281, 283

Infraorder Caridea

Family Atyidae

52. *Potimirim potimirim* (Muller, 1881)..... 11, 127, 129

Family Oplophoridae

53. *Acanthephyra purpurea* A. Milne Edwards, 1881..... 12, 131, 135
 54. *Janicella spinicauda* (A. Milne Edwards, 1883)..... 12, 131, 135
 55. *Oplophorus gracilirostris* A. Milne Edwards, 1881..... 12, 131, 133
 56. *Oplophorus spinosus* (Brullé, 1839)..... 12, 131, 133
 57. *Systellaspis debilis* (A. Milne Edwards, 1881)..... 12, 131, 135

Family Pasiphaeidae

58. *Leptochela bermudensis* Gurney, 1939..... 12, 137, 139
 59. *Leptochela carinata* Ortmann, 1893..... 12, 137, 139
 60. *Leptochela papulata* Chace, 1976..... 12, 137, 139
 61. *Leptochela serratorbita* Bate, 1888..... 12, 137, 139

Family Bresiliidae

62. *Discias atlanticus* Gurney, 1939..... 12, 141, 143
 63. *Discias serratirostris* Lebour, 1949..... 13, 141, 143
 64. *Pseudosquilla chacei* Kensley, 1983..... 13, 141, 143

Family Eugonatonotidae

65. *Eugonatonotus crassus* (A. Milne Edwards, 1881)..... 13, 145, 147

Family Rhynchocinetidae

66. *Rhynchocinetes rigens* Gordon, 1936..... 13, 145, 147

Family Gnathophyllidae

67. *Gnathophyllodes mineri* Schmitt, 1933..... 13, 149, 151
 68. *Gnathophyllum americanum* Guérin-Méneville, 1855..... 13, 149, 151
 69. *Gnathophyllum circellum* Manning, 1963..... 13, 149, 151
 70. *Gnathophyllum modestum* Hay, 1917..... 13, 149, 151

Family Palaemonidae

71. <i>Anchistioides antiquensis</i> (Schmitt, 1924).....	13, 153, 191
72. <i>Brachycarpus biunguiculatus</i> (Lucas, 1849).....	13, 152, 191
73. <i>Leander paulensis</i> Ortmann, 1897.....	13, 154, 163
74. <i>Leander tenuicornis</i> (Say, 1818).....	13, 154, 163
75. <i>Lipkebe holthuisi</i> Chace, 1969.....	14, 153, 191
76. <i>Macrobrachium acanthurus</i> (Wiegmann, 1836).....	14, 154, 165
77. <i>Macrobrachium carcinus</i> (Linnaeus, 1758).....	14, 154, 165
78. <i>Macrobrachium crenulatum</i> Holthuis, 1950.....	14, 154, 167
79. <i>Macrobrachium ohione</i> (Smith, 1874).....	14, 154, 165
80. <i>Macrobrachium olfersii</i> (Wiegmann, 1836).....	14, 154, 165
81. <i>Neopontonides beaufortensis</i> (Borradaile, 1920).....	14, 153, 191
82. <i>Palaemon floridanus</i> Chace, 1942.....	14, 155, 169
83. <i>Palaemon northropi</i> (Rankin, 1898).....	14, 155, 169
84. <i>Palaemonetes intermedius</i> Holthuis, 1949.....	14, 155, 171
85. <i>Palaemonetes paludosus</i> (Gibbes, 1850).....	14, 155, 171
86. <i>Palaemonetes pugio</i> Holthuis, 1949.....	14, 155, 171
87. <i>Palaemonetes vulgaris</i> Say, 1818.....	15, 155, 171
88. <i>Periclimenaeus ascidiarum</i> Holthuis, 1951.....	15, 156, 173
89. <i>Periclimenaeus atlanticus</i> (Rathbun, 1901).....	15, 157, 177
90. <i>Periclimenaeus bermudensis</i> (Armstrong, 1940).....	15, 156, 175
91. <i>Periclimenaeus caraibicus</i> Holthuis, 1951.....	15, 156, 173
92. <i>Periclimenaeus chacei</i> Abele, 1971.....	15, 156, 173
93. <i>Periclimenaeus maxillulidens</i> (Schmitt, 1936).....	15, 157, 177
94. <i>Periclimenaeus pearsei</i> (Schmitt, 1932).....	15, 156, 173
95. <i>Periclimenaeus perlatus</i> (Boone, 1930).....	15, 156, 175
96. <i>Periclimenaeus schmitti</i> Holthuis, 1951.....	15, 157, 175
97. <i>Periclimenaeus wilsoni</i> (Hay, 1917).....	15, 156, 175
98. <i>Periclimenes americanus</i> (Kingsley, 1878).....	15, 158, 179
99. <i>Periclimenes harringtoni</i> Lebour, 1949.....	16, 159, 181
100. <i>Periclimenes iridescens</i> Lebour, 1949.....	16, 159, 181
101. <i>Periclimenes longicaudatus</i> (Stimpson, 1860).....	16, 158, 179
102. <i>Periclimenes magnus</i> Holthuis, 1951.....	16, 158, 179
103. <i>Periclimenes pandionis</i> Holthuis, 1951.....	16, 158, 181
104. <i>Periclimenes pedersoni</i> Chace, 1958.....	16, 158, 179
105. <i>Periclimenes perryae</i> Chace, 1942.....	16, 159, 183
106. <i>Periclimenes rathbunae</i> Schmitt, 1924.....	16, 159, 181
107. <i>Periclimenes yucatanicus</i> (Ives, 1891).....	16, 159, 183
108. <i>Pontonia domestica</i> Gibbes, 1850.....	16, 160, 185
109. <i>Pontonia margarita</i> Smith, 1869.....	16, 160, 185
110. <i>Pontonia unidens</i> Kingsley, 1880.....	16, 160, 185
111. <i>Pontiopsis paulae</i> Gore, 1981.....	16, 153, 193
112. <i>Pseudocoutierea antillensis</i> Chace, 1972.....	17, 153, 193
113. <i>Tuleariocaris neglecta</i> Chace, 1969.....	17, 152, 193
114. <i>Typton carneus</i> Holthuis, 1951.....	17, 161, 187
115. <i>Typton distinctus</i> Chace, 1972.....	17, 161, 189
116. <i>Typton gnathophylloides</i> Holthuis, 1951.....	17, 161, 187
117. <i>Typton prionurus</i> Holthuis, 1951.....	17, 161, 187
118. <i>Typton tortugae</i> McClendon, 1911.....	17, 161, 187
119. <i>Typton vulcanus</i> Holthuis, 1951.....	17, 161, 189
120. <i>Veleroniopsis kimallynae</i> Gore, 1981.....	17, 153, 193

Family Alpheidae

121. <i>Alpheopsis labis</i> Chace, 1972.....	17, 195, 205
122. <i>Alpheopsis trispinosus</i> (Stimpson, 1861).....	17, 195, 205
123. <i>Alpheus amblyonyx</i> Chace, 1972.....	17, 197, 209
124. <i>Alpheus armatus</i> Rathbun, 1901.....	17, 196, 207

125. <i>Alpheus armillatus</i> H. Milne Edwards, 1837.....	17, 198, 213
126. <i>Alpheus bouvieri</i> A. Milne Edwards, 1878.....	18, 199, 215
127. <i>Alpheus candei</i> Guérin-Méneville, 1855.....	18, 197, 209
128. <i>Alpheus cristulifrons</i> Rathbun, 1900.....	18, 198, 211
129. <i>Alpheus cylindricus</i> Kingsley, 1878.....	18, 196, 207
130. <i>Alpheus estuariensis</i> Christoffersen, 1984.....	18, 198, 211
131. <i>Alpheus floridanus</i> Kingsley, 1878.....	18, 199, 215
132. <i>Alpheus formosus</i> Gibbes, 1850.....	18, 197, 207
133. <i>Alpheus heterochaelis</i> Say, 1818.....	18, 198, 213
134. <i>Alpheus malleator</i> Dana, 1852.....	18, 197, 207
135. <i>Alpheus normanni</i> Kingsley, 1878.....	18, 198, 211
136. <i>Alpheus nuttingi</i> (Schmitt, 1924).....	18, 199, 215
137. <i>Alpheus paracrinitus</i> Miers, 1881.....	18, 199, 215
138. <i>Alpheus peasei</i> (Armstrong, 1940).....	19, 197, 211
139. <i>Alpheus schmitti</i> Chace, 1972.....	19, 199, 213
140. <i>Alpheus thomasi</i> Hendrix and Gore, 1973.....	19, 197, 209
141. <i>Alpheus viridari</i> (Armstrong, 1949).....	19, 198, 213
142. <i>Alpheus websteri</i> Kingsley, 1880.....	19, 197, 209
143. <i>Automate evermanni</i> Rathbun, 1901.....	19, 200, 217
144. <i>Automate gardineri</i> Coutière, 1902.....	19, 200, 217
145. <i>Automate rectifrons</i> Chace, 1972.....	19, 200, 217
146. <i>Leptalphes forceps</i> Williams, 1965.....	19, 194, 229
147. <i>Metalpheus rostratipes</i> (Pocock, 1890).....	19, 194, 229
148. <i>Synalphes agelas</i> Pequegnat and Heard, 1979.....	19, 201, 219
149. <i>Synalphes apioceros</i> Coutière, 1909.....	19, 203, 227
150. <i>Synalphes bousfieldi</i> Chace, 1972.....	20, 202, 221
151. <i>Synalphes brevicarpus</i> (Herrick, 1891).....	20, 203, 225
152. <i>Synalphes brooksi</i> Coutière, 1909.....	20, 202, 221
153. <i>Synalphes curacaoensis</i> Schmitt, 1924.....	20, 203, 225
154. <i>Synalphes fritzmuelleri</i> Coutière, 1909.....	20, 203, 227
155. <i>Synalphes goodei</i> Coutière, 1909.....	20, 202, 225
156. <i>Synalphes heardi</i> Dardeau, 1984.....	20, 201, 219
157. <i>Synalphes hemphilli</i> Coutière, 1909.....	20, 203, 227
158. <i>Synalphes herricki</i> Coutière, 1909.....	20, 202, 223
159. <i>Synalphes longicarpus</i> (Herrick, 1891).....	20, 202, 223
160. <i>Synalphes mcclendonii</i> Coutière, 1910.....	20, 202, 221
161. <i>Synalphes minus</i> (Say, 1818).....	20, 203, 225
162. <i>Synalphes pandionis</i> Coutière, 1909.....	21, 202, 223
163. <i>Synalphes paranepetunus</i> Coutière, 1909.....	21, 202, 223
164. <i>Synalphes pectiniger</i> Coutière, 1907.....	21, 201, 219
165. <i>Synalphes rathbunae</i> Coutière, 1909.....	21, 201, 219
166. <i>Synalphes sanctithomae</i> Coutière, 1909.....	21, 202, 221
167. <i>Synalphes townsendi</i> Coutière, 1909.....	21, 203, 227
168. <i>Thunor simus</i> (Guérin-Méneville, 1856).....	21, 194, 229

Family Hippolytidae

169. <i>Bythocaris nana</i> Smith, 1885.....	21, 230, 249
170. <i>Exhippolysmata oplophoroides</i> (Holthuis, 1948).....	21, 230, 249
171. <i>Hippolyte coerulescens</i> (Fabricius, 1775).....	21, 231, 237
172. <i>Hippolyte curacaoensis</i> Schmitt, 1924.....	21, 231, 237
173. <i>Hippolyte nicholsoni</i> Chace, 1972.....	21, 231, 237
174. <i>Hippolyte pleuracanthus</i> (Stimpson, 1871).....	21, 231, 237
175. <i>Hippolyte zostericola</i> (Smith, 1873).....	22, 231, 239
176. <i>Latreutes fucorum</i> (Fabricius, 1798).....	22, 232, 241
177. <i>Latreutes parvulus</i> (Stimpson, 1866).....	22, 232, 241
178. <i>Lysmata amboinensis</i> (De Man, 1888).....	22, 233, 243
179. <i>Lysmata intermedia</i> (Kingsley, 1878).....	22, 233, 243

180. <i>Lysmata rathbunae</i> Chace, 1970.....	22, 233, 243
181. <i>Lysmata wurdemanni</i> (Gibbes, 1850).....	22, 233, 243
182. <i>Merhippolyte americana</i> Holthuis, 1961.....	22, 230, 249
183. <i>Thor amboinensis</i> (De Man, 1888).....	22, 234, 245
184. <i>Thor dobkini</i> Chace, 1972.....	22, 234, 245
185. <i>Thor floridanus</i> Kingsley, 1878.....	22, 234, 245
186. <i>Thor manningi</i> Chace, 1972.....	22, 234, 245
187. <i>Tozeuma carolinense</i> Kingsley, 1878.....	22, 235, 247
188. <i>Tozeuma cornutum</i> A. Milne Edwards, 1881.....	22, 235, 247
189. <i>Tozeuma serratum</i> A. Milne Edwards, 1881.....	23, 235, 247
190. <i>Trachycaris restrictus</i> (A. Milne Edwards, 1878).....	23, 230, 249

Family Ogyrididae

191. <i>Ogyrides alphaerostris</i> (Kingsley, 1880).....	23, 251, 253
192. <i>Ogyrides hayi</i> Williams, 1981.....	23, 251, 253

Family Processidae

193. <i>Ambidexter symmetricus</i> Manning and Chace, 1971.....	23, 254, 261
194. <i>Nikoides schmitti</i> Manning and Chace, 1971.....	23, 254, 261
195. <i>Processa bermudensis</i> (Rankin, 1900).....	23, 255, 257
196. <i>Processa fimbriata</i> Manning and Chace, 1971.....	23, 255, 257
197. <i>Processa guyanae</i> Holthuis, 1959.....	23, 255, 259
198. <i>Processa hemphilli</i> Manning and Chace, 1971.....	23, 255, 259
199. <i>Processa profunda</i> Manning and Chace, 1971.....	23, 255, 259
200. <i>Processa riveroi</i> Manning and Chace, 1971.....	24, 255, 257
201. <i>Processa vicina</i> Manning and Chace, 1971.....	24, 255, 257

Family Pandalidae

202. <i>Pantomus parvulus</i> A. Milne Edwards, 1883.....	24, 262, 269
203. <i>Plesionika acanthonotus</i> (Smith, 1882).....	24, 263, 265
204. <i>Plesionika edwardsii</i> (Brandt, 1851).....	24, 263, 265
205. <i>Plesionika ensis</i> (A. Milne Edwards, 1881).....	24, 263, 265
206. <i>Plesionika escatilis</i> (Stimpson, 1860).....	24, 263, 267
207. <i>Plesionika longicauda</i> (Rathbun, 1901).....	24, 263, 267
208. <i>Plesionika martia</i> (A. Milne Edwards, 1883).....	24, 263, 267
209. <i>Plesionika tenuipes</i> (Smith, 1881).....	24, 263, 265
210. <i>Stylopandalus richardi</i> (Coutière, 1905).....	24, 262, 269

Family Crangonidae

211. <i>Crangon septemspinosa</i> Say, 1818.....	25, 271, 273
212. <i>Metacrangon jacqueti agassizii</i> (Smith, 1882).....	25, 271, 273
213. <i>Parapontocaris caribbaea</i> (Boone, 1927).....	25, 271, 273
214. <i>Philoceras gorei</i> (Dardeau, 1980).....	25, 271, 275
215. <i>Pontophilus brevisrostris</i> Smith, 1881.....	25, 271, 275

Family Glyphocrangonidae

216. <i>Glyphocrangon haematonotus</i> Holthuis, 1971.....	25, 277, 279
217. <i>Glyphocrangon longleyi</i> Schmitt, 1931.....	25, 277, 279
218. <i>Glyphocrangon spinicauda</i> A. Milne Edwards, 1881.....	25, 277, 279

Infraorder Astacidea

Family Nephropidae

219. <i>Acanthacaris caeca</i> (A. Milne Edwards, 1881).....	25, 285, 287
220. <i>Metanephrops binghami</i> (Boone, 1927).....	26, 285, 287
221. <i>Nephropsis aculeata</i> Smith, 1881.....	26, 285, 287

Infraorder Thalassinidea**Family Axiidae**

222. *Axiopsis hirsutimana* (Boesch and Smalley, 1972)..... 26, 289, 291
 223. *Axiopsis oxypleura* (Williams, 1974)..... 26, 289, 291
 224. *Axiopsis serratifrons* (A. Milne Edwards, 1873)..... 26, 289, 291
 225. *Coralaxius abelei* Kensley and Gore, 1981..... 26, 289, 291

Family Callianassidae

226. *Callianassa acanthochirus* (Stimpson, 1866)..... 26, 294, 296, 299
 227. *Callianassa atlantica* Rathbun, 1926..... 26, 294, 297, 301
 228. *Callianassa biformis* Biffar, 1971..... 26, 295, 305
 229. *Callianassa branneri* (Rathbun, 1900)..... 26, 295, 297, 303
 230. *Callianassa fragilis* Biffar, 1970..... 26, 294, 297, 301
 231. *Callianassa guassutina* Rodrigues, 1966..... 26, 294, 296, 299
 232. *Callianassa jamaicense* Schmitt, 1935..... 27, 295, 296, 303
 233. *Callianassa longiventris* A. Milne Edwards, 1870..... 27, 294, 296, 299
 234. *Callianassa marginata* Rathbun, 1901..... 27, 295, 296, 305
 235. *Callianassa quadracuta* Biffar, 1970..... 27, 295, 303
 236. *Callianassa rathbunae* Schmitt, 1935..... 27, 294, 296, 299
 237. *Callianassa trilobata* Biffar, 1970..... 27, 295, 297, 303
 238. *Callichirus islagrande* (Schmitt, 1935)..... 27, 295, 296, 307
 239. *Callichirus major* (Say, 1818)..... 27, 295, 296, 297, 307
 240. *Gourretia latispina* (Dawson, 1967)..... 27, 293, 296, 307

Family Upogebiidae

241. *Upogebia affinis* (Say, 1818)..... 27, 309, 311
 242. *Upogebia operculata* Schmitt, 1924..... 27, 309, 311

Infraorder Palinura**Family Palinuridae**

243. *Justitia longimanus* (H. Milne Edwards, 1837)..... 27, 313, 315
 244. *Panulirus argus* (Latreille, 1804)..... 27, 313, 315
 245. *Panulirus guttatus* (Latreille, 1804)..... 28, 313, 315
 246. *Panulirus laevicauda* (Latreille, 1817)..... 28, 313, 315

Family Scyllaridae

247. *Parribacus antarcticus* (Lund, 1793)..... 28, 316, 321
 248. *Scyllarides aequinoctialis* (Lund, 1793)..... 28, 317, 319
 249. *Scyllarides nodifer* (Stimpson, 1866)..... 28, 317, 319
 250. *Scyllarus americanus* (Smith, 1869)..... 28, 317, 321
 251. *Scyllarus chacei* Holthuis, 1960..... 28, 317, 321
 252. *Scyllarus depressus* (Smith, 1881)..... 28, 317, 321

Family Synaxidae

253. *Palinurellus gundlachi* (Von Martens, 1881)..... 28, 323, 325

Infraorder Anomura**Family Coenobitidae**

254. *Coenobita clypeatus* (Herbst, 1791)..... 28, 327, 329

Family Diogenidae

255. *Calcinus tibicen* (Herbst, 1791)..... 28, 330, 353
 256. *Cancellus ornatus* Benedict, 1901..... 28, 331, 337
 257. *Cancellus viridis* Mayo, 1973..... 29, 331, 337

258. <i>Clibanarius antillensis</i> Stimpson, 1862.....	29, 331, 339
259. <i>Clibanarius cubensis</i> (Saussure, 1858).....	29, 331, 339
260. <i>Clibanarius tricolor</i> (Gibbes, 1850).....	29, 331, 339
261. <i>Clibanarius vittatus</i> (Bosc, 1802).....	29, 331, 339
262. <i>Dardanus fucosus</i> Biffar and Provenzano, 1972.....	29, 332, 341
263. <i>Dardanus insignis</i> (Saussure, 1858).....	29, 332, 341
264. <i>Dardanus venosus</i> (H. Milne Edwards, 1848).....	29, 332, 341
265. <i>Isocheles wurdemanni</i> Stimpson, 1862.....	29, 330, 353
266. <i>Paguristes anomalus</i> Bouvier, 1918.....	29, 334, 349
267. <i>Paguristes cadenati</i> Forest, 1954.....	29, 334, 349
268. <i>Paguristes erythrops</i> Holthuis, 1959.....	29, 334, 347
269. <i>Paguristes grayi</i> Benedict, 1901.....	29, 334, 345
270. <i>Paguristes hernancortezii</i> McLaughlin and Provenzano, 1974.....	29, 334, 349
271. <i>Paguristes hummi</i> Wass, 1955.....	30, 333, 343
272. <i>Paguristes inconstans</i> McLaughlin and Provenzano, 1974.....	30, 333, 345
273. <i>Paguristes invisissacculus</i> McLaughlin and Provenzano, 1974.....	30, 335, 353
274. <i>Paguristes laticlavus</i> McLaughlin and Provenzano, 1974.....	30, 333, 343
275. <i>Paguristes limonensis</i> McLaughlin and Provenzano, 1974.....	30, 334, 351
276. <i>Paguristes lymani</i> A. Milne Edwards and Bouvier, 1893.....	30, 333, 343
277. <i>Paguristes moorei</i> Benedict, 1901.....	30, 333, 345
278. <i>Paguristes oxyophthalmus</i> Holthuis, 1959.....	30, 333, 343
279. <i>Paguristes puncticeps</i> Benedict, 1901.....	30, 334, 347
280. <i>Paguristes sericeus</i> A. Milne Edwards, 1880.....	30, 334, 347
281. <i>Paguristes spinipes</i> A. Milne Edwards, 1880.....	30, 334, 347
282. <i>Paguristes starcki</i> Provenzano, 1965.....	30, 334, 351
283. <i>Paguristes tenuirostris</i> Benedict, 1901.....	30, 334, 349
284. <i>Paguristes tortugae</i> Schmitt, 1933.....	31, 335, 351
285. <i>Paguristes triangulatus</i> A. Milne Edwards and Bouvier, 1893.....	31, 333, 345
286. <i>Paguristes wassi</i> Provenzano, 1961.....	31, 334, 351
287. <i>Petrochirus diogenes</i> (Linnaeus, 1758).....	31, 330, 353

Family Lithodidae

288. <i>Paralomis cubensis</i> Chace, 1939.....	31, 355, 357
---	--------------

Family Paguridae

289. <i>Agaricochirus acanthinus</i> McLaughlin, 1982.....	31, 361, 367
290. <i>Agaricochirus alexandri</i> (A. Milne Edwards and Bouvier, 1893).....	31, 361, 367
291. <i>Agaricochirus boletifer</i> (A. Milne Edwards and Bouvier, 1893).....	31, 361, 367
292. <i>Agaricochirus gibbosimanus</i> (A. Milne Edwards, 1880).....	31, 361, 367
293. <i>Anisopagurus bartletti</i> (A. Milne Edwards, 1880).....	31, 361, 369
294. <i>Anisopagurus pygmaeus</i> (Bouvier, 1918).....	31, 361, 369
295. <i>Catapagurus sharrei</i> A. Milne Edwards, 1880.....	32, 359, 387
296. <i>Iridopagurus caribbensis</i> (A. Milne Edwards and Bouvier, 1893).....	32, 362, 371
297. <i>Iridopagurus globulus</i> De Saint Laurent-Dechancé, 1966.....	32, 362, 371
298. <i>Iridopagurus iris</i> (A. Milne Edwards, 1880).....	32, 362, 371
299. <i>Iridopagurus reticulatus</i> García Gómez, 1983.....	32, 362, 371
300. <i>Iridopagurus violaceus</i> De Saint Laurent-Dechancé, 1966.....	32, 362, 373
301. <i>Manucomplanus corallinus</i> (Benedict, 1892).....	32, 360, 387
302. <i>Nematopaguroides pusillus</i> Forest and De Saint Laurent, 1967.....	32, 359, 387
303. <i>Ostraconotus spatulipes</i> A. Milne Edwards, 1880.....	32, 359, 389
304. <i>Pagurus annulipes</i> (Stimpson, 1860).....	32, 363, 379
305. <i>Pagurus brevidactylus</i> (Stimpson, 1859).....	33, 363, 375
306. <i>Pagurus carolinensis</i> McLaughlin, 1975.....	33, 363, 375
307. <i>Pagurus criniticornis</i> (Dana, 1852).....	33, 363, 379
308. <i>Pagurus defensus</i> (Benedict, 1892).....	33, 364, 381
309. <i>Pagurus gymnodactylus</i> Lemaitre, 1982.....	33, 363, 377
310. <i>Pagurus impressus</i> (Benedict, 1892).....	33, 363, 377
311. <i>Pagurus longicarpus</i> Say, 1817.....	33, 364, 381

312. <i>Pagurus macLaughlinae</i> García Gómez, 1982.....	33, 363, 377
313. <i>Pagurus marshi</i> Benedict, 1901.....	33, 363, 377
314. <i>Pagurus piercei</i> Wass, 1963.....	33, 364, 381
315. <i>Pagurus politus</i> (Smith, 1882).....	33, 364, 379
316. <i>Pagurus pollicaris</i> Say, 1817.....	33, 363, 375
317. <i>Pagurus provenzanoi</i> Forest and De Saint Laurent, 1967.....	34, 362, 375
318. <i>Pagurus stimpsoni</i> (A. Milne Edwards and Bouvier, 1893).....	34, 364, 379
319. <i>Phimochirus holthuisi</i> (Provenzano, 1961).....	34, 365, 383
320. <i>Phimochirus leurocarpus</i> McLaughlin, 1981.....	34, 365, 383
321. <i>Phimochirus operculatus</i> (Stimpson, 1859).....	34, 365, 383
322. <i>Phimochirus randalli</i> (Provenzano, 1961).....	34, 365, 383
323. <i>Pylopaguropsis atlantica</i> Wass, 1963.....	34, 359, 389
324. <i>Pylopagurus discoidalis</i> (A. Milne Edwards, 1880).....	34, 359, 389
325. <i>Rhodochirus rosaceus</i> (A. Milne Edwards and Bouvier, 1893).....	34, 360, 389
326. <i>Solenopagurus lineatus</i> (Wass, 1963).....	34, 359, 391
327. <i>Tomopaguropsis problematica</i> (A. Milne Edwards and Bouvier, 1893).....	34, 359, 391
328. <i>Tomopagurus chacei</i> (Wass, 1963).....	34, 365, 387
329. <i>Tomopagurus cokeri</i> (Hay, 1917).....	34, 365, 385
330. <i>Tomopagurus cubensis</i> (Wass, 1963).....	35, 365, 385
331. <i>Tomopagurus rubropunctatus</i> A. Milne Edwards and Bouvier, 1893.....	35, 365, 385
332. <i>Tomopagurus wassi</i> McLaughlin, 1981.....	35, 365, 385

Family Chirostylidae

333. <i>Uroptychus armatus</i> (A. Milne Edwards, 1880).....	35, 393, 395
--	--------------

Family Galatheididae

334. <i>Galathea rostrata</i> A. Milne Edwards, 1880.....	35, 397, 409
335. <i>Munida angulata</i> Benedict, 1902.....	35, 398, 405
336. <i>Munida affinis</i> A. Milne Edwards, 1880.....	35, 398, 405
337. <i>Munida forceps</i> A. Milne Edwards, 1880.....	35, 398, 403
338. <i>Munida iris iris</i> A. Milne Edwards, 1880.....	35, 398, 403
339. <i>Munida irrasa</i> A. Milne Edwards, 1880.....	35, 398, 403
340. <i>Munida longipes</i> A. Milne Edwards, 1880.....	35, 398, 405
341. <i>Munida miles</i> A. Milne Edwards, 1880.....	36, 397, 401
342. <i>Munida pusilla</i> Benedict, 1902.....	36, 398, 403
343. <i>Munida santipauli</i> Henderson, 1885.....	36, 397, 401
344. <i>Munida simplex</i> Benedict, 1902.....	36, 398, 405
345. <i>Munida spinifrons</i> Henderson, 1885.....	36, 397, 401
346. <i>Munida stimpsoni</i> A. Milne Edwards, 1880.....	36, 398, 407
347. <i>Munida valida</i> Smith, 1883.....	36, 398, 401
348. <i>Munidopsis armata</i> (A. Milne Edwards, 1880).....	36, 399, 407
349. <i>Munidopsis platirostris</i> (A. Milne Edwards and Bouvier, 1894).....	36, 399, 407
350. <i>Munidopsis polita</i> (Smith, 1883).....	36, 399, 407

Family Porcellanidae

351. <i>Euceramus praelongus</i> Stimpson, 1860.....	36, 410, 425
352. <i>Megalobrachium poeyi</i> (Guérin-Méneville, 1855).....	37, 412, 415
353. <i>Megalobrachium soriatum</i> (Say, 1818).....	37, 412, 415
354. <i>Neopisosoma angustifrons</i> (Benedict, 1901).....	37, 410, 425
355. <i>Pachycheles ackleianus</i> A. Milne Edwards, 1880.....	37, 412, 417
356. <i>Pachycheles monilifer</i> (Dana, 1852).....	37, 412, 419
357. <i>Pachycheles pilosus</i> (H. Milne Edwards, 1837).....	37, 412, 417
358. <i>Pachycheles riisei</i> (Stimpson, 1858).....	37, 412, 417
359. <i>Pachycheles rugimanus</i> A. Milne Edwards, 1880.....	37, 412, 417
360. <i>Parapetrolisthes tortugensis</i> (Glassell, 1945).....	37, 410, 425
361. <i>Petrolisthes armatus</i> (Gibbes, 1850).....	37, 413, 421
362. <i>Petrolisthes galathinus</i> (Bosc, 1802).....	37, 413, 421
363. <i>Petrolisthes jugosus</i> Streets, 1872.....	37, 413, 421

364. *Petrolisthes politus* (Gray, 1831)..... 38, 413, 421
 365. *Polyonyx gibbesi* Haig, 1956..... 38, 410, 425
 366. *Porcellana sayana* (Leach, 1820)..... 38, 413, 423
 367. *Porcellana sigsbeiana* A. Milne Edwards, 1880..... 38, 413, 423
 368. *Porcellana stimpsoni* A. Milne Edwards, 1880..... 38, 413, 423

Family Albuneidae

369. *Albunea gibbesii* Stimpson, 1859..... 38, 427, 429
 370. *Albunea paretii* Guérin-Méneville, 1853..... 38, 427, 429
 371. *Lepidopa benedicti* Schmitt, 1935..... 38, 427, 429
 372. *Lepidopa websteri* Benedict, 1903..... 38, 427, 429
 373. *Zygopa michaelis* Holthuis, 1960..... 38, 427, 431

Family Hippidae

374. *Emerita benedicti* Schmitt, 1935..... 38, 433, 435
 375. *Emerita portoricensis* Schmitt, 1935..... 38, 433, 435
 376. *Emerita talpoida* (Say, 1817)..... 38, 433, 435
 377. *Hippa cubensis* (Saussure, 1857)..... 38, 433, 435

Infraorder Brachyura

Family Dromiidae

378. *Dromia erythropus* (George Edwards, 1771)..... 39, 437, 441
 379. *Dromidia antillensis* Stimpson, 1858..... 39, 437, 441
 380. *Hypoconcha arcuata* Stimpson, 1858..... 39, 437, 439
 381. *Hypoconcha sabulosa* (Herbst, 1799)..... 39, 437, 439
 382. *Hypoconcha spinosissima* Rathbun, 1933..... 39, 437, 439

Family Homolodromiidae

383. *Dicranodromia ovata* A. Milne Edwards, 1880..... 39, 443, 445

Family Cymonomidae

384. *Cymonomus quadratus* A. Milne Edwards, 1880..... 39, 443, 445
 385. *Cymopolus agassizi* A. Milne Edwards and Bouvier, 1899..... 39, 443, 445

Family Cyclodorippidae

386. *Clythrocerus granulatus* (Rathbun, 1898)..... 39, 447, 449
 387. *Clythrocerus nitidus* (A. Milne Edwards, 1880)..... 39, 447, 449
 388. *Clythrocerus stimpsoni* Rathbun, 1937..... 39, 447, 449
 389. *Tymolus antennaria* (A. Milne Edwards, 1880)..... 39, 447, 449

Family Homolidae

390. *Homola barbata* (Fabricius, 1793)..... 40, 451, 453

Family Latreilliidae

391. *Latreillia manningi* Williams, 1982..... 40, 451, 453

Family Raninidae

392. *Lyreidus nitidus* (A. Milne Edwards, 1880)..... 40, 455, 459
 393. *Ranilia constricta* (A. Milne Edwards, 1880)..... 40, 455, 457
 394. *Ranilia muricata* H. Milne Edwards, 1837..... 40, 455, 457
 395. *Raninoides loevis* (Latreille, 1825)..... 40, 455, 457
 396. *Raninoides louisianensis* Rathbun, 1933..... 40, 455, 457
 397. *Symethis variolosa* (Fabricius, 1793)..... 40, 455, 459

Family Dorippidae

398. *Ethusa mascarone americana* A. Milne Edwards, 1880 40, 461, 463
 399. *Ethusa microphthalmia* Smith, 1881..... 40, 461, 463
 400. *Ethusa tenuipes* Rathbun, 1897..... 41, 461, 463
 401. *Ethusa truncata* A. Milne Edwards and Bouvier, 1899..... 41, 461, 463

Family Calappidae

402. *Acanthocarpus alexandri* Stimpson, 1871..... 41, 465, 469
 403. *Acanthocarpus bispinosus* A. Milne Edwards, 1880..... 41, 465, 469
 404. *Calappa angusta* A. Milne Edwards, 1880..... 41, 466, 471
 405. *Calappa flammea* (Herbst, 1794)..... 42, 466, 471
 406. *Calappa gallus* (Herbst, 1803)..... 41, 466, 471
 407. *Calappa ocellata* Holthuis, 1958..... 41, 466, 473
 408. *Calappa sulcata* Rathbun, 1898..... 41, 466, 471
 409. *Cycloes bairdii* Stimpson, 1860..... 41, 465, 477
 410. *Hepatus epheliticus* (Linnaeus, 1763)..... 41, 466, 475
 411. *Hepatus pudibundus* (Herbst, 1785)..... 41, 466, 475
 412. *Osachila antillensis* Rathbun, 1898..... 42, 467, 477
 413. *Osachila semilevis* Rathbun, 1916..... 42, 467, 477
 414. *Osachila tuberosa* Stimpson, 1871..... 42, 467, 477

Family Leucosiidae

415. *Callidactylus asper* Stimpson, 1871..... 42, 479, 491
 416. *Ebalia cariosa* (Stimpson, 1860)..... 42, 480, 483
 417. *Ebalia stimpsonii* A. Milne Edwards, 1880..... 42, 480, 483
 418. *Iliacantha intermedia* Miers, 1886..... 42, 480, 485
 419. *Iliacantha liodactylus* Rathbun, 1898..... 42, 480, 485
 420. *Iliacantha sparsa* Stimpson, 1871..... 42, 480, 485
 421. *Iliacantha subglobosa* Stimpson, 1871..... 42, 480, 485
 422. *Lithadia cadaverosa* Stimpson, 1871..... 42, 480, 487
 423. *Lithadia granulosa* A. Milne Edwards, 1880..... 42, 480, 487
 424. *Myropsis quinquespinosa* Stimpson, 1871..... 42, 479, 491
 425. *Persephona crinita* Rathbun, 1931..... 42, 481, 487
 426. *Persephona mediterranea* (Herbst, 1794)..... 43, 481, 487
 427. *Speloeophorus elevatus* Rathbun, 1898..... 43, 481, 489
 428. *Speloeophorus nodosus* (Bell, 1855)..... 43, 481, 489
 429. *Speloeophorus pontifer* (Stimpson, 1871)..... 43, 481, 489
 430. *Uhlias limbatus* Stimpson, 1871..... 43, 479, 491

Family Majidae

431. *Acanthonyx petiverii* H. Milne Edwards, 1834..... 43, 495, 545
 432. *Achaeopsis thomsoni* (Norman, 1873)..... 43, 493, 545
 433. *Aepinus septemspinosus* (A. Milne Edwards, 1879)..... 43, 493, 547
 434. *Anasimus latus* Rathbun, 1894..... 43, 494, 547
 435. *Anomalothir furcillatus* (Stimpson, 1871)..... 43, 493, 547
 436. *Arachnopsis filipes* Stimpson, 1871..... 43, 493, 547
 437. *Batrachonotus fragosus* Stimpson, 1871..... 43, 493, 549
 438. *Chorinus heros* (Herbst, 1790)..... 44, 494, 549
 439. *Coelocerus spinosus* A. Milne Edwards, 1875..... 44, 496, 549
 440. *Collodes leptocheles* Rathbun, 1894..... 44, 497, 509
 441. *Collodes nudus* Stimpson, 1871..... 44, 497
 442. *Collodes obesus* A. Milne Edwards, 1878..... 44, 497, 509
 443. *Collodes robustus* Smith, 1883..... 44, 497, 509
 444. *Collodes trispinosus* Stimpson, 1871..... 44, 497, 509
 445. *Epialtus bituberculatus* H. Milne Edwards, 1834..... 44, 498, 511
 446. *Epialtus dilatatus* A. Milne Edwards, 1878..... 44, 498, 511
 447. *Epialtus dilatatus* forma *elongata* Rathbun, 1923..... 44, 498, 513

448. <i>Epialtus kingsleyi</i> Rathbun, 1923.....	44, 498, 511
449. <i>Epialtus longirostris</i> Stimpson, 1860.....	44, 498, 511
450. <i>Euprognatha gracilipes</i> A. Milne Edwards, 1878.....	44, 499, 513
451. <i>Euprognatha rastellifera</i> Stimpson, 1871.....	45, 499, 513
452. <i>Hemus cristulipes</i> A. Milne Edwards, 1875.....	45, 495, 549
453. <i>Inachoides forceps</i> A. Milne Edwards, 1879.....	45, 494, 551
454. <i>Leptopisa setirostris</i> (Stimpson, 1871).....	45, 496, 551
455. <i>Libinia dubia</i> H. Milne Edwards, 1834.....	45, 499, 515
456. <i>Libinia emarginata</i> Leach, 1815.....	45, 499, 515
457. <i>Libinia erinacea</i> (A. Milne Edwards, 1879).....	45, 499, 515
458. <i>Macrocoeloma camptocerum</i> (Stimpson, 1871).....	45, 500, 519
459. <i>Macrocoeloma diplacanthum</i> (Stimpson, 1860).....	45, 500, 519
460. <i>Macrocoeloma eutheca</i> (Stimpson, 1871).....	45, 500, 519
461. <i>Macrocoeloma laevigatum</i> (Stimpson, 1860).....	45, 500, 519
462. <i>Macrocoeloma septemspinum</i> (Stimpson, 1871).....	45, 500, 521
463. <i>Macrocoeloma subparallelum</i> (Stimpson, 1860).....	46, 500, 517
464. <i>Macrocoeloma trispinosum trispinosum</i> (Latreille, 1825).....	46, 500, 517
465. <i>Macrocoeloma trispinosum nodipes</i> (Desbonne, 1867).....	46, 500, 517
466. <i>Macrocoeloma trispinosum</i> , variety.....	46, 500, 517
467. <i>Metoporphaphis calcarata</i> (Say, 1818).....	46, 494, 551
468. <i>Microphrys antillensis</i> Rathbun, 1920.....	46, 501, 521
469. <i>Microphrys bicornutus</i> (Latreille, 1825).....	46, 501, 521
470. <i>Mithrax acuticornis</i> Stimpson, 1870.....	46, 501, 523
471. <i>Mithrax caribbaeus</i> Rathbun, 1900.....	46, 502, 527
472. <i>Mithrax cinctimanus</i> (Stimpson, 1860).....	46, 502, 527
473. <i>Mithrax cornutus</i> Saussure, 1857.....	46, 501, 523
474. <i>Mithrax coryphe</i> (Herbst, 1801).....	46, 503, 529
475. <i>Mithrax forceps</i> (A. Milne Edwards, 1875).....	47, 503, 529
476. <i>Mithrax hemphilli</i> Rathbun, 1892.....	47, 502, 525
477. <i>Mithrax hispidus</i> (Herbst, 1790).....	47, 502, 525
478. <i>Mithrax holderi</i> Stimpson, 1871.....	47, 502, 525
479. <i>Mithrax pilosus</i> Rathbun, 1892.....	47, 501, 523
480. <i>Mithrax pleuracanthus</i> Stimpson, 1871.....	47, 502, 527
481. <i>Mithrax ruber</i> (Stimpson, 1871).....	47, 503, 529
482. <i>Mithrax sculptus</i> (Lamarck, 1818).....	47, 503, 529
483. <i>Mithrax spinosissimus</i> (Lamarck, 1818).....	47, 501, 523
484. <i>Mithrax tortugae</i> Rathbun, 1920.....	47, 502, 527
485. <i>Mithrax verrucosus</i> H. Milne Edwards, 1832.....	47, 501, 502, 525
486. <i>Mocosoia crebripunctata</i> Stimpson, 1871.....	47, 495, 551
487. <i>Nibilia antilocapra</i> (Stimpson, 1871).....	47, 494, 553
488. <i>Oplopisa spinipes</i> A. Milne Edwards, 1879.....	48, 494, 553
489. <i>Pelia mutica</i> (Gibbes, 1850).....	48, 494, 553
490. <i>Picroceroides tubularis</i> Miers, 1886.....	48, 495, 553
491. <i>Pitho aculeata</i> (Gibbes, 1850).....	48, 504, 531
492. <i>Pitho anisodon</i> (Von Martens, 1872).....	48, 504, 531
493. <i>Pitho laevigata</i> (A. Milne Edwards, 1875).....	48, 504, 531
494. <i>Pitho lherminieri</i> (Schramm, 1867).....	48, 504, 531
495. <i>Pitho mirabilis</i> (Herbst, 1794).....	48, 504, 533
496. <i>Pitho quadridentata</i> (Miers, 1879).....	48, 504, 533
497. <i>Podochela curvirostris</i> (A. Milne Edwards, 1879).....	48, 505, 535
498. <i>Podochela gracilipes</i> Stimpson, 1871.....	48, 505, 535
499. <i>Podochela lamelligera</i> (Stimpson, 1871).....	48, 505, 535
500. <i>Podochela macrodera</i> Stimpson, 1860.....	49, 505, 535
501. <i>Podochela riisei</i> Stimpson, 1860.....	49, 505, 537
502. <i>Podochela sidneyi</i> Rathbun, 1924.....	49, 505, 537
503. <i>Pyromaia arachna</i> Rathbun, 1924.....	49, 506, 539
504. <i>Pyromaia cuspidata</i> Stimpson, 1871.....	49, 506, 539

505. <i>Rochinia crassa</i> (A. Milne Edwards, 1879).....	49, 506, 541
506. <i>Rochinia hystrix</i> (Stimpson, 1871).....	49, 506, 541
507. <i>Rochinia tanneri</i> (Smith, 1883).....	49, 506, 541
508. <i>Rochinia umbonata</i> (Stimpson, 1871).....	49, 506, 541
509. <i>Sphenocarcinus corrosus</i> A. Milne Edwards, 1875.....	49, 495, 555
510. <i>Stenocionops furcata coelata</i> (A. Milne Edwards, 1878).....	49, 507, 543
511. <i>Stenocionops furcata furcata</i> (Olivier, 1791).....	49, 507, 543
512. <i>Stenocionops spinimana</i> (Rathbun, 1892).....	50, 507, 543
513. <i>Stenocionops spinosissima</i> (Saussure, 1857).....	50, 507, 543
514. <i>Stenorhynchus seticornis</i> (Herbst, 1788).....	50, 494, 555
515. <i>Stilbomastax margaritifera</i> (Monod, 1939).....	50, 495, 555
516. <i>Thoe puella</i> Stimpson, 1860.....	50, 495, 555
517. <i>Tyche emarginata</i> White, 1847.....	50, 495, 557

Family Parthenopidae

518. <i>Cryptopodia concava</i> Stimpson, 1871.....	50, 558, 565
519. <i>Heterocrypta granulata</i> (Gibbes, 1850).....	50, 558, 565
520. <i>Leiolambrus nitidus</i> Rathbun, 1901.....	50, 558, 565
521. <i>Mesorhoea sexspinoso</i> Stimpson, 1871.....	50, 558, 565
522. <i>Parthenope agona</i> (Stimpson, 1871).....	50, 559, 561
523. <i>Parthenope fraterculus</i> (Stimpson, 1871).....	51, 559, 563
524. <i>Parthenope granulata</i> (Kingsley, 1879).....	51, 559, 561
525. <i>Parthenope pourtalesii</i> (Stimpson, 1871).....	51, 559, 561
526. <i>Parthenope serrata</i> (H. Milne Edwards, 1834).....	51, 559, 561
527. <i>Solenolambrus decemspinosus</i> Rathbun, 1894.....	51, 559, 563
528. <i>Solenolambrus typicus</i> Stimpson, 1871.....	51, 559, 563
529. <i>Solenolambrus tenellus</i> Stimpson, 1871.....	51, 559, 563
530. <i>Tutankhamen cristatipes</i> (A. Milne Edwards, 1880).....	51, 558, 567

Family Atelecyclidae

531. <i>Trichopeltarion nobile</i> A. Milne Edwards, 1880.....	51, 569, 571
--	--------------

Family Cancridae

532. <i>Cancer borealis</i> Stimpson, 1859.....	51, 569, 571
533. <i>Cancer irroratus</i> Say, 1817.....	52, 569, 571

Family Geryonidae

534. <i>Geryon fenneri</i> Manning and Hothuis, 1984.....	52, 569, 571
---	--------------

Family Portunidae

535. <i>Arenaeus cribrarius</i> (Lamarck, 1818).....	52, 572, 589
536. <i>Bathynectes longispina</i> Stimpson, 1871.....	52, 572, 589
537. <i>Benthochason schmitti</i> Rathbun, 1931.....	52, 572, 589
538. <i>Callinectes bocourti</i> A. Milne Edwards, 1879.....	52, 573, 577
539. <i>Callinectes danae</i> Smith, 1869.....	52, 573, 579
540. <i>Callinectes exasperatus</i> (Gerstaecker, 1856).....	52, 573, 579
541. <i>Callinectes larvatus</i> Ordway, 1863.....	52, 573, 577
542. <i>Callinectes ornatus</i> Ordway, 1863.....	52, 573, 579
543. <i>Callinectes sapidus</i> Rathbun, 1896.....	52, 573, 577
544. <i>Callinectes similis</i> Williams, 1966.....	53, 573, 577
545. <i>Cronius ruber</i> (Lamarck, 1818).....	53, 573, 581
546. <i>Cronius tumidulus</i> (Stimpson, 1871).....	53, 573, 581
547. <i>Ovalipes floridanus</i> Hay and Shore, 1918.....	53, 574, 581
548. <i>Ovalipes stephensoni</i> Williams, 1976.....	53, 574, 581
549. <i>Portunus anceps</i> (Saussure, 1858).....	53, 574, 583
550. <i>Portunus binoculus</i> Holthuis, 1969.....	53, 575, 587
551. <i>Portunus depressifrons</i> (Stimpson, 1859).....	53, 575, 585
552. <i>Portunus floridanus</i> Rathbun, 1930.....	53, 575, 585
553. <i>Portunus gibbesii</i> (Stimpson, 1859).....	53, 574, 583

554. <i>Portunus ordwayi</i> (Stimpson, 1860).....	53, 575, 587
555. <i>Portunus sayi</i> (Gibbes, 1850).....	53, 574, 583
556. <i>Portunus sebae</i> (H. Milne Edwards, 1834).....	54, 575, 585
557. <i>Portunus spinicarpus</i> (Stimpson, 1871).....	54, 575, 587
558. <i>Portunus spinimanus</i> Latreille, 1819.....	54, 575, 587
559. <i>Portunus ventralis</i> (A. Milne Edwards, 1879).....	54, 574, 585
560. <i>Portunus vocans</i> (A. Milne Edwards, 1878).....	54, 574, 583

Family Goneplacidae

561. <i>Chacellus filiformis</i> Guinot, 1969.....	54, 591, 595
562. <i>Eucratopsis crassimanus</i> (Dana, 1852).....	54, 592, 595
563. <i>Euphrosynoplax clausa</i> Guinot, 1969.....	54, 591, 597
564. <i>Euryplax nitida</i> Stimpson, 1859.....	54, 592, 597
565. <i>Frevillea barbata</i> A. Milne Edwards, 1880.....	54, 593, 595
566. <i>Frevillea hirsuta</i> (Borradaile, 1916).....	54, 593, 595
567. <i>Glyptoplax smithii</i> A. Milne Edwards, 1880.....	54, 592, 597
568. <i>Goneplax sigsbei</i> (A. Milne Edwards, 1880).....	54, 591, 597
569. <i>Nanoplax xanthiformis</i> (A. Milne Edwards, 1881).....	55, 591, 599
570. <i>Neopilumnoplax americana</i> (Rathbun, 1898).....	55, 591, 599
571. <i>Panoplax depressa</i> Stimpson, 1871.....	55, 592, 599
572. <i>Pilumnoplax elata</i> (A. Milne Edwards, 1880).....	55, 591
573. <i>Pseudorhombila quadridentata</i> (Latreille, 1828).....	55, 592, 599
574. <i>Sotoplax robertsi</i> Guinot, 1984.....	55, 592, 601
575. <i>Speocarcinus lobatus</i> Guinot, 1969.....	55, 592, 601
576. <i>Thalassoplax angusta</i> Guinot, 1969.....	55, 591, 601
577. <i>Trapezioplax tridentata</i> (A. Milne Edwards, 1880).....	55, 592, 601

Family Xanthidae

578. <i>Actaea acantha</i> (H. Milne Edwards, 1834).....	55, 607, 615
579. <i>Actaea bifrons</i> Rathbun, 1898.....	55, 607, 615
580. <i>Allactaea lithostrota</i> Williams, 1974.....	55, 604, 649
581. <i>Banareia palmeri</i> (Rathbun, 1894).....	56, 604, 649
582. <i>Carpilius corallinus</i> (Herbst, 1783).....	56, 603, 651
583. <i>Carpoporus papulosus</i> Stimpson, 1871.....	56, 604, 651
584. <i>Cataleptodius floridanus</i> (Gibbes, 1850).....	56, 604, 651
585. <i>Chlorodiella longimana</i> (H. Milne Edwards, 1834).....	56, 605, 653
586. <i>Domecia acanthophora acanthophora</i> (Desbonne and Schramm, 1867).....	56, 606, 653
587. <i>Eriphia gonagra</i> (Fabricius, 1781).....	56, 606, 653
588. <i>Etisus maculatus</i> (Stimpson, 1860).....	56, 605, 653
589. <i>Eurypanopeus abbreviatus</i> (Stimpson, 1860).....	56, 607, 617
590. <i>Eurypanopeus depressus</i> (Smith, 1869).....	56, 607, 617
591. <i>Eurypanopeus dissimilis</i> (Benedict and Rathbun, 1891).....	56, 607, 617
592. <i>Eurypanopeus turgidus</i> (Rathbun, 1930).....	56, 607, 617
593. <i>Eurytium limosum</i> (Say, 1818).....	56, 606, 655
594. <i>Glyptoxanthus erosus</i> (Stimpson, 1859).....	57, 603, 655
595. <i>Heteractaea ceratopus</i> (Stimpson, 1860).....	57, 604, 655
596. <i>Hexapanopeus angustifrons</i> (Benedict and Rathbun, 1891).....	57, 608, 621
597. <i>Hexapanopeus caribbaeus</i> (Stimpson, 1871).....	57, 608, 619
598. <i>Hexapanopeus hemphillii</i> (Benedict and Rathbun, 1891).....	57, 608, 621
599. <i>Hexapanopeus lobipes</i> (A. Milne Edwards, 1880).....	57, 608, 619
600. <i>Hexapanopeus paulensis</i> Rathbun, 1930.....	57, 608, 619
601. <i>Hexapanopeus quinquedentatus</i> Rathbun, 1901.....	57, 608, 621
602. <i>Leptodius parvulus</i> (Fabricius, 1793).....	57, 604, 655
603. <i>Lobopilumnus agassizii</i> (Stimpson, 1871).....	57, 606, 657
604. <i>Melybia thalamita</i> Stimpson, 1871.....	57, 606, 657
605. <i>Menippe mercenaria</i> (Say, 1818).....	57, 608, 623
606. <i>Menippe nodifrons</i> Stimpson, 1859.....	58, 608, 623
607. <i>Micropanope barbadensis</i> (Rathbun, 1921).....	58, 609, 627

608. <i>Micropanope lobifrons</i> A. Milne Edwards, 1880.....	58, 609, 625
609. <i>Micropanope nuttingi</i> (Rathbun, 1898).....	58, 609, 627
610. <i>Micropanope pusilla</i> A. Milne Edwards, 1880.....	58, 609, 625
611. <i>Micropanope sculptipes</i> Stimpson, 1871.....	58, 609, 625
612. <i>Micropanope spinipes</i> A. Milne Edwards, 1880.....	58, 609, 625
613. <i>Micropanope urinator</i> (A. Milne Edwards, 1881).....	58, 609, 627
614. <i>Neopanope packardii</i> (Kingsley, 1879).....	58, 609, 629
615. <i>Neopanope sayi</i> (Smith, 1869).....	58, 609, 629
616. <i>Neopanope texana</i> (Stimpson, 1859).....	58, 609, 629
617. <i>Panopeus americanus</i> Saussure, 1857.....	58, 610, 631
618. <i>Panopeus bermudensis</i> Benedict and Rathbun, 1891.....	59, 610, 631
619. <i>Panopeus hartii</i> Smith, 1869.....	59, 611, 635
620. <i>Panopeus herbstii</i> H. Milne Edwards, 1834.....	59, 610, 631
621. <i>Panopeus lacustris</i> Desbonne, 1867.....	59, 610, 633
622. <i>Panopeus obesus</i> Smith, 1869.....	59, 610, 633
623. <i>Panopeus occidentalis</i> Saussure, 1857.....	59, 611, 635
624. <i>Panopeus rugosus</i> A. Milne Edwards, 1880.....	59, 610, 633
625. <i>Panopeus simpsoni</i> Rathbun, 1930.....	59, 610, 631
626. <i>Paractaea rufopunctata nodosa</i> (Stimpson, 1860).....	59, 604, 657
627. <i>Paraliomera dispar</i> (Stimpson, 1871).....	59, 611, 637
628. <i>Paraliomera longimana</i> (A. Milne Edwards, 1865).....	60, 611, 637
629. <i>Pilumnoides nudifrons</i> (Stimpson, 1871).....	60, 606, 657
630. <i>Pilumnus caribaeus</i> Desbonne and Schramm, 1867.....	60, 612, 641
631. <i>Pilumnus dasypodus</i> Kingsley, 1879.....	60, 612, 641
632. <i>Pilumnus floridanus</i> Stimpson, 1871.....	60, 612, 643
633. <i>Pilumnus gemmatus</i> Stimpson, 1860.....	60, 612, 641
634. <i>Pilumnus holosericus</i> Rathbun, 1898.....	60, 612, 643
635. <i>Pilumnus lacteus</i> Stimpson, 1871.....	60, 612, 643
636. <i>Pilumnus longleyi</i> Rathbun, 1930.....	60, 611, 639
637. <i>Pilumnus marshi</i> Rathbun, 1901.....	60, 611, 639
638. <i>Pilumnus nudimanus</i> Rathbun, 1900.....	60, 611, 639
639. <i>Pilumnus pannosus</i> Rathbun, 1896.....	60, 612, 643
640. <i>Pilumnus sayi</i> Rathbun, 1897.....	60, 612, 641
641. <i>Pilumnus spinosissimus</i> Rathbun, 1898.....	60, 611, 639
642. <i>Platyactaea setigera</i> (H. Milne Edwards, 1834).....	61, 603, 659
643. <i>Platypodiella spectabilis</i> (Herbst, 1794).....	61, 603, 659
644. <i>Pseudomedaes agassizii</i> (A. Milne Edwards, 1880).....	61, 613, 645
645. <i>Pseudomedaes distinctus</i> (Rathbun, 1898).....	61, 613, 645
646. <i>Rhithropanopeus harrisii</i> (Gould, 1841).....	61, 605, 659
647. <i>Tetraxanthus bidentatus</i> (A. Milne Edwards, 1880).....	61, 613, 647
648. <i>Tetraxanthus rathbunae</i> Chace, 1939.....	61, 613, 647
649. <i>Xantho denticulata</i> White, 1847.....	61, 603, 659

Family Gecarcinidae

650. <i>Cardisoma guanhumi</i> Latreille, 1825.....	61, 661, 663
651. <i>Gecarcinus lateralis</i> (Fremenville, 1835).....	61, 661, 663
652. <i>Gecarcinus ruricola</i> (Linnaeus, 1758).....	62, 661, 663

Family Grapsidae

653. <i>Aratus pisonii</i> (H. Milne Edwards, 1837).....	62, 665, 675
654. <i>Cyclograpsus integer</i> H. Milne Edwards, 1837.....	62, 665, 675
655. <i>Euchirograpsus americanus</i> A. Milne Edwards, 1880.....	62, 666, 669
656. <i>Euchirograpsus antillensis</i> Turkay, 1975.....	62, 666, 669
657. <i>Geograpsus lividus</i> (H. Milne Edwards, 1837).....	62, 665, 675
658. <i>Goniopsis cruentata</i> (Latreille, 1802).....	62, 665, 675
659. <i>Grapsus grapsus</i> (Linnaeus, 1758).....	62, 665, 677
660. <i>Pachygrapsus gracilis</i> (Saussure, 1858).....	62, 666, 669
661. <i>Pachygrapsus transversus</i> (Gibbes, 1850).....	63, 666, 669

662. <i>Percnon gibbesi</i> (H. Milne Edwards, 1853).....	63, 665, 677
663. <i>Plagusia depressa</i> (Fabricius, 1775).....	63, 665, 677
664. <i>Planes minutus</i> (Linnaeus, 1758).....	63, 665, 679
665. <i>Platychoirograpsus spectabilis</i> De Man, 1896.....	63, 665, 679
666. <i>Sesarma benedicti</i> Rathbun, 1897.....	63, 667, 671
667. <i>Sesarma cinereum</i> (Bosc, 1802).....	63, 667, 671
668. <i>Sesarma curacaoense</i> De Man, 1892.....	63, 667, 673
669. <i>Sesarma miersii</i> Rathbun, 1897.....	63, 667, 671
670. <i>Sesarma reticulatum</i> (Say, 1817).....	63, 667, 673
671. <i>Sesarma ricordi</i> H. Milne Edwards, 1853.....	63, 667, 671

Family Pinnotheridae

672. <i>Dissodactylus borradalei</i> Rathbun, 1918.....	64, 682, 687
673. <i>Dissodactylus crinitichelis</i> Moreira, 1901.....	64, 682, 689
674. <i>Dissodactylus mellitae</i> (Rathbun, 1900).....	64, 682, 689
675. <i>Dissodactylus primitivus</i> Bouvier, 1917.....	64, 682, 687
676. <i>Dissodactylus rugatus</i> Bouvier, 1917.....	64, 682, 687
677. <i>Dissodactylus stebbingi</i> Rathbun, 1918.....	64, 682, 687
678. <i>Fabia byssomiae</i> (Say, 1818).....	64, 682, 691
679. <i>Fabia tellinae</i> Cobb, 1973.....	64, 682, 691
680. <i>Orthotheres strombi</i> (Rathbun, 1905).....	64, 681, 705
681. <i>Parapinnixa bouvieri</i> Rathbun, 1918.....	64, 683, 693
682. <i>Parapinnixa hendersoni</i> Rathbun, 1918.....	64, 683, 693
683. <i>Pinnaxodes floridensis</i> Wells and Wells, 1961.....	64, 681, 705
684. <i>Pinnixa chacei</i> Wass, 1955.....	64, 683, 695
685. <i>Pinnixa chaetoptera</i> Stimpson, 1860.....	64, 684, 699
686. <i>Pinnixa cristata</i> Rathbun, 1900.....	65, 683, 695
687. <i>Pinnixa cylindrica</i> (Say, 1818).....	65, 684, 697
688. <i>Pinnixa floridana</i> Rathbun, 1918.....	65, 684, 697
689. <i>Pinnixa leptosynaptae</i> Wass, 1968.....	65, 683, 695
690. <i>Pinnixa lunzi</i> Glassell, 1937.....	65, 684, 697
691. <i>Pinnixa pearsei</i> Wass, 1955.....	65, 684, 699
692. <i>Pinnixa retinens</i> Rathbun, 1918.....	65, 684, 695
693. <i>Pinnixa sayana</i> Stimpson, 1860.....	65, 684, 697
694. <i>Pinnotheres hemphilli</i> Rathbun, 1918.....	65, 685, 703
695. <i>Pinnotheres maculatus</i> Say, 1818.....	65, 685, 701, 703
696. <i>Pinnotheres moseri</i> Rathbun, 1918.....	65, 685, 701
697. <i>Pinnotheres ostreum</i> Say, 1817.....	65, 685, 701, 703
698. <i>Pinnotheres shoemakeri</i> Rathbun, 1918.....	65, 685, 703

Family Ocypodidae

699. <i>Ocypode quadrata</i> (Fabricius, 1787).....	66, 707, 717
700. <i>Uca burgersi</i> Holthuis, 1967.....	66, 709, 715
701. <i>Uca leptodactyla</i> Rathbun, 1898.....	66, 707, 711
702. <i>Uca longisignalis</i> Salmon and Atsides, 1968.....	66, 708, 713
703. <i>Uca minax</i> (Le Conte, 1855).....	66, 709, 715
704. <i>Uca panacea</i> Novak and Salmon, 1974.....	66, 707, 711
705. <i>Uca pugilator</i> (Bosc, 1802).....	66, 707, 711
706. <i>Uca pugnax</i> (Smith, 1870).....	66, 709, 715
707. <i>Uca rapax</i> (Smith, 1870).....	66, 709, 715
708. <i>Uca speciosa</i> (Ives, 1891).....	66, 708, 713
709. <i>Uca spinicarpa</i> Rathbun, 1900.....	66, 708, 713
710. <i>Uca thayeri</i> Rathbun, 1900.....	67, 708, 711
711. <i>Uca vocator</i> (Herbst, 1804).....	67, 708, 713
712. <i>Ucides cordatus</i> (Linnaeus, 1763).....	67, 707, 717

Family Palicidae

713. *Palicus affinis* A. Milne Edwards and Bouvier, 1899..... 67, 718, 721
714. *Palicus alternatus* Rathbun, 1897..... 67, 718, 721
715. *Palicus cristatipes* (A. Milne Edwards, 1880)..... 67, 718, 721
716. *Palicus cursor* (A. Milne Edwards, 1880)..... 67, 719, 723
717. *Palicus dentatus* A. Milne Edwards, 1880..... 67, 718, 723
718. *Palicus faxoni* Rathbun, 1897..... 67, 718, 723
719. *Palicus floridana* (Rathbun, 1918)..... 67, 719, 725
720. *Palicus gracilis* (Smith, 1883)..... 67, 719, 725
721. *Palicus obesus* (A. Milne Edwards, 1880)..... 67, 718, 723
722. *Palicus sica* (A. Milne Edwards, 1880)..... 68, 718, 721

Family Cryptochiridae

723. *Pseudocryptochirus corallicola* (Verrill, 1908)..... 68, 727, 729
724. *Pseudocryptochirus hypostegus* Shaw and Hopkins, 1977..... 68, 727, 729

Introduction

There are more species of shrimps, lobsters, and crabs in the marine waters of the state of Florida than in any other region of the continental United States. This great diversity is the result of three factors: (1) Florida's coastline is extensive; (2) a tremendous number of marine habitats occurs in Florida; and (3) two biogeographic regions come together in Florida, the northern *Carolinian* and the southern *Caribbean* (or *West Indian*). The total number of decapods in the marine

shallow waters of Florida is probably close to 900 (see Methods and Materials).

The literature on Florida's decapods is scattered and incomplete and often lacks keys or illustrations. The present contribution is an attempt to remedy this situation. We have compiled a checklist, keys, and illustrations of all marine, shallow-water (≤ 300 m) decapod crustaceans known to occur in Florida, a total of 724 species.

Brief Review of the Literature

Published work on the decapod crustaceans occurring in Florida dates from the earliest explorations by Europeans in the Caribbean and western Atlantic. Many of these contained brief descriptions and even some illustrations (e. g., Sloane, 1725), which were given nomenclatural status by Linnaeus (1758). Many of the species described by Gibbes (1850), Stimpson (1860, 1871), and Kingsley (1878) occurred in the marine waters of Florida. Professor H. E. Webster collected along the gulf coast of Florida, and these collections were published on by Kingsley (1879). Additional Florida collections were reported on by Ives (1891). The *Blake* expeditions included Florida material, which was published by A. Milne Edwards and Bouvier (1893, 1897, 1902, 1909, 1923).

Modern published work dealing explicitly with Florida marine decapods is scattered. Chace (1942a) described five new species of decapods from the west coast of Florida. Wass (1955) published an annotated list of the decapods of northwestern Florida and described three new species. Provenzano (1959) reviewed the hermit crabs of Florida and described one new species. Hulings (1961) added several new records from deeper water in the northeastern gulf. Wells and Wells (1961) described a new species of crab from the northern gulf, as did Salmon and Atsides (1968) and Novak and Salmon (1974). Although not dealing specifically

with Florida, William's (1965a) excellent volume on the decapods of the Carolinas contains a tremendous amount of information relevant to the Florida fauna. The 1984 revision of this work (Williams, 1984) extended the coverage to include the northeastern Atlantic coast of Florida.

Beginning in the 1970's, there was a virtual explosion of information on Florida decapods. Biffar (1971a, 1971b) reviewed the callinassids of Florida. Abele (1971, 1972a, 1972b, 1972c, 1973) reviewed the status of some palaemonids, xanthids, atyids, and grapsids in Florida. McLaughlin and Provenzano (1974a, 1974b), and McLaughlin and co-workers (McLaughlin, 1981a, 1981b, 1982; García-Gómez, 1982; Lemaitre, 1982; Lemaitre et al., 1982) provided extensive information on the hermit crabs of Florida. Beginning in 1973, Robert H. Gore published a series of papers on the decapods of the Indian River region of Florida, including a major work on parthenopid crabs (e. g., Hendrix and Gore, 1973; Gore 1977, 1979, 1981; Gore and Wilson, 1978; Gore and Scotto, 1979; Kensley and Gore, 1981). Mayo (1973) reviewed the genus *Cancellus* including Florida material. Additional data on Florida decapods can be found in Rouse (1970) and in the results of the *Hourglass* cruises (e. g., Lyons, 1970; Cobb, 1971, 1973; Cobb et al., 1973).

Methods and Materials

The checklist is derived from several sources. First, it is based on a search of the literature, and we have cited these sources in the text. Additions were made based on our own collections, those of Dr. Patsy McLaughlin, and those housed in the following institutions: the Smithsonian Institution's National Museum of Natural History (SI-NMNH); the State of Florida's Department of Natural Resources Collections in St. Petersburg, Florida; and the collections of Harbor Branch Foundation's Indian River Coastal Zone Museum in Fort Pierce, Florida.

Keys were prepared from the literature sources cited in the headings of the keys and verified by us in so far as was possible.

This volume is intended as an identification guide, and we wished to provide a standard format for the illustrations. The illustrations were therefore re-drawn from the sources

cited. We were able to locate source illustrations for 722 of the 724 species, although they varied in quality. We were unable to locate specimens or illustrations of the majid crab *Collodes nudus* and the goneplacid crab *Pilumnoplax elata*. We urge our readers to verify their identifications using the primary literature. The scaling is in a standard format: a single line indicates that the scale is in millimeters, and a double line indicates centimeters; the number of units is indicated by the number of tick marks shown on the scale.

We made every effort to indicate all species known from Florida's marine waters. The checklist, however, is not complete for three reasons: (1) we were working on a time schedule and could not search indefinitely; (2) there are numerous undescribed species that are either currently being studied or in museums, and we did not believe it appropriate to include these; and (3) we are sure that we simply missed some species.

Classification and Arrangement of Taxa and Illustrations

We have generally followed the classification given in Bowman and Abele (1982) and the arrangement in Abele and Felgenhauer (1982). Within families we have arranged the genera (and species within genera) alphabetically. The illustrations are grouped by family and follow the key to that family. They are grouped by genera and within genera by the sequence that they occur in the key with the

following exception: Genera containing a single species are grouped together at the end of the family. This was done simply to save space.

Taxonomic nomenclature follows the most recent revision available unless general usage dictates otherwise. This is an identification guide not a revision.



Acknowledgements

We owe a great debt to the many authors whose work is cited in this volume. We thank the following individuals for their comments: David Camp, Fenner A. Chace, Jr., Michael Dardeau, Isabel Pérez Farfante, Darryl Felder, Bruce Felgenhauer, Brian Kensley, Patsy McLaughlin, Raymond Manning, Jody Martin, Paula Mikkelsen, and Austin Williams. A special word of thanks is due David Camp, Patsy McLaughlin, and Paula Mikkelsen for comparing our checklist with collections under their care as well as for providing detailed information on species and keys. All errors are our own.

A special thanks also to Landon Ross of the Florida Department of Environmental Regulation for efforts above and beyond the

call of duty in assisting with the completion of this project. Similar thanks to Steve Wolfe for the extensive work involved in the production of this volume.

We thank Lisa Velez for her help in library research and for entering thousands of references into our computerized literature system. Anne Thistle typed the keys and legends, patiently revising them numerous times. This work was supported, in part, by the Department of Environmental Regulation, State of Florida, through a contract administered by the Florida Institute of Government. Partial support was provided by the National Science Foundation grant no. BSR 85-08430.



Schematic drawing of brachyuran crab

Checklist of the Decapod Crustaceans of Florida

SUBORDER DENDROBRANCHIATA

FAMILY ARISTEIDAE

Aristaeomorpha foliacea (Risso, 1827)

Description: Zariquiey Alvarez, 1968:42, figs. 22a, b, 24a.--Holthuis, 1980b:8.

Type-locality: Vicinity of Nice.

Distribution: Eastern Atlantic: Bay of Biscay to NW Africa and the entire Mediterranean. Western Atlantic: South of Massachusetts to the Straits of Florida, Gulf of Mexico, Caribbean Sea and off Venezuela. Indo-West Pacific: East Africa to Japan, Australia, New Zealand and Fiji (Holthuis, 1980b).

Plesiopenaeus edwardsianus (Johnson, 1867)

Description: Crosnier and Forest, 1973:292, figs. 98, 99a, b.--Holthuis, 1980b:11.

Type-locality: Madeira

Distribution: Eastern Atlantic: Portugal to South Africa; (not in the Mediterranean). Western Atlantic: Grand Bank (43°42'N) to Gulf of Mexico, Caribbean Sea and north coast of South America (Holthuis, 1980b). Indo-West Pacific: off east Africa to Japan, Australia.

FAMILY BENTHESICYMIDAE

Bentheogennema intermedia (Bate, 1888)

Description: Bate, 1888:343, pl. 58: fig. 3.--Roberts and Pequegnat, 1970:39.

Type-locality: Off Sierra Leone, 01°47'N, 24°26'W; 3386 m; *Challenger* station 106.

Distribution: Appears to be distributed worldwide; probably pelagic (Roberts and Pequegnat, 1970); Florida (personal communication, P. M. Mikkelsen).

FAMILY PENAEIDAE

Funchalia villosa (Bouvier, 1905)

Description: Burkenroad, 1936:129

Type-locality: Between Canary Islands and Azores.

Distribution: Adults are known from the eastern and western North Atlantic, Mediterranean, the south central South Atlantic, and the Caribbean; larvae known from the western North Atlantic and South Pacific. Florida (personal communication, D. K. Camp, P. M. Mikkelsen).

Metapenaeopsis gerardoi Pérez Farfante, 1971

Description: Pérez Farfante, 1971:20, figs. 11, 12, 13c.

Type-locality: Off Mayaguez, Puerto Rico, 18°8.5'N, 67°23'W.

Distribution: Bahamas, Florida Keys, West Indies, and Caribbean coast of Central and South America (Pérez Farfante, 1971).

Metapenaeopsis goodei (Smith, 1885)

Description: Pérez Farfante, 1971:9, figs. 4-8.--Williams, 1984:36, 37, figs. 17-18.

Type-locality: Bermudas.

Distribution: Bermuda; between Capes Hatteras and Lookout, North Carolina, through Florida Straits and along west Florida to Pensacola; Isla de Lobos reef, Veracruz (Ray, 1974), around Yucatan Peninsula, through Caribbean Sea, and along South American coast to Espírito Santo, Brazil (Coelho and Ramos, 1972; Williams, 1984).

Metapenaeopsis smithi (Schmitt, 1924)

Description: Schmitt, 1924a:62, figs. 1b, c, 2a, c.--Pérez Farfante, 1971: 29, figs. 13E, 18-22.

Type-locality: Caracas Baai, Curaçao.

Distribution: Bermudas and southeastern Florida to Curaçao, mostly near islands; western Caribbean; Lesser Antilles (Chace, 1972).

Parapenaeus americanus (Rathbun, 1901)

Description: A. Milne Edwards and Bouvier, 1909:231.

Type-locality: Off Mayaguez Harbor, Puerto Rico, 412 m. *Fish Hawk* Stn. 6070.

Distribution: 40°03'N, 70°49'W northern Uruguay, 33°26'S, 59°58'W; Puerto Rico; St. Lucia; Martinique.

Parapenaeus politus Smith, 1881

Description: Pérez Farfante, 1982:200, figs. 3-5.--Williams, 1984:37, fig. 19.

Type-locality: Off "Martha's Vineyard"

(Smith, 1885) 39°55'00"N, 70°54'15"W, 260 m.

Distribution: Martha's Vineyard, Massachusetts, through Gulf of Mexico; Gulf of Paria off Venezuela (Williams, 1984). Florida (personal communication, P. M. Mikkelsen).

***Penaeopsis serrata* Bate, 1881**

Description: Pérez Farfante, 1980b:748, figs. 28-36.

Type-locality: Off Barbados, "Gulf of Mexico", Blake Stn. 275, 399 m.

Distribution: Western Atlantic: from east of Barnegate, New Jersey south of Martha's Vineyard, Massachusetts, through the Gulf of Mexico and the Caribbean south to French Guiana; off Rio Grande do Sul, Brazil.

Eastern Atlantic: from south of Cabo San Vicente, Portugal, to off Cadiz, Spain and off the northwest coast of Africa to Tamzak ("Tamxat"), Mauritania (Pérez Farfante, 1980b).

***Penaeus aztecus* Ives, 1891**

Description: Pérez Farfante, 1969:527, figs. 39-43, 46-48, 50.--Williams, 1984:24, figs. 9, 10.

Type-locality: Veracruz, Mexico.

Distribution: Martha's Vineyard, Massachusetts around Florida Peninsula to northwestern Yucatan (Williams, 1984).

***Penaeus brasiliensis* Latreille, 1817**

Description: Pérez Farfante, 1969:562, figs. 68, 75, 76.--Williams, 1984:28, figs. 11-12.

Type-locality: Brazil.

Distribution: Bermudas; off Cape Hatteras, North Carolina to Florida Keys, off Campeche and Yucatan; through Caribbean Sea to Rio Grande do Sul, Brazil (Williams, 1984).

***Penaeus duorarum* Burkenroad, 1939**

Description: Pérez Farfante, 1969:499, figs. 20-22, 25-27, 30-31.-- Williams, 1984:28, 29, figs. 13-14.

Type-locality: Off Mobile Bay, Alabama (29°15'N and 88°11'W, 36.5 m *Atlantis* Stn. 2813).

Distribution: Lower Chesapeake Bay through Florida Straits, around Mexico to Cape Catoche and Isla Mujeres at the tip of Yucatan Peninsula (Williams, 1984).

***Penaeus setiferus* (Linnaeus, 1767)**

Description: Pérez Farfante, 1969:468, figs. 4-11.--Williams, 1984:32, figs. 15-16.

Type-locality: Off Matanzas Inlet, Florida.

Distribution: Fire Island, New York, to Saint Lucie Inlet, Florida; near Dry Tortugas, Florida (rarely); Gulf of Mexico from Ochlocknee River, Florida, to Campeche, Mexico (Williams, 1984).

***Trachypenaeus constrictus* (Stimpson, 1874)**

Description: Williams, 1984:38, figs. 20, 21.

Type-locality: Beaufort, North Carolina.

Distribution: Tangier Sound, Chesapeake Bay, to Veracruz, Mexico; Bermuda; Cuba, Puerto Rico and Sombbrero Island; Surinam; off Ceara, São Paulo, and Santa Catarina, Brazil (Williams, 1984).

***Trachypenaeus similis* (Smith, 1885)**

Description: Burkenroad, 1934:96, figs. 10, 11.

Type-locality: Gulf of Paria, Venezuela, 10°37'40"N, 61°42'40"W.

Distribution: Gulf of Mexico to Estado do Pará, Brazil.

***Trachypeneopsis mobilispinis* (Rathbun, 1920)**

Description: Rathbun, 1920:320, figs. 1, 2a-c --Chace, 1972:10.

Type-locality: Cave Round Bay, Saba.

Distribution: Bermudas and Bay of Campeche eastward to Cuba, Cay Sal Bank, Turks, Saba, Barbuda, Jamaica, Virgin Islands and Saint Christopher Islands (Chace, 1972).

***Xiphopenaeus kroyeri* (Heller, 1862)**

Description: Williams, 1984:40, figs. 22, 23.

Type-locality: Rio de Janeiro, Brazil.

Distribution: Between Capes Hatteras and Lookout, North Carolina, through Gulf of Mexico and Caribbean Sea to Ponta do Zimbrow, Santa Catarina, Brazil (Pérez Farfante, 1978).

FAMILY SOLENOCERIDAE***Hadropenaeus affinis* (Bouvier, 1906)**

Description: Pérez Farfante, 1977:317 figs. 9, 43, 44A, 45-49.

Type-locality: Off Cape Verde Island (16°53'N, 25°10'W, 410-460 m).

Distribution: Western Atlantic: from off Cape Lookout, North Carolina southward to the Straits of Florida, in the northeastern part of the Gulf of Mexico and throughout Caribbean. Eastern Atlantic: off Cape Verde Islands (Pérez Farfante, 1977).

***Hadropenaeus modestus* (Smith, 1885)**

Description: Pérez Farfante, 1977:323, figs. 9, 44, 46, 49-52.

Type-locality: Off Bethany Beach, Delaware, 38°31'N, 73°21'W, 285 m. *Fish Hawk* Stn. 1047.

Distribution: Off Delaware Bay, to the Straits

of Florida, and in the Gulf of Mexico, northwest of Charlotte Harbor, Florida; Bahamas throughout the Caribbean to off Barra Grande, Brazil (Pérez Farfante, 1977).

Hymenopenaeus aphoticus Burkenroad, 1936

Description: Pérez Farfante, 1977:275, figs. 4C, 8-12.

Type-locality: Turks Island Passage, 1,646-1,728 m, 21°15'40"N, 71°17'06"W, *Pawnee* Stn. 54.

Distribution: Western Atlantic: southwest Florida throughout the Gulf of Mexico, and Caribbean Sea. Eastern Atlantic: south of the Azores Islands and off Morocco. (Pérez Farfante, 1977).

Hymenopenaeus debilis Smith, 1882

Description: Pérez Farfante, 1977:268 figs. 2, 3, 4B, 5-9.

Type-locality: Syntypes: SE of Savannah Beach, Georgia, 31°57'00"N, 78°18'35"W, 609 m, *Blake* Stn. 317; SE of Cape Fear, North Carolina, 33°19'00"N, 76°13'30"W, 836 m, *Blake* Stn. 323; E of Cape Fear, North Carolina 33°42'15"N, 76°00'50"W, 849 m, *Blake* Stn. 326.

Distribution: Western Atlantic: from Hudson Canyon, New Jersey (39°55'N, 70°31'W) through Gulf of Mexico and Caribbean Sea to Guyana (08°14'N, 57°38'W). Eastern Atlantic: Azores Islands and northwest Africa from Cape Spartel, Morocco, to Cape Verde Islands, including Canary Islands (Pérez Farfante, 1977).

Mesopenaeus tropicalis (Bouvier, 1905)

Description: Pérez Farfante, 1977:332, figs. 56-58, 60-63.--Williams, 1984:18, fig. 5.

Type-locality: Mer des Antilles (Pérez Farfante 1977).

Distribution: Northeast of Cape Lookout, North Carolina, 34°43'N, 76°40'W, through Florida Straits to Alabama; off Cape Catoche and Bahamas through Caribbean Sea and along coast of South America to Rio Grande do Sul, Brazil (Williams, 1984).

Pleoticus robustus (Smith, 1885)

Description: Pérez Farfante, 1977:297, figs. 9, 29-36.

Type-locality: South of Curaçao 11°43'00"N, 69°09'30"W, 380 m., *Albatross* Stn. 2125.

Distribution: South of Martha's Vineyard,

Massachusetts, through Gulf of Mexico, and the Caribbean to French Guiana (Pérez Farfante, 1977).

Solenocera atlantidis Burkenroad, 1939

Description: Pérez Farfante and Bullis, 1973:20, figs. 11, 13, 14.-- Williams, 1984:19, fig. 6.

Type-locality: Gulf of Mexico off Alabama, 29°45'N, 88°11'W, 37 m. *Atlantis* Stn. 281.

Distribution: Off Oregon Inlet, North Carolina, around Gulf of Mexico and Caribbean Sea to Cananeia, São Paulo, Brazil (Pérez Farfante and Bullis, 1973).

Solenocera necopina Burkenroad, 1939

Description: Pérez Farfante and Bullis, 1973:14, figs. 7, 9, 10.-- Williams, 1984:20, fig. 7.

Type-locality: Off Mobile Bay, Alabama, 29°16'N, 87°54'W, 229 m. *Atlantis* Stn. 2377.

Distribution: South of New England (40°04'N, 70°29'W) through Gulf of Mexico and Caribbean Sea to Rio Grande do Sul, Brazil and Uruguay (Williams, 1984; Pérez Farfante, 1977).

Solenocera vioscai Burkenroad, 1939

Description: Pérez Farfante and Bullis, 1973:3, figs. 1A, B, 3.--Williams, 1984:21, fig. 8.

Type-locality: About 5 mi. (8km) off Pass a L'outre, Louisiana, 27 m.

Distribution: Southeast of Cape Lookout, North Carolina, to Dry Tortugas, Florida (but rare off Florida); northern and western Gulf of Mexico to Tabasco (Williams, 1984).

FAMILY SICYONIIDAE

Sicyonia brevirostris Stimpson, 1871

Description: Williams, 1984:43, figs. 25, 26.

Type-locality: Cuba.

Distribution: Off Norfolk, Virginia, through Bahamas to Southern Cuba; around Gulf of Mexico to southern Texas; Campeche Banks to Isla Contoy, Yucatan; doubtful record of its occurrence on Pacific Coast of southern Mexico (Williams, 1984).

Sicyonia burkenroadii Cobb, 1971

Description: Williams, 1984:46, fig. 27.

Type-locality: Gulf of Mexico off Port Isabel, Texas, 26°13'N, 96°45'W, 42 m.

Distribution: Off Cape Lookout, North

Carolina, 34°12'N, 76°11'W, through Gulf of Mexico to Bahia, Brazil (Pérez Farfante 1980a); occasionally as deep as 585 m.

Sicyonia dorsalis Kingsley, 1878

Description: Williams, 1984:46, fig. 28.

Type-locality: Fort Jefferson, Dry Tortugas, Florida.

Distribution: Cape Hatteras, North Carolina to Texas; Colombia to French Guiana; Ceara to Santos and Santa Catarina, Brazil (Williams, 1984).

Sicyonia laevigata Stimpson, 1871

Description: Williams, 1984:47, fig. 29.

Type-locality: Charleston, South Carolina.

Distribution: Cape Hatteras, North Carolina, to northwest Florida; through West Indies to Colombia, and Santa Catarina, Brazil (Pérez Farfante 1980a). Pacific Coast of Panama.

Sicyonia parri (Burkenroad, 1934)

Description: Williams, 1984:48, fig. 30.

Type-locality: Crooked Island, Bahamas.

Distribution: Beaufort, North Carolina through Gulf of Mexico and West Indies to Bahia, Brazil (Williams, 1984).

Sicyonia stimpsoni Bouvier, 1905

Description: Williams, 1984:49, fig. 31.

Type-locality: Off Barbados, 13°03'05"N, 59°36'18"W, 185 m.

Distribution: Near Cape Hatteras through Florida Straits, and including west Florida, to Barbados and Columbia to Surinam (Williams, 1984).

Sicyonia typica (Boeck, 1864)

Description: Williams, 1984:49, fig. 32.

Type-locality: Molde Fjord, west coast of Norway (erroneous locality, evidently incorrectly labeled).

Distribution: Off Wrightsville Beach, North Carolina, through Gulf of Mexico; Cuba through West Indies to near Ilha de Santa Catarina, Brazil (Williams, 1984).

FAMILY SERGESTIDAE

Acetes americanus carolinae Hansen, 1933

Description: Williams, 1984:50, fig. 33.

Type-locality: Beaufort Inlet (about 34°47'N), North Carolina.

Distribution: Lower Chesapeake Bay (Mobjack Bay and York River) through Gulf of Mexico to Panama, Surinam and French Guiana (Williams, 1984).

Sergestes armatus Kröyer, 1855

Description: Kensley, 1971:232, fig. 8.

Type-locality: Tropical Atlantic.

Distribution: North Atlantic, Mediterranean, off Azores and Canaries, South Atlantic. Florida (85-250 m) (personal communication, P. M. Mikkelsen).

Sergestes atlanticus H. Milne Edwards, 1830

Description: Kensley, 1971:234, fig. 9.

Type-locality: Near Azores.

Distribution: Mediterranean, North Atlantic, off Azores and Canaries, Sargassum Sea, Off Cape Point. Florida (75-250 m) (personal communication, P. M. Mikkelsen).

Sergestes edwardsii Kröyer, 1855

Description: Crosnier and Forest, 1973:320, figs. 108b, 100b-c, 110c-d.

Type-locality: Tropical Atlantic from 3 degrees S to 10 degrees N.

Distribution: Atlantic Ocean; also reported from the Indo-Pacific and Eastern Pacific but there is some question about the identity of material outside of the Atlantic (Crosnier and Forest, 1973). Florida (65-300 m) (personal communication, P. M. Mikkelsen).

Sergestes henseni (Ortmann, 1893)

Description: Ortmann, 1893:38, pl. 3: fig. 3.

Type-locality: North Atlantic off West Africa.

Distribution: Mediterranean; north Atlantic; western and southern Atlantic; (Crosnier and Forest, 1973). Florida (85-250 m) (personal communication, P. M. Mikkelsen).

Sergestes paraseminudus Crosnier and Forest, 1973

Description: Crosnier and Forest, 1973:313, figs. 105d, 106c-d,f.

Type-locality: Tropical eastern Atlantic; 01°55'S, 8°30'E; 0-50 m.

Distribution: Tropical eastern Atlantic; Florida (85-250 m) (personal communication, P. M. Mikkelsen).

Sergestes pectinatus Sund, 1920

Description: Kensley, 1971:240, fig. 13.

Type-locality: "Atlantic Ocean, equatorial region."

Distribution: Sargassum Sea, North Atlantic, Off Canaries, Azores. Florida (85-250 m) (personal communication, P. M. Mikkelsen).

Sergestes sargassi Ortmann, 1893

Description: Hansen, 1922:148, pl. 9: fig. 2a-p

Type-locality: Off Florida, Sargassum Sea.
Distribution: NE Atlantic, Sargassum Sea;
Florida (75-750 m) (personal communication,
P. M. Mikkelsen).

***Sergestes vigilax* Stimpson, 1860**

Description: Hansen, 1922:159, pl. 1: fig. 8.
Type-locality: Off the Azores.
Distribution: NE Atlantic, Indian Ocean from
Gulf of Aden to Ceylon and Cocos-Keeling
Island; Florida (65-300 m) (personal
communication, P. M. Mikkelsen).

***Sergia extenuatus* Burkenroad, 1940**

Description: Crosnier and Forest, 1973:338,
figs. 112e-f, 113b, 114g.
Type-locality: Off the Ivory Coast, 03°45'S,
10°00'W.
Distribution: Tropical eastern Atlantic (Gabon
to Angola); Florida (85-250 m) (personal
communication, P. M. Mikkelsen).

***Sergia splendens* Sund, 1920**

Description: Kensley, 1971:260, fig. 23.
Type-locality: Unknown
Distribution: North Atlantic, Mediterranean
(off Monaco), off Table Bay. Florida (85-250
m) (personal communication, P. M.
Mikkelsen).

FAMILY LUCIFERIDAE

***Lucifer faxoni* Borradaile, 1915**

Description: Williams, 1984:52, fig. 34.
Type-locality: Hampton Roads, Virginia
(Chace 1972).
Distribution: "Coastal waters of North and
South America from Long Island Sound to
Rio de Janeiro," around Gulf of Mexico and
throughout Caribbean Sea (probably
widespread); Bermuda and scattered mid-
Atlantic occurrences in path of Gulf Stream;
Bay of Dakar, Senegal (Bowman and
McCain, 1967).

***Lucifer typus* H. Milne Edwards, 1837**

Description: Bowman and McCain,
1967:660, figs. 1a, b, 2-7.
Type-locality: Tropical North Atlantic.
Distribution: Warmer open waters of the
north and south Atlantic oceans approximately
between the 40 degrees parallels. Unlike
L. faxoni, this species is not usually found in
inshore waters (Bowman and McCain, 1967).

SUBORDER PLEOCYEMAT

**INFRAORDER
STENOPODIDEA**

FAMILY STENOPODIDAE

***Microprosthema semilaeve* (Von Martens,
1872)**

Description: Holthuis, 1946:54, pl. 3: fig. 1.
Type-locality: Cuba.
Distribution: Bahamas, southern Florida, and
Yucatan to Fernando de Noronha, Brazil
(Chace, 1972).

***Odontozona libertae* Gore, 1981**

Description: Gore, 1981:153, figs. 4a-c, 5a-l.
Type-locality: Elbow Reef, off Key Largo,
Monroe County, Florida.
Distribution: Known only from the type-
locality.

***Stenopus hispidus* (Olivier, 1811)**

Description: Williams, 1984:54, fig. 35.
Type-locality: "Australasiatic seas."
Distribution: Western Atlantic from North
Carolina (Kruczynski and Jenner, 1969) and
Bermuda, southern Florida through Gulf of
Mexico (Ray, 1974) to Fernando de Noronha
and Espirito Santo, Brazil (Coelho and
Ramos, 1972). Indo-Pacific from Durban,
South Africa, and Red Sea, to Japan, Hawaii,
western Australia south to ~24 degrees South
and eastern Australia as far south as
Shellharbour, New South Wales, through
New Caledonia, New Hebrides, Lord Howe
Island, northern New Zealand, to Tuamotu
Archipelago (Yaldwyn, 1968; Williams,
1984).

***Stenopus scutellatus* Rankin, 1898**

Description: Williams, 1984:56, fig. 36.
Type-locality: Silver Cay, New Providence,
Bahamas.
Distribution: Bermuda; South Carolina
(Wenner and Read, 1982); Gulf of Mexico to
Fernando de Noronha and Rio Grande do
Norte, Brazil (Williams, 1984).

INFRAORDER CARIDEA

FAMILY ATYIDAE

***Potimirim potimirim* (Müller, 1881)**

Description: Villalobos, 1960:327.--Abele,
1972c:215, figs. 3D, 4.

Type-locality: Rio Itahai Itahai, state of Santa Catarina, Brazil.

Distribution: Rio Itahai Itahai, State of Santa Catarina, and Rio Gurjau, Recife, state of Pernambuco, Brazil; introduced into the freshwaters of southern Florida (Abele, 1972c).

FAMILY OPLOPHORIDAE

Acanthephyra purpurea A. Milne Edwards, 1881

Description: Chace, 1940a:134, figs. 11-17.

Type-locality: Berlengas Island, off the West Coast of Portugal, 2590 m.

Distribution: North Atlantic between about 20° and 53° N latitude (Chace, 1940a).

Florida (85-250 m over 750 m) (personal communication, P. M. Mikkelsen).

Janicella spinicauda (A. Milne Edwards, 1883)

Description: Chace, 1986: figs. 23, 24.

Type-locality: Off Casablanca, Morocco: *Travailleur* Stn. 65; 34°13'30"N, 7°43'00"W, 636 m, muddy sand.

Distribution: Apparently widespread mesopelagically in the tropical seas of the world (except in the eastern Pacific off the Americas) (Chace, 1986).

Oplophorus gracilirostris A. Milne Edwards, 1881

Description: Chace, 1986: figs. 32a-32c.

Type-locality: Off Dominica, Lesser Antilles, 216 m.

Distribution: Off southeastern Africa, Indian Ocean, Indonesia, Philippines, southern Japan, Fiji Islands, Hawaii, Gulf of Mexico, Bahamas, Caribbean Sea; mesopelagic (Chace, 1986).

Oplophorus spinosus (Brullé, 1839)

Description: Crosnier and Forest, 1973:25.

Type-locality: Canary Islands (Holthuis, 1949b).

Distribution: Indian Ocean, southern Japan, off Hawaii, seamounts west of North America, and northeast of Easter Island, western and eastern subtropical North Atlantic, central South Atlantic; mesopelagic (Chace, 1986). Florida (150 m) (personal communication, P. M. Mikkelsen).

Systellaspis debilis (A. Milne Edwards, 1881)

Description: Chace, 1986: figs. 34g-i, 35 e-f.

Type-locality: "... trouvée à une profondeur de 500 brasses dans le canal de Bahama."

Distribution: South Africa, Indian Ocean, Philippines, Indonesia, Hawaii, western Atlantic from south of Greenland to Gulf of Mexico and Bahamas and eastern Atlantic from the Faeroe Islands to Angola; mesopelagic (Chace, 1986).

FAMILY PASIPHAEIDAE

Leptochela bermudensis Gurney, 1939

Description: Chace, 1976:7, figs. 5-7.

Type-locality: Seven miles south of Bermuda between 1000 m and surface.

Distribution: Bermuda to Barbados and southwestern Gulf of Mexico (Chace, 1976).

Leptochela carinata Ortmann, 1893

Description: Chace, 1976:45, figs. 35-37.-- Williams, 1984:59, fig. 39.

Type-locality: Off Baía de Marajo, Para, Brazil, 50-100 m.

Distribution: Georges Bank; South Carolina; Gulf of Mexico through Bahamas to Para, Brazil (Williams, 1984).

Leptochela papulata Chace, 1976

Description: Chace 1976:26, figs. 22-24.

Type-locality: East of Cape Lookout, North Carolina, 34°35'30"N, 75°45'30"W, 59 m.

Distribution: Georges Bank off Cape Cod, Massachusetts (Fontaine, 1977); North Carolina to Georgia; eastern Gulf of Mexico (Williams, 1984).

Leptochela serratorbita Bate, 1888

Description: Chace, 1976:36, figs. 29-31.-- Williams, 1984:58, fig. 38.

Type-locality: Saint Thomas, Virgin Islands. Distribution: Beaufort, North Carolina; South Carolina; western Gulf of Mexico and Florida Keys to Leeward Islands (Williams, 1984).

FAMILY BRESILIIDAE

Discias atlanticus Gurney, 1939

Description: Williams, 1984:62, fig. 41.

Type-locality: The Reach, Bermuda.

Distribution: Bermuda; off Savannah, Georgia, off Fort Pierce, Florida; Guadeloupe; Cape Verde Islands and Gabon; off northern Kenya (Bruce, 1975; Gore and Wilson, 1978); (Williams, 1984).

***Discias serratirostris* Lebour, 1949**

Description: Wilson and Gore, 1979:311, fig. 1
 Type-locality: Off Castle Roads, Bermuda.
 Distribution: 3 mi. off Castle Roads, Bermuda, Vero Beach, Indian River County, on the central eastern coast of Florida.

***Pseudochelones chacei* Kensley, 1983**

Description: Kensley, 1983:22, figs. 18-22.
 Type-locality: Looe Key, Florida.
 Distribution: Looe Key, Florida to Carrie Bow Cay, Belize.

FAMILY EUGONATONOTIDAE***Eugonatonotus crassus* (A. Milne Edwards, 1881)**

Description: A. Milne Edwards, 1881:10.-- Boone, 1927:106, figs. 22, 23.
 Type-locality: Caribbean Sea, off Grenada, 479 m, *Blake* Stn. 249.
 Distribution: Bahamas, Guianas, and westward into eastern Gulf of Mexico, Yucatan, and Honduras; 26°28'N, 84°42'W, 228 m; 26°20'N, 84°42'W, 216 m, from stomach of *Epinephalus flavolimbatus*. (personal communication, D. K. Camp).
 Pacific: from Borneo through the Philippines to Japan.

FAMILY RHYNCHOCINETIDAE***Rhynchocinetes rigens* Gordon, 1936**

Description: Gordon, 1936:75, figs. 1-7.-- Manning, 1961a:1, figs. 1, 2.
 Type-locality: Pontinha Bay, Madeira Island.
 Distribution: Florida; Bahamas; Virgin Islands; Bermuda; Madeira and Azores.

FAMILY GNATHOPHYLLIDAE***Gnathophyllodes mineri* Schmitt, 1933**

Description: Schmitt, 1933:7, fig. 3.
 Type-locality: Coral reefs at Ballena Point, Ensenada, Puerto Rico.
 Distribution: Southeastern Florida, Yucatan, and Caribbean Sea (Chace, 1972).

***Gnathophyllum americanum* Guérin-Méneville, 1855**

Description: Manning, 1963:58, figs. 5, 6.
 Type-locality: Cuba.
 Distribution: Bermudas, southern Florida, Gulf of Mexico, and Caribbean Sea; Canary Islands and Indo-Pacific region from the Red Sea to the Tuamotu Archipelago (Chace, 1972).

***Gnathophyllum circellum* Manning, 1963**

Description: Manning, 1963:54, figs. 3, 4.
 Type-locality: Alligator Reef, Monroe County, Florida.
 Distribution: Florida Keys and Great Exuma Island, Bahamas (Chace, 1972).

***Gnathophyllum modestum* Hay, 1917**

Description: Williams, 1984:90, fig. 62.
 Type-locality: 20 mi. SW Beaufort, North Carolina.
 Distribution: Off Beaufort, North Carolina (Williams, 1984); Florida Middle Grounds, Panama City and Biscayne Bay Florida (Dardeau et al., 1980).

FAMILY PALAEMONIDAE***Anchistioides antiguensis* (Schmitt, 1924)**

Description: Williams, 1984:78, fig. 52.
 Type-locality: English Harbor, Antigua.
 Distribution: Off Charleston, South Carolina (Wenner and Read, 1982); off west Florida through West Indies to Maranhao, Pernambuco, and Alagoas, Brazil (Coelho and Ramos, 1972); Bermuda (Williams, 1984).

***Brachycarpus biunguiculatus* (Lucas, 1849)**

Description: Williams, 1984:63, fig. 42.
 Type-locality: Oran and Bone, Algeria.
 Distribution: Virtually pantropical (Bruce, 1974); east and west American coasts, Mediterranean; West Africa; and Indo-Pacific region. Western Atlantic distribution: Cape Fear, North Carolina, western Gulf of Mexico (Ray, 1974; Pequegnat and Ray, 1974) through West Indies to Curaçao and Old Providence Island; Bermuda (Williams, 1984).

***Leander paulensis* Ortmann, 1897**

Description: Manning, 1961b:525.-- Chace, 1972: 19.
 Type-locality: São Paulo, Brazil.
 Distribution: Sandy Key Basin, in Florida Bay off Flamingo, Florida; São Paulo, Brazil.

***Leander tenuicornis* (Say, 1818)**

Description: Williams, 1984:65, fig. 43.
 Type-locality: Newfoundland Banks.
 Distribution: Tropical and subtropical waters all over world except for west coast of Americas; Newfoundland Banks (occasionally mouth of Bay of Fundy and New England; Wigley, 1970; Williams and Wigley, 1977) to

Falkland Islands in Western Atlantic (Holthuis, 1952; Bruce, 1974; Williams, 1984).

Lipkebe holthuisi Chace, 1969

Description: Chace, 1969:263, figs. 8, 9.
--Shaw et al., 1977:284, figs. 1-2.

Type-locality: Gulf of Mexico: west-northwest of Dry Tortugas, 25°13'N, 83°55'W.

Distribution: Northeastern Gulf of Mexico off Florida; Dry Tortugas; Brazil (Shaw et al., 1977).

Macrobrachium acanthurus (Wiegmann, 1836)

Description: Williams, 1984:66, fig. 44-45.

Type-locality: Brazilian coast.

Distribution: Neuse River estuary, North Carolina, to Rio Grande do Sul, Brazil (Williams, 1984).

Macrobrachium carcinus (Linnaeus, 1758)

Description: Holthuis, 1952:114, pl. 30, pl. 31: figs. a-c.

Type-locality: "in Americal fluviis" (restricted to Jamaica by Holthuis, 1952).

Distribution: Distribution of this species lies largely beyond the temperate east coast of the United States. The range extends from St. Augustine, St. Johns County, and Silver Glen Springs, Marion County, Florida, southward around the Gulf of Mexico and Caribbean Sea to Santa Catarina, Brazil (Williams, 1984).

Macrobrachium crenulatum Holthuis, 1950

Description: Holthuis, 1952:107, pl. 27: figs. a-d, pl. 28.

Type-locality: Rio Peje Bobo, Panama.

Distribution: West Indies, Panama, and Venezuela. Indian River, Ft. Pierce, Florida (personal communication, P. M. Mikkelsen).

Macrobrachium ohione (Smith, 1874)

Description: Williams, 1984:68, fig. 46.

Type-locality: Ohio River at Cannelton, Indiana.

Distribution: A narrow zone along Atlantic seaboard from James River, Hopewell, Virginia (Hobbs and Massmann, 1952), to southern Georgia; widespread from coastal Alabama to Aransas Bay, Texas; Mississippi River and tributaries upstream to McCurtain County, Oklahoma, Fort Smith, Arkansas; St. Louis, Missouri; Washington County, Ohio (Hedgpeth, 1949; Williams, 1984).

Macrobrachium olfersii (Wiegmann, 1836)

Description: Williams, 1984:70, figs. 47-48.

Type-locality: "Brazilian Coast."

Distribution: Lower Cape Fear River near Southport, North Carolina; Florida; Louisiana; Texas; Veracruz, Mexico, to Santa Catarina, Brazil. Villalobos (1969) gave a distributional map for this and related species (Williams, 1984).

Neopontonides beaufortensis Borradaile, 1920

Description: Williams, 1984:80, fig. 53.

Type-locality: Beaufort, North Carolina.

Distribution: Beaufort, North Carolina, to Grand Isle, Louisiana; Caledonia Bay, Panama; Antigua (Williams, 1984).

Palaemon floridanus Chace, 1942

Description: Holthuis, 1952:197, pl. 48: figs. a-j.

Type-locality: Captiva Island, W Florida.

Distribution: West coast of Florida.

Palaemon northropi (Rankin, 1898)

Description: Holthuis, 1952:192, pl. 47.-- Chace, 1972:21.

Type-locality: Nassau, New Providence, Bahama Islands.

Distribution: Bermudas and Florida to Estado de São Paulo, Brazil (Chace, 1972).

Palaemonetes intermedius Holthuis, 1949

Description: Williams, 1984:75, fig. 50.

Type-locality: Iron Box Bay, Chincoteague Bay, Virginia.

Distribution: Vineyard Sound, Massachusetts, to Port Aransas, Texas (Holthuis, 1952); Bahia de la Ascension, Quintana Roo, Mexico (Chace, 1972; Williams, 1984).

Palaemonetes paludosus (Gibbes, 1850)

Description: Holthuis, 1952:207, pl. 51: figs. e-j.

Type-locality: St. Andrews, Charleston County, South Carolina.

Distribution: Fresh-waters E of Alleghenies, from New Jersey to Florida. Indian River lagoon, Florida (personal communication, P. M. Mikkelsen).

Palaemonetes pugio Holthuis, 1949

Description: Williams, 1984:76, fig. 51.

Type-locality: Lagoon near Cove Point Light, Chesapeake Bay.

Distribution: Intermittent from Verte River, 3 mi. W St. Modeste (47°51'N, 69°26'W), Quebec, to near Yarmouth, Nova Scotia, Newcastle and East Brunswick, Maine (Bousfield and Laubitz, 1972; Knowlton, 1973; Williams, 1974c), Southward to Corpus Christi, Texas (Holthuis, 1952; Williams, 1984).

Palaemonetes vulgaris (Say, 1818)

Description: Williams, 1984:72, fig. 49.
Type-locality: Atlantic coast of United States.
Distribution: Southern Gulf of St. Lawrence from northern Cape Breton Island (Bousfield, 1956) through Northumberland Strait to St. Simons Inlet and Miscou Harbor near Portage Bay (Bousfield and Laubitz, 1972), southward to Cameron County, Texas (Williams, 1984).

Periclimenaeus ascidiarum Holthuis, 1951

Description: Holthuis, 1951b:80, pl. 22: figs. g-l, pl. 23: figs. a-i.
Type-locality: Bird Key Reef, Dry Tortugas, Florida.
Distribution: Colombia, 2 mi. SW of Cape la Vela. 38-40 m; Florida Middle Grounds (Hopkins et al., 1977) and Bird Key Reef, Tortugas, Florida; Dominica.

Periclimenaeus atlanticus (Rathbun, 1901)

Description: Holthuis, 1951b:83, pl. 24: figs. a-p.
Type-locality: Off St. Thomas, Virgin Islands.
Distribution: Known from type-locality, and South Florida, Isla de Cozumel, Bahía de la Ascención.

Periclimenaeus bermudensis (Armstrong, 1940)

Description: Holthuis, 1951b:107, pl. 32: figs. d-g, pl. 33: figs. a-i.
Type-locality: The Reach, St. Georges Island, Bermuda.
Distribution: Coastal waters from Bermuda, Bahamas and Dry Tortugas, Florida.

Periclimenaeus caraibicus Holthuis, 1951

Description: Holthuis, 1951b:110, pl. 32: figs. h-j, pl. 34.
Type-locality: Buccoo Reef, Tobago.
Distribution: Tobago, Barbuda, Antigua Island, Dominica, Saint Lucia Island, Bahia de la Ascension. Florida (personal communication, D. K. Camp).

Periclimenaeus chacei Abele, 1971

Description: Abele, 1971:38, figs. 1, 2.
Type-locality: Northeastern Gulf of Mexico off the west coast of Florida.
Distribution: Known from the type-locality and Indian River, Florida (Personal communication with R. H. Gore).

Periclimenaeus maxillulidens (Schmitt, 1936)

Description: Holthuis, 1951b:87, pl. 26: figs. a-o.
Type-locality: Entrance to Lac, Bonaire.
Distribution: Northeastern Gulf of Mexico and Bonaire (Chace, 1972).

Periclimenaeus pearsei (Schmitt, 1932)

Description: Holthuis, 1951b:93, pl. 28: figs. a-r.
Type-locality: Tortugas, Florida.
Distribution: Tortugas, Florida.

Periclimenaeus perlatus (Boone, 1930)

Description: Holthuis, 1951b:99, pl. 30: figs. a-l, pl. 32: fig. a.
Type-locality: Gonave Bay, Haiti.
Distribution: Dry Tortugas, Florida to Panama (Chace, 1972).

Periclimenaeus schmitti Holthuis, 1951

Description: Williams, 1984:81, fig. 54.
Type-locality: Tortugas, Florida.
Distribution: Bogue Sound, Black Rocks off New River, and Lockwoods Folly River, North Carolina; Tortugas, Florida (Williams, 1984).

Periclimenaeus wilsoni (Hay, 1917)

Description: Williams, 1984:82, fig. 55.
Type-locality: Fishing grounds, 20 mi. off Beaufort Inlet, North Carolina.
Distribution: Off Beaufort, North Carolina; off Sapelo Island, Georgia; off Loggerhead Key, near Tortugas, and Franklin County, Florida (Williams, 1984).

Periclimenes americanus (Kingsley, 1878)

Description: Williams, 1984:83, fig. 56.
Type-locality: Key West, Florida.
Distribution: Beaufort, North Carolina; to western Gulf of Mexico (Felder and Chaney, 1979), and through West Indies to Aruba; Para to São Paulo, Brazil (Coelho and Ramos, 1972; Williams, 1984).

***Periclimenes harringtoni* Lebour, 1949**

Description: Holthuis, 1951b:35, pl. 9: figs. a-k.

Type-locality: Harrington Sound, Bermuda.
Distribution: Harrington Sound, Bermuda, and Dry Tortugas, Florida.

***Periclimenes iridescent* Lebour, 1949**

Description: Williams, 1984:85, fig. 57.

Type-locality: Off Castle Roads, Bermuda.
Distribution: Northeast off Cape Hatteras, 35°32.9'N, 75°11.9'W (Herbst et al. 1979); southern and northwestern Florida; Tobago; Cubagua Island, Venezuela; Bermuda (Chace 1972; Williams, 1984).

***Periclimenes longicaudatus* (Stimpson, 1860)**

Description: Williams, 1984:86, fig. 58.

Type-locality: Coast of Carolina.

Distribution: Cape Hatteras, North Carolina, to southwestern Florida; West Indies to São Paulo, Brazil. There are doubtful records from the Indian Ocean and deeper waters of the Gulf of Mexico (Holthuis, 1951b; Williams, 1984).

***Periclimenes magnus* Holthuis, 1951**

Description: Holthuis, 1951b:52, pl. 15: figs. a-f.

Type-locality: Gulf of Mexico, off Aransas, Texas, 27°40'N, 96°34'W, 50 m, *Pelican* Stn. 42.
Distribution: Type-locality and Murray Key, Florida Bay (Rouse, 1970).

***Periclimenes pandionis* Holthuis, 1951**

Description: Holthuis, 1951b:41, pl. 11: figs. a-i.--Gore et al., 1981:254, fig. 1.

Type-locality: Gulf stream off Key West, Florida, 24°21'55"N, 81°58'25"W (179 m. *Fish Hawk* Stn. 7279).

Distribution: Indian River and Key West, Florida (Gore et al., 1981).

***Periclimenes pedersoni* Chace, 1958**

Description: Williams, 1984:87, fig. 59.

Type-locality: Simms (Lyford) Cay, New Providence Island, Bahamas.

Distribution: East of Cape Lookout, North Carolina, 34°35.5'N, 75°5.5'W (Herbst et al. 1979); off northwest Florida, Bahamas, through West Indies to Bonaire; Belize (Williams, 1984).

***Periclimenes perryae* Chace, 1942**

Description: Holthuis, 1951b:31, pl. 7: figs. a-o.

Type-locality: Off Sanibel Island, Lee County,

W Florida.

Distribution: Florida Middle Grounds (Hopkins et al., 1977) and from shallow water (10 m) off Sanibel Island, Lee County, West Florida.

***Periclimenes rathbunae* Schmitt, 1924**

Description: Holthuis, 1951b:58, pl. 17: figs. a-h.

Type-locality: Spanish Port, Curaçao.

Distribution: Netherlands West Indies, Spanish Port, Curaçao; tentatively Dry Tortugas, Florida.

***Periclimenes yucatanicus* (Ives, 1891)**

Description: Limbaugh, Pederson, and Chace, 1961:240, fig. 2.

Type-locality: Off Progreso, Estado de Yucatan, Mexico.

Distribution: Southern Florida to Colombia (Chace, 1972).

***Pontonia domestica* Gibbes, 1850**

Description: Williams, 1984:88, fig. 60.

Type-locality: South Carolina.

Distribution: Atlantic Beach near Beaufort Inlet, North Carolina, to Gulf of Mexico S of Houma, Terrebonne Parish, Louisiana (USNM); Bahamas; Madeira (Williams, 1984).

***Pontonia margarita* Smith, 1869**

Description: Williams, 1984:89, fig. 61.

Type-locality: Bay of Panama.

Distribution: Atlantic coast: Drumm Inlet to Beaufort Inlet, North Carolina; east and west Florida. Pacific coast: Gulf of California to Colombia; Galapagos Islands (Williams, 1984).

***Pontonia unidens* Kingsley, 1880**

Description: Holthuis, 1951b:150, pl. 47: figs. J-k.

Type-locality: Key West, Florida.

Distribution: Known only from the original type-series from Key West, Florida.

***Pontoniopsis paulae* Gore, 1981**

Description: Gore, 1981:139, figs. 1a-p.

Type-locality: Carys Fort Reef, off Key Largo, Monroe county, Florida.

Distribution: Known only from the type-locality.

***Pseudocoutierea antillensis* Chace, 1972**

Description: Chace, 1972:43, figs. 11a-b.

Type-locality: Saba Bank at 17°28'N, 63°13'W.

Distribution: Known from the type-locality; eastern Florida.

***Tuleariocaris neglecta* Chace, 1969**

Description: Chace, 1969:266, figs. 10, 11.

Type-locality: St. James, Barbados.

Distribution: Florida Keys, Dominica, Barbados, Curaçao, Madeira, on *Diadema antillarum* (Chace, 1972).***Typton carneus* Holthuis, 1951**

Description: Chace, 1972:46, fig. 12.

Type-locality: Tortugas, Florida.

Distribution: South and west coasts of Florida and Bahamas to Tobago (Chace, 1972).

***Typton distinctus* Chace, 1972**

Description: Chace, 1972:49, figs. 13, 14.

Type-locality: Los Arroyos, Provincia de Pinar del Rio, Cuba.

Distribution: Western Cuba and Yucatan Peninsula, Mexico (Chace, 1972). Florida (personal communication, D. A. Camp).

***Typton gnathophylloides* Holthuis, 1951**

Description: Holthuis, 1951b:159, pl. 50: figs. a-l.

Type-locality: Dry Tortugas, Florida.

Distribution: Same as the type-locality.

***Typton prionurus* Holthuis, 1951**

Description: Holthuis, 1951b:165, pl. 52: figs. a-l.

Type-locality: Tortugas, Florida between Middle Ground and White Shoal, 18 m. (W. L. Schmitt coll., Stn 45-30).

Distribution: Same as the type-locality.

***Typton tortugae* McClendon, 1911**

Description: Holthuis, 1951b:153, pl. 48: figs. a-o.

Type-locality: Dry Tortugas, Florida.

Distribution: Outside Castle Harbour, Bermuda (Gurney, 1936), Dry Tortugas, Florida (McClendon, 1911; Schmitt, 1930; Pearse, 1932); Gulf of California (Holthuis, 1951).

***Typton vulcanus* Holthuis, 1951**

Description: Holthuis, 1951b:157, pl. 49: figs. a-n.

Type-locality: Dry Tortugas, Florida.

Distribution: South of Tortugas, Florida.

***Veleroniopsis kimallynae* Gore, 1981**

Description: Gore, 1981:147.

Type-locality: Elbow Reef, off Key Largo, Monroe County, Florida.

Distribution: Known only from the type-locality.

FAMILY ALPHEIDAE***Alpheopsis labis* Chace, 1972**

Description: Chace, 1972:55, fig. 15.

Type-locality: Charlotte Point, English Harbor, Antigua Island.

Distribution: Bermuda, Cuba, Hispaniola and Antigua Island (Chace, 1972). Florida Middle Grounds (Hopkins et al., 1977); Florida. (27°37'N, 83°59'W, 54.8 m) (personal communication, D. K. Camp). Looe Key, Florida (personal communication, D. L. Felder).

***Alpheopsis trispinosus* (Stimpson, 1861)**

Description: Banner and Banner, 1973:337, fig. 14.

Type-locality: Port Jackson, Australia.

Distribution: Probably pantropical (Gore, 1981).

***Alpheus amblyonyx* Chace, 1972**

Description: Chace, 1972:59, fig. 16.

Type-locality: Near center of Arrecife Nicchehabin, Bahia de la Ascension, Territorio de Quintana Roo, Mexico, on or under coral in 1-5 feet of water.

Distribution: Territorio de Quintana Roo, Mexico; Puerto Rico; Saint Thomas; and Dominica; (Chace, 1972). Eastern and Southern Florida (personal communication, P. M. Mikkelsen).

***Alpheus armatus* Rathbun, 1901**Description: Zimmer, 1913:295, figs. W-Z, A¹, B¹.

Type-locality: Ponce, Puerto Rico.

Distribution: Throughout the West Indian region from the Bahamas and southern Florida to Tobago and westward to the Yucatan Peninsula (Chace, 1972).

***Alpheus armillatus* H. Milne Edwards, 1837**

Description: Williams, 1984:92, fig. 63.

Type-locality: Antilles.

Distribution: North Carolina, through Gulf of

Mexico and West Indies to Cananeia, São Paulo, Brazil; Bermuda (Holthuis, 1956).

Alpheus bouvieri A. Milne Edwards, 1878
Description: Crosnier and Forest, 1966:273, fig. 22.

Type-locality: Cape Verde Islands.

Distribution: Bermudas and Antigua Island to Tobago and Fernando de Noronha; eastern Atlantic from the Cape Verde Islands and Guinea to São Tomé and Congo (Chace, 1972). Eastern and Southern Florida.

Alpheus candei Guérin-Méneville, 1855

Description: Coutière, 1910:486, fig. 1.

Type-locality: Cuba.

Distribution: Apparently known only from the Dry Tortugas, Florida and Cuba (Chace, 1972).

Alpheus cristulifrons Rathbun, 1900

Description: Crosnier and Forest, 1966:260, figs. 17, 18.

Type-locality: Fernando de Noronha.

Distribution: Western tropical Atlantic from the Dry Tortugas, Florida to Fernando de Noronha and westward to the Yucatan Peninsula; also the islands of São Tomé and Príncipe in the eastern Atlantic (Chace, 1972).

Alpheus cylindricus Kingsley, 1878

Description: Crosnier and Forest, 1966:257, fig. 16.

Type-locality: Archipelago de las Perlas, Gulf of Panama.

Distribution: Bermudas and Florida to Barbados; eastern Atlantic from the islands of Príncipe, São Tomé, and Annobon; eastern Pacific from the Gulf of California, the Gulf of Panama, and the Galapagos Islands (Chace, 1972).

Alpheus estuariensis Christoffersen, 1984

Description: Christoffersen, 1984:191, figs. 1, 2.

Type-locality: Rio Potengi estuary, Natal, State of Rio Grande do Norte, Brazil.

Distribution: Florida; Mississippi to Texas; Cuba; Dominican Republic; Curaçao; Trinidad; Ceara to Parana, Brazil.

Alpheus floridanus Kingsley, 1878

Description: Chace, 1972:65, figs. 17-20.-- Crosnier and Forest, 1966:267, 269, figs. 20, 21.

Type-locality: Fort Jefferson, Dry Tortugas, Florida.

Distribution: Gulf of Mexico to Estado da Bahia, Brazil; eastern Atlantic from Guinea to Congo (Chace, 1972).

Alpheus formosus Gibbes, 1850

Description: Williams, 1984:94, fig. 64.

Type-locality: Key West, Florida.

Distribution: Near Beaufort, North Carolina through Gulf of Mexico (Ray, 1974; Felder and Chaney, 1979) and West Indies to São Paulo, Brazil.

Alpheus heterochaelis Say, 1818

Description: Williams, 1984:95, fig. 65.--

Christoffersen, 1984:200, figs. 5-7.

Type-locality: Amelia Island, Nassau County, Florida.

Distribution: North Carolina to the State of Paraíba, Brazil (Christoffersen, 1984).

Alpheus malleator Dana, 1852

Description: Crosnier and Forest, 1966:240, fig. 10.

Type-locality: Rio de Janeiro, Brazil?

Distribution: Puerto Rico to Estado de São Paulo, Brazil; eastern Atlantic from Senegal to Congo; eastern Pacific from the Gulf of California, Ecuador, and the Galapagos Islands (Chace, 1972).

Alpheus normanni Kingsley, 1878

Description: Williams, 1984:97, fig. 66.

Type-locality: Pacific coast of Panama.

Distribution: Bermuda; around Cape Charles, Virginia, and lower Chesapeake Bay through Gulf of Mexico (Ray, 1974) and West Indies to São Paulo, Brazil (Christoffersen, 1979); Gulf of California and Panama (Chace, 1972; Williams, 1984).

Alpheus nuttingi (Schmitt, 1924)

Description: Schmitt, 1924b:78, pl. 2: figs. 4-6.

Type-locality: Pelican Island, Barbados.

Distribution: Florida Keys to Estado de Alagoas, Brazil and westward to Isla de Providencia and Panama (Chace, 1972).

Alpheus paracrinitus Miers, 1881

Description: Crosnier and Forest, 1966:253, fig. 15.

Type-locality: Goree, Senegal.

Distribution: Virtually pantropical; to a depth of 18 m. In Western Atlantic, from the Bermudas and the northeastern Gulf of Mexico to Tobago (Chace, 1972).

***Alpheus peasei* (Armstrong, 1940)**

Description: Verrill, 1922:68, fig. 68, fig. 5b, pl. 19: figs. 3a-d, pl. 20: fig. 1, pl. 21: figs. 6, 6a, pl. 24: figs. 2-4, pl. 29: fig. 1a-t.

Type-locality: Castle Harbour, Bermudas.

Distribution: Bermudas and Florida Keys to Tobago and westward to Isla de Providencia and the Yucatan Peninsula (Chace, 1972).

***Alpheus schmitti* Chace, 1972**

Description: Chace, 1972:70, figs. 21, 22.

Type-locality: Grand Anse Bay outside Saint Georges Harbour, Grenada, in partially exposed conglomerate rock and coral ledge along shore.

Distribution: Known from the type-series, the Florida Keys, Antigua Island, and Tobago (Chace, 1972).

***Alpheus thomasi* Hendrix and Gore, 1973**

Description: Hendrix and Gore, 1973:413, figs. 1-3.

Type-locality: Virginia Beach, Virginia Key, Miami, Florida.

Distribution: Cape Florida, Key Biscayne, Dade County; and from Jupiter Inlet in Palm Beach County, and Walton Rocks, St. Lucie County, about 5 miles south of Ft. Pierce, Florida (Hendrix and Gore, 1973).

***Alpheus viridari* (Armstrong, 1949)**

Description: Armstrong, 1949:8, fig. 2.

Type-locality: Barahona, Dominican Republic.

Distribution: Florida Keys to Trinidad and westward to Curaçao and the Yucatan Peninsula (Chace, 1972).

***Alpheus websteri* Kingsley, 1880**

Description: Rankin, 1898:249, pl. 30: fig. 6 (as *Alpheus nigrospinatus*).--Crosnier and Forest, 1966:236 (as *Alpheus ridleyi*).

Type-locality: Key West, Florida.

Distribution: Bahamas to Fernando de Noronha, Brazil and westward to the Yucatan peninsula (Chace, 1972). Looe Key, Florida (personal communication, D. L. Felder).

***Automate evermanni* Rathbun, 1901**

Description: Williams, 1984:99, fig. 67.

Type-locality: Off Aguadilla, Puerto Rico.

Distribution: North Carolina(?); Georgia to Texas and Puerto Rico; eastern Atlantic from Cape Verde Islands and Liberia to Nigeria (Chace, 1972).

***Automate gardineri* Coutière, 1902**

Description: Banner and Banner, 1966:17, fig. 8.--Chace, 1972:74, fig. 23.--Williams, 1984:100, fig. 68.

Type-locality: Maldives and Laccadive Islands.

Distribution: North Carolina; Virgin Islands, Antigua Island, Barbados and the Yucatan Peninsula; Indo-Pacific region from the Red Sea to Samoa (Chace, 1972). Florida.

***Automate rectifrons* Chace, 1972**

Description: Chace, 1972:75, fig. 24.

Type-locality: Inner side of Arrecife

Nicchehabin, Bahia de la Ascension, Territorio de Quintana Roo, Mexico.

Distribution: The type-locality; possibly Antigua Island (Chace, 1972). Southern Florida (personal communication, P. A. McLaughlin).

***Leptalphes forceps* Williams, 1965**

Description: Williams, 1984:101, fig. 69.

Type-locality: Gallants Point, Newport River, Carteret County, North Carolina.

Distribution: Drum Inlet, Beaufort, Banks Channel near Wrightsville Beach, and Lockwoods Folly Inlet, North Carolina; Old Tampa Bay, Florida (Saloman, 1971; Simon and Dauer, 1977); Davis Bayou, Mississippi (Dawson, 1967a; Williams, 1984).

***Metalpheus rostratipes* (Pocock, 1890)**

Description: Crosnier and Forest, 1966:246, figs. 12-14.

Type-locality: Fernando de Noronha.

Distribution: Southern Florida; Puerto Rico and the Yucatan Peninsula to Fernando de Noronha; probably pantropical (Chace, 1972).

***Synalphes agelas* Pequegnat and Heard, 1979**

Description: Pequegnat and Heard, 1979:110, figs. 1-4.--Dardeau, 1984:12, figs. 3-6.

Type-locality: West Flower Garden Bank, Gulf of Mexico (27°52'N, 93°48'W) in 25 m.

Distribution: Atlantic: Grand Bahama Island; Gulf of Mexico: Florida Middle Ground, Sonnier Bank, 28 Fathom Bank, West Flower Garden Bank, North Hospital Bank and Hospital Rock (Pequegnat and Heard, 1979); Caribbean, off Puerto Rico (Dardeau, 1984).

***Synalphes apioceros* Coutière, 1909**

Description: Coutière, 1909:27, fig. 9.

Type-locality: Marco, Florida.

Distribution: Southern Florida to Surinam

westward to the Yucatan Peninsula (Chace, 1972).

Synalpheus bousfieldi Chace, 1972

Description: Chace, 1972:86, figs. 29, 30.

Type-locality: West side of reef east of anchorage, Bahía del Espíritu Santo, Territorio de Quintana Roo, Mexico.

Distribution: Atlantic: Grand Bahama Island (Dardeau, 1984) and possibly south to Brazil (Christoffersen, 1979); Gulf of Mexico: Florida Middle Ground, Sonnier Bank, Bright Bank, and West Flower Garden Bank (Dardeau, 1984); Caribbean: Yucatan Peninsula and Virgin Islands (Chace, 1972).

Synalpheus brevicarpus (Herrick, 1891)

Description: Coutière, 1909:50, fig. 29.--

Christoffersen, 1979:333, fig. 19.

Type-locality: Nassau, New Providence, Bahamas; in green sponge.

Distribution: Bermudas; east Florida to Dry Tortugas; southwest Florida; Bahama Islands; Cuba to Virgin Islands; Los Roques Islands; Curaçao; Panama; Pernambuco to the north of Rio Grande do Sul; eastern Pacific, Bay of Panama (Christoffersen, 1979).

Synalpheus brooksi Coutière, 1909

Description: Coutière, 1909:69, fig. 41.--

Dardeau, 1984:26, figs. 11-14.

Type-locality: Sugar Loaf Key, Florida.

Distribution: Gulf of Mexico, Florida Keys, Bahamas and the Yucatan Peninsula to Estado do Rio Grande do Norte Brazil (Chace, 1972).

Synalpheus curacaoensis Schmitt, 1924

Description: Schmitt, 1924a:66, fig. 3.

Type-locality: Spaansche Water, Curaçao.

Distribution: Curaçao; Bonaire, (Chace, 1972). Eastern Florida.

Synalpheus fritzmuelleri Coutière, 1909

Description: Williams, 1984:102, fig. 70.

Type-locality: Marco, Florida.

Distribution: Off Beaufort, North Carolina to Santa Catarina, Brazil; Bermuda; St. Helena Island, South Atlantic; Baja California (Chace, 1972).

Synalpheus goodei Coutière, 1909

Description: Coutière, 1909:58, fig. 33.--

Dardeau, 1984:40, figs. 18-21.

Type-locality: Bermudas.

Distribution: Bermudas and the Gulf of

Mexico to Curaçao and Panama (Chace, 1972).

Synalpheus heardi Dardeau, 1984

Description: Dardeau, 1984:47, figs. 23-26.

Type-locality: Florida Middle Ground, Gulf of Mexico.

Distribution: Eastern Gulf of Mexico, off central western Florida; Grand Bahama Island (Dardeau, 1984).

Synalpheus hemphilli Coutière, 1909

Description: Coutière, 1909:38, fig. 20.

Type-locality: Off the west coast of Florida (27°04'N, 83°21'W).

Distribution: Bermudas and the eastern Gulf of Mexico to Curaçao and Islas Los Roques (Chace, 1972).

Synalpheus herricki Coutière, 1909

Description: Coutière, 1909:74, fig. 44.--

Dardeau, 1984:55, figs. 27-30.

Type-locality: "Off Anclote, Florida" presumably Anclote Key off Tarpon Springs.

Distribution: Atlantic: questionably from Eleuthera, Bahama Islands (Chace, 1972); Gulf of Mexico: questionably from Florida Bay (Tabb and Manning, 1961); off central western Florida from Sanibel Island to Cape San Blas (Coutière, 1909; Dardeau, 1984).

Synalpheus longicarpus (Herrick, 1891)

Description: Williams, 1984:104, fig. 71.--

Dardeau, 1984:64, figs. 32-35.

Type-locality: Bahamas, probably Nassau, New Providence Island.

Distribution: Beaufort, North Carolina to west Flower Garden Reef, SSE of Galveston, Texas; Yucatan, Mexico through West Indies to Rio de Janeiro, Brazil (Williams, 1984).

Synalpheus mcclendoni Coutière, 1910

Description: Chace, 1972:95, figs. 33, 34.--

Dardeau, 1984:74, figs. 37-39.

Type-locality: Dry Tortugas, Florida.

Distribution: Atlantic: Grand Bahama Island; Gulf of Mexico: Dry Tortugas (Coutière, 1910) and Isla de Lobos Reef, off Veracruz (Ray, 1974); Caribbean: Yucatan Peninsula, Windward Islands (Chace, 1972) and Barbados (Schmitt, 1924a; Dardeau, 1984).

Synalpheus minus (Say, 1818)

Description: Williams, 1984:105, fig. 72.

Type-locality: "Coasts of the southern states and off East Florida."

Distribution: Near Cape Hatteras, North Carolina to São Paulo, Brazil (Christoffersen, 1979); Bermuda (Williams, 1984).

***Synalpheus pandionis* Coutière, 1909**

Description: Coutière, 1909:67, fig. 39.--

Dardeau, 1984:78, figs. 40-43.

Type-locality: Off Saint Thomas, 36-42 m.

Distribution: Eastern Gulf of Mexico to Barbados and Curaçao (Chace, 1972).

***Synalpheus paranepetunus* Coutière, 1909**

Description: Coutière, 1909:86, fig. 52.--

Dardeau, 1984:92, figs. 47-50.

Type-locality: Off Golfo de Morrosquillo, Colombia (9°30'15"N, 76°20'30" W) in 77 m.

Distribution: Dry Tortugas, Florida and the Yucatan Peninsula to the Grenadines and Colombia (Chace, 1972). Possibly from the Gulf of Mexico, off central western Florida, and the West Flower Garden Bank (Dardeau, 1984).

***Synalpheus pectiniger* Coutière, 1907**

Description: Coutière, 1909:78, figs. 48, 49.--

Dardeau, 1984:98, figs. 51-53.

Type-locality: Curaçao.

Distribution: Gulf of Mexico, Florida keys and Bahamas to Curaçao (Chace, 1972).

***Synalpheus rathbunae* Coutière, 1909**

Description: Coutière, 1909:84, fig. 51.

Type-locality: Off Saint Thomas in 37-55 m.

Distribution: Bahamas to the Grenadines westward to the Yucatan Peninsula; (Chace, 1972). Looe Key, Florida (personal communication, D. L. Felder).

***Synalpheus sanctithomae* Coutière, 1909**

Description: Coutière, 1909:61, fig. 35.--

Christoffersen, 1979:352, figs. 29-30.

Type-locality: Off Saint Thomas in 37-42 m.

Distribution: Saint Thomas; Atol dos Rocas and Pernambuco to the south of Bahia, Brazil (Christoffersen, 1979). Florida Keys (Gore, 1981).

***Synalpheus townsendi* Coutière, 1909**

Description: Williams, 1984:106, fig. 73.

Type-locality: Gulf of Mexico south of Cape San Blas, Florida (29°14'00"N, 85°29'15"W) in 46 m.

Distribution: Off Beaufort, North Carolina to Rio de Janeiro, Brazil; Bermuda; Gulf of California (Williams, 1984)

***Thunor simus* (Guérin-Méneville, 1856)**

Description: Armstrong, 1949:13, figs. 3, 4A-

J, L, (as *Thunor rathbunae*).--Chace,

1972:104, fig. 39, (as *Thunor rathbunae*).

Type-locality: Cuba.

Distribution: Key West, Florida and Yucatan

Peninsula to Barbados (Chace, 1972);

Piscadera Bay, Curaçao (Holthuis, 1980a).

FAMILY HIPPOLYTIDAE

***Bythocaris nana* Smith, 1885**

Description: Smith, 1885:449; 1886:600, pl. 12: fig. 2.

Type-locality: Massachusetts, off Martha's Vineyard; 263 m.

Distribution: Off Martha's Vineyard, Massachusetts to Southern Florida and northeastern Gulf of Mexico.

***Exhippolysmata oplophoroides* (Holthuis, 1948)**

Description: Williams, 1984:114, fig. 79.

Type-locality: Mouth of Suriname River near Resolutie, Surinam.

Distribution: Off Cape Fear River, North Carolina, to Port Aransas, Texas; Guyana to the north of Uruguay (Williams, 1984).

***Hippolyte coerulescens* (Fabricius, 1775)**

Description: Williams, 1984:116, fig. 80.

Type-locality: "Pelago inter Tropicos"

Distribution: Widespread in tropical and subtropical Atlantic Ocean (Chace 1972).

***Hippolyte curacaoensis* Schmitt, 1924**

Description: Williams, 1984:117, fig. 81.

Type-locality: West Punt, Curaçao.

Distribution: Beaufort and Sneads Ferry, North Carolina; West Indies from Cuba to Curaçao (Williams, 1984).

***Hippolyte nicholsoni* Chace, 1972**

Description: Chace, 1972:113, figs. 46, 47.

Type-locality: Milford Bay, between Pigeon Point and Crown Point, Tobago, in 9-12 m.

Distribution: Puerto Rico, Antigua Island, Saint Lucia Island, and Tobago (Chace, 1972). Looe Key, Florida (personal communication, D. L. Felder).

***Hippolyte pleuracanthus* (Stimpson, 1871)**

Description: Williams, 1984:117, fig. 82.

Type-locality: Norfolk Harbor, Virginia, and Somers Point, Great Egg Harbor, New Jersey.

Distribution: Connecticut to North Carolina (Williams, 1984).

Hippolyte zostericola (Smith, 1873)

Description: Chace, 1972:118, figs. 49-50.

Type-locality: Vineyard Sound, Massachusetts.

Distribution: Southern Massachusetts; North Carolina to Yucatan; Trinidad and Curaçao; Ceara, Brazil (Fausto-Filho, 1975); Bermuda.

Latreutes fucorum (Fabricius, 1798)

Description: Williams, 1984:119, fig. 84.

Type-locality: Floating gulfweed.

Distribution: Western North Atlantic between 10° and 50° N; Azores and Cape Verde Islands (Chace, 1972).

Latreutes parvulus (Stimpson, 1866)

Description: Williams, 1984:120, fig. 85.

Type-locality: St. Joseph Island, Texas.

Distribution: Beaufort, North Carolina, to Rio de Janeiro, Brazil; West Africa (Williams, 1984).

Lysmata amboinensis (De Man, 1888)

Description: Limbaugh, Pederson, and Chace, 1961:247, fig. 5.--Hayashi, 1975:285, figs. 1-4, pl. 4.

Type-locality: Amboina, Indonesia.

Distribution: Circumtropical (Bruce, 1974; Hayashi, 1975). In Florida: northern Gulf of Mexico through the Florida Keys.

Lysmata intermedia (Kingsley, 1878)

Description: Sivertsen, 1933:5, pl. 2: fig. 9-15.

Type-locality: Dry Tortugas, Florida.

Distribution: The Florida Middle Grounds (Dardeau et al., 1980) to the Florida Keys to Tobago and Curaçao; Azores; Galapagos Islands (Chace, 1972). The Galapagos Islands record is questionable (Abele, 1975).

Lysmata rathbunae Chace, 1970

Description: Williams, 1984:126, fig. 89.

Type-locality: Off Boynton Beach, Florida, 26°31'N, 80°1'W, 55-64 m.

Distribution: Typical form: SE Cape Fear, North Carolina, 33°30'24"N, 77°24'30"W, 25 m; east coast of Florida to Yucatan. Bermuda; Miami, Florida, Venezuela (Williams, 1984).

Lysmata wurdemanni (Gibbes, 1850)

Description: Williams, 1984:127, fig. 90.

Type-locality: Key West, Florida.

Distribution: Great Egg Harbor, New Jersey,

to Port Aransas, Texas; Surinam; French Guiana; Mamanguape, São Paulo, Brazil (Williams, 1984).

Merhippolyte americana Holthuis, 1961

Description: Holthuis, 1961:1, fig. 1.

Type-locality: 20°59'30"N, 86°23'45"W, Yucatan Channel, 237.6 m, coral.

Distribution: North Carolina; South Florida; Yucatan Channel; São Paulo; Rio Grande do Sul to Province of Buenos Aires (Christoffersen, 1979).

Thor amboinensis (De Man, 1888)

Description: Chace, 1972:130, figs. 55, 56.

Type-locality: Ambon, Indonesia.

Distribution: Florida Keys to Tobago and Yucatan; Bay of Bengal, Indonesia, and Caroline Islands (Chace, 1972).

Thor dobkini Chace, 1972

Description: Williams, 1984:134, fig. 94.

Type-locality: Punta Rassa (near mouth of Caloosahatchee River), Lee County, Florida.

Distribution: Off Shackleford Bank, North Carolina, to Yucatan; Louisiana; north coast of Cuba (Williams, 1984).

Thor floridanus Kingsley, 1878

Description: Williams, 1984:135, fig. 95.

Type-locality: Key West, Florida.

Distribution: Black Rocks off New River, North Carolina (?) to Yucatan (Williams, 1984).

Thor manningi Chace, 1972

Description: Williams, 1984:137, fig. 96.

Type-locality: English Harbour, Antigua Island.

Distribution: Beaufort, North Carolina, to Yucatan and through West Indies to Curaçao; Islas Tres Marias, off west coast of Mexico (Chace 1972).

Tozeuma carolinense Kingsley, 1878

Description: Williams, 1984:138, fig. 97.

Type-locality: Fort Macon, North Carolina.

Distribution: Vineyard Sound, Massachusetts, through Gulf of Mexico to Yucatan and southward to Colon, Panama; through West Indies to Curaçao; Pernambuco to Bahia, Brazil (Coelho and Ramos 1972).

Tozeuma cornutum A. Milne Edwards, 1881

Description: A. Milne Edwards, 1881:16;

1883, pl. 32.--Chace, 1972:141.

Type-locality: Off Barbados.

Distribution: Off Barbados, in 73 m; in deep water east of Florida Keys; in Great Lameshur Bay, St. John, Virgin Islands (Chace, 1972).

Tozeuma serratum A. Milne Edwards, 1881

Description: Williams, 1984:140, fig. 98.

Type-locality: Off Barbados.

Distribution: Nonamesset Island, Massachusetts; Off Capes Hatteras and Lookout, North Carolina (Herbst et al., 1979); Cape Canaveral, extreme southern and northwestern Florida, Colombia and Barbados (Chace, 1972; Williams, 1984).

Trachycaris restrictus (A. Milne Edwards, 1878)

Description: Holthuis, 1949b:233, figs. 2, 3.

Type-locality: Cape Verde Islands.

Distribution: Bermudas; Cape San Blas, Florida (Dardeau et al., 1980) south to Estado do Para, Brazil; eastern Atlantic from the Canary Islands to St. Helena Island (Chace, 1972).

FAMILY OGYRIDIDAE

Ogyrides alphaerostris (Kingsley, 1880)

Description: Williams, 1984:107, fig. 74.

Type-locality: Northampton County, Virginia, eastern shore, Atlantic side.

Distribution: Eastern shore of Accomack County, and lower James River, Virginia, through Gulf of Mexico to Rio Grande do Sul, Brazil (Christoffersen 1979). Florida (26 m) (personal communication, P. M. Mikkelsen).

Ogyrides hayi Williams, 1981

Description: Williams, 1984:109, fig. 75.

Type-locality: Off Bogue Bank west of Ft. Macon, North Carolina, ~ 3.5 m.

Distribution: Beaufort, North Carolina, to Sebastian Inlet, Florida; northwestern Florida to Mississippi; Puerto Rico (Williams, 1984).

FAMILY PROCESSIDAE

Ambidexter symmetricus Manning and Chace, 1971

Description: Manning and Chace, 1971:3, figs. 1, 2.

Type-locality: Matheson Hammock Wading Beach, Biscayne Bay, Miami, Florida.

Distribution: Gulf of Mexico to Trinidad (Chace, 1972).

Nikoides schmitti Manning and Chace, 1971

Description: Williams, 1984:141, fig. 99.

Type-locality: 1.25 km south of Garden Key, Tortugas, Monroe County, Florida.

Distribution: East of Cape Lookout, North Carolina (Herbst et al., 1979), Biscayne Bay and Dry Tortugas; Guadeloupe and the Guianas (Williams, 1984).

Processa bermudensis (Rankin, 1900)

Description: Williams, 1984:143, fig. 100.

Type-locality: Harrington Sound, Bermuda.

Distribution: Bermuda; North Carolina near Cape Hatteras to northwestern Florida; Veracruz, Mexico (Ray, 1974); Cuba; Puerto Rico; Guadeloupe; Peninsula de Arago, Estado Sucre, Venezuela, in *Sargassum*; Bahia and Rio de Janeiro, Brazil (Christoffersen, 1979; Williams, 1984).

Processa fimbriata Manning and Chace, 1971

Description: Williams, 1984:144, fig. 101.

Type-locality: Off East Key, Tortugas, Monroe County, Florida.

Distribution: Off New River, North Carolina, to Rio de Janeiro, Brazil (Christoffersen 1979).

Processa guyanae Holthuis, 1959

Description: Williams, 1984:145, fig. 102.

Type-locality: NW of the Coppename River (Surinam) 6°54'N, 56°14'W, 49 m.

Distribution: Off Cape Hatteras, North Carolina, to eastern Gulf of Mexico, including northern coast of Cuba; Surinam, Ceara, Brazil, to Uruguay (Williams, 1984). Florida (40-200 m) (personal communication, P. M. Mikkelsen).

Processa hemphilli Manning and Chace, 1971

Description: Williams, 1984:146, fig. 103.

Type-locality: Marco, Collier County, Florida.

Distribution: E Cape Lookout, and Bogue Sound, North Carolina; E coast of Florida; NW Florida (Saloman, 1979); Guadeloupe; Rio de Janeiro, Brazil, to Province of Buenos Aires, Argentina (Christoffersen, 1979; Williams, 1984).

Processa profunda Manning and Chace, 1971

Description: Williams, 1984:147, fig. 104.

Type-locality: Gulf of Mexico off the west coast of Florida, 202 m.

Distribution: Southeast of Cape Hatteras; off South Carolina; Gulf of Mexico off southern

and western Florida; Surinam (Williams, 1984).

Processa riveroi Manning and Chace, 1971
Description: Manning and Chace, 1971:28, fig. 16.

Type-locality: Maguey Island, La Parguera, Puerto Rico.

Distribution: Puerto Rico. Looe Key, Florida (personal communication, D. L. Felder).

Processa vicina Manning and Chace, 1971

Description: Williams, 1984:148, fig. 105.

Type-locality: Off North Carolina, 34°35'30"N, 75°45'30"W, 59 m.

Distribution: SE Cape Lookout, North Carolina, northwest Florida, off Isla Margarita, Venezuela (Williams, 1984).

FAMILY PANDALIDAE

Pantomus parvulus A. Milne Edwards, 1883

Description: Williams, 1984:157, fig. 110.

Type-locality: Northern part of Yucatan Bank, 23°13'N, 89°16'W, 153.6 m.

Distribution: Cape Lookout, North Carolina, to Yucatan, Mexico; Puerto Rico; St. Croix, Virgin Island; Surinam (Williams, 1984).

Plesionika acanthonotus (Smith, 1882)

Description: Holthuis, 1951a:62, figs. 13b-t.--Pequegnat, 1970:91.

Type-locality: Off South Carolina, 32°43' N, 77°21' W, 426 m. *Blake* Stn. 321.

Distribution: Western Atlantic: from off South Carolina to off southern Florida and off Nicaragua and Brazil, NE and NW Gulf of Mexico. Eastern Atlantic: off Portugal and Spain in the north; and off Angola and Rio Mundi, Africa, in the south and in the Mediterranean (Pequegnat, 1970).

Plesionika edwardsii (Brandt, 1851)

Description: Chace, 1985:62, fig. 26.

Type-locality: Unknown.

Distribution: In the western and eastern Atlantic (including the Gulf of Mexico and the Mediterranean) and in the IndoPacific region (Chace, 1985).

Plesionika ensis (A. Milne Edwards, 1881)

Description: Holthuis, 1951a:55, fig. 11.

Type-locality: Off Barbados, 434m. *Blake* Stn. 283.

Distribution: Western Atlantic: east coast of Florida, Florida Straits and eastern Gulf of

Mexico, off Barbados, and NE Brazil. Eastern Atlantic: off Morocco and the Gulf of Guinea. Indian Ocean near Andaman Island. Pacific: Hawaiian Islands (Pequegnat, 1970).

Plesionika escatilis (Stimpson, 1860)

Description: Crosnier and Forest, 1973:221, fig. 69a.

Type-locality: Madeira Islands, Eastern Atlantic.

Distribution: The exact status and range of this species is yet to be determined. It occurs in the eastern and western Atlantic regions. Mesopelagic.

Plesionika longicauda (Rathbun, 1901)

Description: Rathbun, 1901:117, fig. 24.--Pequegnat, 1970:86.

Type-locality: Northeast Gulf of Mexico, 161 m, *Albatross* Ocean Stn. 2403, 28°42.5'N, 85°29'W.

Distribution: Indigenous to SW North Atlantic: NE Gulf of Mexico, Florida straits, Bahamas, off Puerto Rico, in the Caribbean off Honduras, and off Surinam (Pequegnat, 1970).

Plesionika martia (A. Milne Edwards, 1883)

Description: Holthuis, 1951a:51, fig. 10.

Type-locality: "east Atlantic," *Travailleur* Stn. 400-1, 200 m.

Distribution: Western Atlantic: off South Carolina to Florida and off Bermuda; NE and SW Gulf of Mexico. Eastern Atlantic: off SW Ireland, Bay of Biscay, throughout Mediterranean, Gulf of Guinea, and Cape of Good Hope. Indo-West Pacific: from Gulf of Aden and east African coast to Japan and Hawaii (Pequegnat, 1970).

Plesionika tenuipes (Smith, 1881)

Description: Smith, 1881:441; 1882:59, pl.

13: fig. 12.--Pequegnat, 1970:103, figs. 4-15.

Type-locality: Off Block Island, Rhode Island, 183-461 m, *Fish Hawk* Stn. 870, 871, 873, 877, 880.

Distribution: Western Atlantic: off east coast of United States from Rhode Island to southern tip of Florida; eastern and western Gulf of Mexico, (Pequegnat, 1970).

Stylopandalus richardi (Coutière, 1905)

Description: Chace, 1985:136, fig. 62.

Type-locality: West of Madeira at 32°13'N, 23°58'W, 2000-0 m, and Canary Islands at

27°43'N, 18°28'W, 3000-0 m.

Distribution: Probably occurs in all major tropical and temperate seas (Chace, 1985). Florida (65-300 m) (personal communication, P. M. Mikkelsen].

FAMILY CRANGONIDAE

Crangon septemspinosa Say, 1818

Description: Williams, 1984:159, fig. 112.

Type-locality: "Bay shores and inlets of the sea" (east coast of the United States).

Distribution: Subarctic-boreal (Haefner, 1979) although extending beyond these limits; northern part of Gulf of St. Lawrence, doubtfully Baffin Bay (Squires, 1965) to east Florida; Arctic Alaska southward to Shumagin Islands, Alaska; Sea of Okhotsk, and Ranshima, Hokkaido, Japan (Williams, 1984).

Metacrangon jacqueti agassizii (Smith, 1882)

Description: Crosnier and Forest, 1973:233, figs. 74b, 75a, 76c.

Type-locality: Smith (1882) did not designate a type but described specimens from Blake Stns. 317, 326, 332 and 329 off the east coast of the United States.

Distribution: Western Atlantic from about 41° N south to Florida.

Parapontocaris caribbaea (Boone, 1927)

Description: Boone, 1927:125, fig. 28.--

Dardeau and Heard, 1983:10, figs. 2f, 3.

Type-locality: Caribbean Sea, off Honduras, north of Glover Reef, 870 m, *Pawnee* Stn. 1

Distribution: Western Atlantic, Bahama Islands and Straits of Florida; northwestern Gulf of Mexico, off Galveston Bay; Caribbean Sea, off Honduras (Dardeau and Heard, 1983).

Philoceras gorei (Dardeau, 1980)

Description: Williams, 1984:161, fig. 114.

Type-locality: 135 km due west of Sanibel Island Light, [Florida], 26°24'N, 83°22'W, 55 m.

Distribution: Off Georgia; off SW Florida, Cape San Blas and Padre Island, Texas (Williams, 1984).

Pontophilus brevirostris Smith, 1881

Description: Williams, 1984:161, fig. 113.

Type-locality: Material described from a series of United States Fish Commission Stns. (*Fish Hawk*) 865-867, 870-874, 877 and 878, 119 to 283 m, off Martha's Vineyard, Massachusetts, constitutes the type-series

(Smith, 1881). In 1882, Smith essentially repeated the original description, gave locality data for specimens studied from Stns. 314-315, 321, 327, 333, 344-345, and illustrated a mature female from Stn. 873, one of the stations listed in the original description. Two females from this lot, 40°02'N, 70°57'W, 183 m, are in the type collection of the USNM as are many syntypes from Stns. 865-67, 871, and 872 (Williams, 1984).

Distribution: Gulf of Maine to Gulf of Mexico off Dry Tortugas and Cuba (Williams and Wigley, 1977; Pequegnat, 1970).

FAMILY GLYPHOCRANGONIDAE

Glyphocrangon haematonotus Holthuis, 1971

Description: Holthuis, 1971:315, figs. 6, 7.

Type-locality: From *Gerda* Stn. 649, Straits of Florida, 26°34'N, 79°43'W, 494 m.

Distribution: From the east coast of South Carolina and the Bahama Islands to the Caribbean coast of Colombia and St. Vincent (Holthuis, 1971).

Glyphocrangon longleyi Schmitt, 1931

Description: Schmitt, 1931:393.--Holthuis, 1971:309, figs. 6, 7.

Type-locality: South of Dry Tortugas, Florida.

Distribution: East coast of Florida, Bahamas, Gulf of Mexico south to Santa Lucia (West Indies), Yucatan and Colombia (Holthuis, 1971).

Glyphocrangon spinicauda A. Milne Edwards, 1881

Description: A. Milne Edwards, 1881:3.--Holthuis, 1971:295, figs. 6, 7.

Type-locality: St. Kitts (17°19'27"N, 62°50'30"W, 458 m, fine gray sand and ooze).

Distribution: Western Atlantic from the east coast of Florida south to Barbados and into the Caribbean area as far west as Yucatan, Honduras, and Nicaragua (Holthuis, 1971).

INFRAORDER ASTACIDEA

FAMILY NEPHROPIDAE

Acanthacaris caeca (A. Milne Edwards, 1881)

Description: Holthuis, 1974:741, figs. 4-8.

Type-locality: Off Grenada, 12°03'15"N, 61°48'30"W, 761 m.

Distribution: Throughout the Gulf of Mexico

and the Caribbean Sea, including the Straits of Florida (Holthuis, 1974).

Metanephrops binghami (Boone, 1927)

Description: Holthuis, 1974:827, figs. 25, 26.

Type-locality: From north of Glover Reef, British Honduras (Belize).

Distribution: From the Bahama Islands to French Guiana, including the Gulf of Mexico and the Caribbean Sea (Holthuis, 1974).

Nephropsis aculeata Smith, 1881

Description: Holthuis, 1974:776, figs. 15, 16A, 16B.

Type-locality: East of New Jersey, United States of America, 40°02'N, 70°57'W, 183 m, bottom soft sticky mud.

Distribution: From east of New Jersey, to French Guiana, including the entire Gulf of Mexico and Caribbean Sea (Holthuis, 1974).

INFRAORDER THALASSINIDEA

FAMILY AXIIDAE

Axiopsis hirsutimana (Boesch and Smalley, 1972)

Description: Boesch and Smalley, 1972:45, figs. 1-9.

Type-locality: Off British Guiana, 6°50'N, 54°47'W.

Distribution: Off British Guiana; SE Pascagoula Sea Buoy, Mississippi; about 80 km northeast of tip of Mississippi River delta (Boesch and Smalley, 1972). Tortugas shrimp grounds, Florida (Williams, 1974c).

Axiopsis oxypleura (Williams, 1974)

Description: Williams, 1974c:457, figs. 11-18.

Type-locality: Straits of Florida west of Riding Rocks, 25°15'N, 79°13'W, 365 m.

Distribution: Known only from the type-locality.

Axiopsis serratifrons (A. Milne Edwards, 1873)

Description: Kensley, 1981:1253, figs. 1-5.

Type-locality: "Upolu [presumably the island in western Samoa] et les îles Sandwich" [presumably the Hawaiian Islands].

Distribution: See Kensley, 1981:1260

Coralaxius abelei Kensley and Gore, 1981

Description: Kensley and Gore, 1981:1278, figs. 1-6.

Type-locality: Atlantic Ocean, French Reef, off Key Largo, Monroe County, Florida; 25°02'N, 80°19'W; 76 m.

Distribution: French Reef, off Key Largo, Florida; Caribbean Sea, Carrie Bow Cay, Belize.

FAMILY CALLIANASSIDAE

Callianassa acanthochirus (Stimpson, 1866)

Description: Biffar, 1971a:655, figs. 3, 4.

Type-locality: Florida Keys.

Distribution: Miami; Florida Keys; Dry Tortugas, Puerto Rico; Jamaica; Barbados; Antigua; Venezuela (Biffar, 1971a).

Callianassa atlantica Rathbun, 1926

Description: Williams, 1984:180, fig. 125.

Type-locality: "Our species ranges from the coast of Southern [United] States north to Long Island Sound" (Smith, 1873).

Distribution: Bass River, Nova Scotia, to Georgia; Franklin County, Florida (Williams, 1984).

Callianassa biformis Biffar, 1971

Description: Williams, 1984:182, fig. 126.

Type-locality: South end of Sapelo Island, mouth of Doboy Sound, McIntosh County, Georgia.

Distribution: Bass River, Yarmouth and Nantucket Sound, Massachusetts (Williams and Wigley, 1977); Chesapeake Bay(?); North Inlet, South Carolina (Holland and Polgar, 1976), to McIntosh County, Georgia; Franklin County, NW Florida (Williams, 1984).

Callianassa branneri (Rathbun, 1900)

Description: Rathbun, 1900:150, pl. 8: figs. 5-8.--Biffar, 1971a: 661, figs. 5, 6.

Type-locality: Mamanguarpe Stone Reef, Brazil.

Distribution: Bermuda, southeastern Florida, including Keys and Dry Tortugas; Bimini; Little San Salvador; Puerto Rico; Barbados; Curaçao; Tobago; Brazil (Biffar, 1971a).

Callianassa fragilis Biffar, 1970

Description: Biffar, 1970:45, fig. 3; 1971a:667, figs. 7, 8.

Type-locality: Punta Arenas, Puerto Rico. Distribution: Southeastern Florida; Puerto Rico; Antigua; Venezuela (Biffar, 1971a).

Callianassa guassutunga Rodríguez, 1966

Description: Rodríguez, 1966:45, figs. 41c-

60.--Biffar, 1971a: 674, figs. 9, 10.
 Type-locality: São Sebastiao, Brazil.
 Distribution: Southeastern Florida, Puerto Rico; Brazil (Biffar, 1971a).

Callianassa jamaicense Schmitt, 1935
 Description: Schmitt, 1935b:9, pl. 1: fig. 2, pl. 2: figs. 6-8, pl. 4: figs. 1, 4.
 Type-locality: Brackish pond at Montego Bay, Jamaica.
 Distribution: Grand Isle, Louisiana to Brazil; Jamaica.

Callianassa longiventris A. Milne Edwards, 1870
 Description: Biffar, 1971a:685, figs. 13, 14.
 Type-locality: Martinique.
 Distribution: Bermuda; southeastern Florida; Jamaica; Martinique (Biffar, 1971a).

Callianassa marginata Rathbun, 1901
 Description: Rathbun, 1901:92, fig. 15.--
 Biffar, 1971a:689, figs. 15, 16.
 Type-locality: Mayaguez Harbor, Puerto Rico, 315 m.
 Distribution: Southeastern Florida, Arrowsmith Bank; Puerto Rico; Barbados (Biffar, 1971a).

Callianassa quadracuta Biffar, 1970
 Description: Biffar, 1970:40, fig. 2; 1971a:694, figs. 17, 18.
 Type-locality: Cumana, Venezuela.
 Distribution: Southeastern Florida; Venezuela (Biffar, 1971a).

Callianassa rathbunae Schmitt, 1935
 Description: Schmitt, 1935b:15, pl. 1: fig. 5, pl. 2: fig. 2, pl. 3: fig. 1, pl. 4: fig. 2.--
 Biffar, 1970:699, figs. 19, 20.
 Type-locality: Bluefields, Jamaica.
 Distribution: Miami; Jamaica (Biffar, 1970).

Callianassa trilobata Biffar, 1970
 Description: Biffar, 1970:36, fig. 1; 1971a:704, figs. 21, 22.
 Type-locality: Off Pinellas Point, Tampa Bay, Florida, 2-3 m.
 Distribution: Miami, Tampa Bay, and Lemon Bay, Florida (Biffar, 1971a).

Callichirus islagrande (Schmitt, 1935)
 Description: Schmitt, 1935b:5, pl. 1: fig. 3, pl. 3: fig. 2, pl. 4: fig. 5.
 Type-locality: Grand Isle, Louisiana.

Distribution: Grand Isle, Louisiana; Alligator Harbor, Florida.

Callichirus major (Say, 1818)
 Description: Williams, 1984:183, fig. 127.
 Type-locality: Bay Shore of St. Johns River in east Florida, near low-water mark.
 Distribution: Beaufort Inlet, North Carolina, to Cape Canaveral, Florida; Grand Terre Island to Timbalier Island, Louisiana; Espirito Santo and São Paulo, Brazil (Rodríguez, 1965, 1971; Williams, 1984).

Gouretia latispina (Dawson, 1967)
 Description: Dawson, 1967b:190, fig. 1.--
 Biffar, 1971a:679, figs. 11, 12.
 Type-locality: Grand Isle, Louisiana, 14 m.
 Distribution: Grand Isle, Louisiana; off southwestern Florida; Honduras (Biffar, 1971a).

FAMILY UPOGEBIIDAE

Upogebia affinis (Say, 1818)
 Description: Williams, 1984:191, fig. 133.
 Type-locality: Georgia.
 Distribution: Wellfleet, Massachusetts, to Rockport, Texas (Hedgpeth, 1950); through West Indies to Estado de São Paulo, Brazil (Coelho, 1966, 1970; Gomes Corrêa, 1968; Williams, 1984).

Upogebia operculata Schmitt, 1924
 Description: Schmitt, 1924b:91, pl. 5: figs. 1-4.
 Type-locality: Okra Reef, Barbados.
 Distribution: Okra Reef, Barbados; St. Thomas, Savannah Passage; Dry Tortugas, Florida (Schmitt, 1935a). Looe Key, Florida (personal communication, D. L. Felder).

INFRAORDER PALINURA

FAMILY PALINURIDAE

Justitia longimanus (H. Milne Edwards, 1837)
 Description: Bouvier, 1925:442, pl. 8: fig. 1.--
 Manning, 1978:24.
 Type-locality: The Antilles.
 Distribution: Bermuda and from southern tip of Florida through most of the Antilles (Manning, 1978).

Panulirus argus (Latreille, 1804)
 Description: Williams, 1984:170, fig. 120.

Type-locality: Erroneously given as East Indies ("des Grandes-Indies").
 Distribution: North Carolina through Gulf of Mexico and West Indies to Rio de Janeiro, Brazil; Bermuda (Williams, 1984).

Panulirus guttatus (Latreille, 1804)
 Description: Gruvel, 1911:29, fig. 12, pl. 3: fig. 3.--Holthuis, 1959:124, fig. 20.
 Type-locality: Surinam.
 Distribution: Western Atlantic Ocean from Bermuda and Florida to Brazil and the West Indies (Holthuis, 1959).

Panulirus laevicauda (Latreille, 1817)
 Description: Gruvel, 1911:45, fig. 21.--Holthuis, 1959:123.
 Type-locality: Brazil.
 Distribution: Bermuda, Florida, Cuba, Jamaica, Curaçao, French Guiana, and Brazil (Holthuis, 1959).

FAMILY SCYLLARIDAE

Parribacus antarcticus (Lund, 1793)
 Description: Holthuis, 1985:73, figs. 21, 25A.
 Type-locality: Amboina Moluccas (Holthuis, 1985).
 Distribution: The species is known both from the western Atlantic from Florida to Brazil including the West Indies and Caribbean Sea, and from the Indo-West Pacific (E and SE Africa to Formosa) (Holthuis, 1985).

Scyllarides aequinoctialis (Lund, 1793)
 Description: Lyons, 1970:15, pl. 2: figs. A, B.
 Type-locality: Jamaica.
 Distribution: West Indies and Caribbean Sea; Gulf of Mexico; southern Florida to Bermuda (Lyons, 1970).

Scyllarides nodifer (Stimpson, 1866)
 Description: Williams, 1984:174, fig. 121.
 Type-locality: Florida Keys.
 Distribution: Bermuda; Cape Lookout, North Carolina, to Florida and throughout Gulf of Mexico to Yucatan (Lyons, 1970); a postlarva from south of Long Island (29°11'N, 71°56'W) was taken in the stomach of a lancetfish (*Alepisaurus*) (Williams, 1984).

Scyllarus americanus (Smith, 1869)
 Description: Williams, 1984:176, fig. 122.
 Type-locality: Egmont Key, Florida.
 Distribution: Off Bouge Inlet, North Carolina, to Campeche Banks off Mexico, and Venezuela (Williams, 1984).

Scyllarus chacei Holthuis, 1960
 Description: Williams, 1984:177, fig. 123.
 Type-locality: North-northwest mouth of Marowijne River, about 20 mi. off coast of Surinam.
 Distribution: Off Cape Hatteras, North Carolina, through Gulf of Mexico and West Indies, and Caribbean Sea to off Cape São Roque, Brazil (Williams, 1984).

Scyllarus depressus (Smith, 1881)
 Description: Williams, 1984:178, fig. 124.
 Type-locality: South of Martha's Vineyard, 40°5'39"N, 70°23'52"W, 157.3 m.
 Distribution: Off Martha's Vineyard, Massachusetts; off Cape Hatteras, North Carolina, through Gulf of Mexico and West Indies to State of São Paulo, Brazil (Williams, 1984). Florida (78 m) (personal communication, P. M. Mikkelsen).

FAMILY SYNAXIDAE

Palinurellus gundlachi (Von Martens, 1881)
 Description: Manning, 1978:35.
 Type-locality: Cuba, Barbados.
 Distribution: Bermuda, southern Florida, most of the West Indies, Yucatan (Manning, 1978).

INFRAORDER ANOMURA

FAMILY COENOBITIDAE

Coenobita clypeatus (Herbst, 1791)
 Description: Provenzano, 1959:359, fig. 3.
 Type-locality: West Indies.
 Distribution: Florida, Bermuda, West Indies to Venezuela (Provenzano, 1959).

FAMILY DIOGENIDAE

Calcinus tibicen (Herbst, 1791)
 Description: Provenzano, 1959:363, fig. 4.
 Type-locality: Not given by Herbst.
 Distribution: Bermuda, West Indian region from NW Florida to Brazil (Provenzano, 1959, Abele, 1970).

Cancellus ornatus Benedict, 1901
 Description: Williams, 1984:193, fig. 134.
 Type-locality: Northeast Gulf of Mexico between Mississippi delta and Cedar Keys, Florida, 28°45'N, 85°02'W, 55 m.
 Distribution: Off Cape Fear, North Carolina, 33°43'N, 76°40'W to 33°42.7'N, 76°40.2'W, 90-110 m (Herbst et al., 1979) through eastern Gulf of Mexico, Greater and Lesser Antilles, to

near Los Abrolhos off central Brazil (Williams, 1984).

Cancellus viridis Mayo, 1973

Description: Mayo, 1973:28, figs. 9-11.

Type-locality: Southwest Caribbean Sea, north of Panama.

Distribution: Known from the type-locality. Southern Florida (personal communication, P. A. McLaughlin).

Clibanarius antillensis Stimpson, 1862

Description: Provenzano, 1959:368, fig. 5B.

Type-locality: Barbados.

Distribution: Southern Florida through West Indies to Curaçao and Brazil (Provenzano, 1959).

Clibanarius cubensis (Saussure, 1858)

Description: Provenzano, 1959:369, fig. 5C.

Type-locality: Cuba.

Distribution: Florida, from Miami southward, West Indies to South America (Provenzano, 1959).

Clibanarius tricolor (Gibbes, 1850)

Description: Provenzano, 1959:366, fig. 5A.

Type-locality: Key West, Florida.

Distribution: Bermuda, Florida from Miami through the Keys, West Indies (Provenzano, 1959).

Clibanarius vittatus (Bosc, 1802)

Description: Williams, 1984:194, fig. 135.

Type-locality: "Les cotes de la Caroline."

Distribution: Potomac River, Gunston, Virginia, to Florianopolis, Santa Catarina, Brazil (Forest and de Saint Laurent, 1967).

Dardanus fucosus Biffar and Provenzano, 1972

Description: Williams, 1984:196, fig. 136.

Type-locality: Off French Guiana-Brazil border, 05°29'N, 51°37'W, 64 m, Oregon Stn. 4202.

Distribution: Near Cape Hatteras, North Carolina, 35°02'N, 75°26'W, to off Amapa, extreme northern Brazil, 04°02'N, 50°33'W (Williams, 1984).

Dardanus insignis (Saussure, 1858)

Description: Williams, 1984:197, fig. 137.

Type-locality: Guadeloupe.

Distribution: Off Oregon Inlet, North Carolina, 31 m. (Cerase-Vivas et al., 1963), to Port

Aransas, Texas; through West Indies to Guadeloupe (Williams, 1984).

Dardanus venosus (H. Milne Edwards, 1848)

Description: Provenzano, 1959:374, fig. 6.

Type-locality: Guadeloupe.

Distribution: Bermuda, southern Florida, West Indies to Brazil (Provenzano, 1959).

Isocheles wurdemanni Stimpson, 1862

Description: Provenzano, 1959:375, fig. 7.

Type-locality: Gulf of Mexico, at mouth of Rio Grande.

Distribution: Texas, Louisiana, west coast of Florida and Venezuela (Provenzano, 1959).

Paguristes anomalus Bouvier, 1918

Description: Provenzano, 1959:391, figs. 12A-C.

Type-locality: Near San Diego de Cuba under old coral.

Distribution: Known only from type-locality and Long Reef, Florida (Provenzano, 1959).

Paguristes cadenati Forest, 1954

Description: Forest, 1954:353, figs. 1, 3.

Type-locality: Fort-de-France (Martinique).

Distribution: Martinique; Florida Keys.

Paguristes erythrops Holthuis, 1959

Description: Holthuis, 1959:138, figs. 24, 25.

Type-locality: Between the mouths of the Coppename and Suriname Rivers, 06°42'N, 55°38'W; bottom mud and fine shells; depth 44 m.

Distribution: From the type-locality and southern Florida (personal communication, P. A. McLaughlin).

Paguristes grayi Benedict, 1901

Description: Provenzano, 1959:387, fig. 10B.

Type-locality: San Antonio Bridge, San Juan, Puerto Rico.

Distribution: Florida Keys, Tortugas, Puerto Rico. Santo Domingo and probably generally throughout the West Indian region (Provenzano, 1959).

Paguristes hernancortezii McLaughlin and Provenzano, 1974

Description: McLaughlin and Provenzano, 1974a:207, figs. 16a, 17d-f, 18c-d, g-h, 19f-j, 20b, f-h, 21.

Type-locality: 7 mi. off Sanibel Island, Florida, M/V *Hernan Cortez* Stn. L., 26°24'N, 83°22'W, 55m.

Distribution: Known only from the type-locality.

***Paguristes hummi* Wass, 1955**

Description: Williams, 1984:200, fig. 139.

Type-locality: Alligator Harbor, Franklin County, Florida.

Distribution: Newport River, North Carolina, to off Sapelo Island, Georgia; Marco Beach, southwestern Florida, to off Isles Dernieres, Louisiana (28°38'N, 90°55'W) (Williams, 1984).

***Paguristes inconstans* McLaughlin and Provenzano, 1974**

Description: McLaughlin and Provenzano, 1974b:919, figs. 13-15.

Type-locality: Off east coast of Florida, 27°55'N, 79°05'W.

Distribution: Western and eastern coasts of Florida, southward through Caribbean to Colombia (McLaughlin and Provenzano, 1974b).

***Paguristes invisissacculus* McLaughlin and Provenzano, 1974**

Description: McLaughlin and Provenzano, 1974a:223, figs. 23b; 24d-f; 25c, d, g, h, 26e-i, 27.

Type-locality: Ragged Key, Florida.

Distribution: Florida Keys to Jamaica (McLaughlin and Provenzano, 1974a).

***Paguristes laticlavus* McLaughlin and Provenzano, 1974**

Description: McLaughlin and Provenzano, 1974b:928, figs. 16-18.

Type-locality: Off Colombia, 11°16.9'N, 74°17'W.

Distribution: Florida through Caribbean to Colombia and Venezuela (McLaughlin and Provenzano, 1974b).

***Paguristes limonensis* McLaughlin and Provenzano, 1974**

Description: McLaughlin and Provenzano, 1974b:902, figs. 7-9.

Type-locality: West side Limon Bay, Pulpit Point, Panama.

Distribution: West coast of Florida, Panama, Colombia (McLaughlin and Provenzano, 1974b).

***Paguristes lymani* A. Milne Edwards and Bouvier, 1893**

Description: Williams, 1984:201, fig. 140.

Type-locality: Sand-Key (Florida), 27 m.

Distribution: Southeast of Cape Lookout, North Carolina (150-180 m); Florida Keys to Swan Island off Honduras; through West Indies to Guyana (Williams, 1984).

***Paguristes moorei* Benedict, 1901**

Description: Williams, 1984:202, fig. 141.

Type-locality: Puerto Rico.

Distribution: Edge of continental shelf off Cape Lookout, North Carolina; Florida Straits (Hazlett, 1966); Puerto Rico (Williams, 1984).

***Paguristes oxyphthalmus* Holthuis, 1959**

Description: Holthuis, 1959:135, figs. 22b, 23.

Type-locality: About 20 mi. NNW of the mouth of the Coppename River, depth 31 m.

Distribution: Suriname. Florida (personal communication, P. A. McLaughlin).

***Paguristes puncticeps* Benedict, 1901**

Description: Provenzano, 1959:384, fig. 10A.

Type-locality: Jamaica.

Distribution: Along northwestern Florida; South Florida from Miami southward, probably generally in the West Indies (Provenzano, 1959).

***Paguristes sericeus* A. Milne Edwards, 1880**

Description: Williams, 1984:203.

Type-locality: 23°34'N, 83°16'W, (near Dry Tortugas, Florida), 66m.

Distribution: Off Cape Lookout, North Carolina; West Flower Garden Bank, NW Gulf of Mexico to Virgin Islands (Williams, 1984).

***Paguristes spinipes* A. Milne Edwards, 1880**

Description: Williams, 1984:204, fig. 143.

Type-locality: Grenada, 168 m.

Distribution: Gulf Stream south of Cape Lookout, North Carolina; off Cape Canaveral to Florida Straits, Sarasota, Florida; Barbados to Pernambuco, Brazil (Williams, 1984).

***Paguristes starcki* Provenzano, 1965**

Description: Provenzano, 1965:726, figs. 1, 2.

Type-locality: One-third mi. south-southwest of Alligator Light, Monroe County, Florida, at a depth of 6 m.

Distribution: From the type-locality.

***Paguristes tenuirostris* Benedict, 1901**

Description: Benedict, 1901:143, pl. 4: fig. 1.

Type-locality: *Grampus* Stn. 5077, 125m,

Gulf of Mexico, off west coast of Florida.
Distribution: Known only from the type-locality.

Paguristes tortugae Schmitt, 1933

Description: Williams, 1984:205, fig. 144.
Type-locality: Off Fort Jefferson Dock, Garden Key, Dry Tortugas, Florida,
Distribution: Reefs off Beaufort, North Carolina, to southern and southeastern Florida; through West Indies to northern Brazil; (?) northern Gulf of Mexico (Williams, 1984).

Paguristes triangulatus A. Milne Edwards and Bouvier, 1893

Description: Williams, 1984:206, fig. 145.
Type-locality: Barbados, 136 m.
Distribution: Off Oregon Inlet, North Carolina, (12m) to Tortugas, Florida; Barbados; Trinidad (Williams, 1984).

Paguristes wassi Provenzano, 1961

Description: Provenzano, 1961:155, fig. 1.
Type-locality: One fourth mi. south-southeast of Alligator Light, Florida Keys, at 8 m, coral and sand bottom.
Distribution: From Virgin Islands and Florida Keys.

Petrochirus diogenes (Linnaeus, 1758)

Description: Williams, 1984:198, fig. 138.
Type-locality: Near shores of Bahama Islands [Catesby, 1743 (1754 ed. as cited in Holthuis 1959)].
Distribution: Off Cape Lookout, North Carolina, through Gulf of Mexico and West Indies to off Ilha de São Sebastiao, Brazil, 23°42.5'S, 45°14.5'W (Forest and De Saint Laurent, 1967).

FAMILY LITHODIDAE

Paralomis cubensis Chace, 1939

Description: Chace, 1939:49.
Type-locality: East of Havana, Cuba, 23°12'30"N, 82°12'00"W, 439-549 m.
Distribution: From the type-locality. Florida (26°45'N, 84°55'W, 255-400 m) (personal communication, D. K. Camp).

FAMILY PAGURIDAE

Agaricochirus acanthinus McLaughlin, 1982

Description: McLaughlin, 1982:838, figs. 1g, 2g, 3g, 4, 5a, b, e.
Type-locality: Gerda Stn. 1301, 24°57'N,

80°14'W.

Distribution: Straits of Florida, western Caribbean (McLaughlin, 1982).

Agaricochirus alexandri (A. Milne Edwards and Bouvier, 1893)

Description: A. Milne Edwards and Bouvier, 1893:87, pl. 6: figs. 23-26.-- McLaughlin, 1982:834, figs. 1b, 2b, 3b.
Type-locality: Blake Stn. 132, off Santa Cruz (St. Croix), Virgin Islands.
Distribution: Straits of Florida, Caribbean to Barbados and northern coast of South America (McLaughlin, 1982).

Agaricochirus boletifer (A. Milne Edwards and Bouvier, 1893)

Description: A. Milne Edwards and Bouvier, 1893:84, pl. 6: figs. 19-22.-- McLaughlin, 1982:825, figs. 1a, 2a, 3a.
Type-locality: Blake Stn. 231, off St. Vincent, West Indies.
Distribution: Eastern Gulf of Mexico, central Caribbean, St. Vincent, West Indies (McLaughlin, 1982).

Agaricochirus gibbosimanus (A. Milne Edwards, 1880)

Description: A. Milne Edwards, 1880:42.-- McLaughlin, 1982:836, figs. 1e, 2e, 3e.
Type-locality: Blake Stn. 206, off Martinique.
Distribution: Western Atlantic off Dominican Republic; Yucatan Channel and northern Caribbean; Windward Islands (McLaughlin, 1982). Florida (personal communication, P. A. McLaughlin).

Anisopagurus bartletti (A. Milne Edwards, 1880)

Description: A. Milne Edwards and Bouvier, 1893:91, pl. 7: figs. 1-9.
Type-locality: Off St. Vincent, 146 fathoms, Blake Stn. 223.
Distribution: Southern Florida and the islands of St. Vincent, Grenada and Barbados in the West Indies.

Anisopagurus pygmaeus (Bouvier, 1918)

Description: Williams, 1984:223, fig. 159.
Type-locality: Bahia de Socapa (Zocappa) near Santiago de Cuba.
Distribution: Off Little River Inlet, South Carolina; southern Florida, including Tortugas, to Puerto Rico; Curaçao (Williams, 1984).

Catapagurus sharrei A. Milne Edwards, 1880
Description: Forest and De Saint Laurent, 1967:151, figs. 124-135.

Type-locality: Antilles

Distribution: Western Atlantic south to Brazil (Forest and De Saint Laurent, 1967).

Iridopagurus caribbensis (A. Milne Edwards and Bouvier, 1893)

Description: Williams, 1984:207, fig. 146.

Type-locality: Flannegan Passage, Virgin Islands, *Blake* Stn. 142.

Distribution: ESE of Charleston, South Carolina (32°34'N, 79°05'W); WSW of Panama City, Florida (30°19'N, 86°15.5'W); southern Florida, Virgin Islands, and Guadeloupe (Williams, 1984). East and west coast of Florida; Alabama; Bahama Islands; Lesser Antilles; Venezuela; Curacao; off Santa Marta, Colombia (García-Gómez, 1983).

Iridopagurus globulus De Saint Laurent-Dechancé, 1966

Description: De Saint Laurent-Dechancé, 1966:169, figs. 28, 33, 38.

Type-locality: Northwest Providence Channel, Bahama Islands, *Gerda* Stn. 522.

Distribution: The Straits of Florida; the Northwest Province Channel, Bahama Islands; off Barbados, and Golfo de Uraba, Colombia (García-Gómez, 1983).

Iridopagurus iris (A. Milne Edwards, 1880)

Description: A. Milne Edwards and Bouvier, 1893:112 (in part), pl. 8: figs. 14-25.--García-Gómez, 1983:16.

Type-locality: Off Barbados, *Blake* Stn. 290.

Distribution: Known in the western Atlantic from southern Florida, off northwest Little Bahama Bank south to North Bahama Islands, Puerto Rico, Saint Vincent in the Windward Islands, Barbados, Trinidad and Tobago, French Guiana, Venezuela, Colombia, off Southwest Cay in the southwestern Caribbean, and Islas Mujeres, off the Yucatan Peninsula, Mexico (García-Gómez, 1983).

Iridopagurus reticulatus García Gómez, 1983

Description: García-Gómez, 1983:37, figs. 3, 4.

Type-locality: Off north coast of Crooked Island, Bahama Islands; 3-5 m.

Distribution: Off the coast of North Carolina; off Bermuda; Hollywood Beach, Florida; Cay Sal; off the Grand Bahama Island, southeast through various localities off the Bahama

Islands, Jamaica, Dominican Republic, the United States Virgin Islands, the Leeward Islands and Bonaire in the Lesser Antilles, Venezuela, Curaçao, Colombia, Surinam and to its southern and eastern range, off French Guiana (García-Gómez, 1983).

Iridopagurus violaceus De Saint Laurent-Dechancé, 1966

Description: De Saint Laurent-Dechancé, 1966:165, figs. 16, 22, 26, 31, 36.

Type-locality: Off Fernando do Norhona, Brazil, *Calypso* Stn. 19.

Distribution: Off the west coast of Florida, the Florida Keys, Little Bahama Bank and off Castle Roads, South coast of Bermuda; through the Antillean arc, from Antigua to the Grenadines, off the north coast of Venezuela, Colombia and Panama, and from off the coast of Tobago southeast to French Guiana; Brazil (García-Gómez, 1983).

Manucomplanus corallinus (Benedict, 1892)

Description: Williams, 1984:224, fig. 160.

Type-locality: Off Key West, Florida.

Distribution: Off Cape Lookout, North Carolina, to Gulf of Mexico between Cedar Keys, Florida, and Mississippi delta; off Cape Catoche, Yucatan (Williams, 1984).

Nematopaguroides pusillus Forest and Saint Laurent, 1967

Description: Forest and St. Laurent 1967:159, figs. 142-146.

Type-locality: *Calypso* Stn. 23, Brazil, 08°19.5'S, 34°39'W, 75 m.

Distribution: Southern Florida and the type-locality.

Ostraconotus spatulipes A. Milne Edwards, 1880

Description: A. Milne Edwards and Bouvier, 1893:169, pl. 12.

Type-locality: *Sigsbee* Stn. 50; 119 fathoms 26°31'N, 85°53'E; Florida.

Distribution: Florida and Barbados.

Pagurus annulipes (Stimpson, 1860)

Description: Williams, 1984:210, fig. 148.

Type-locality: Beaufort Harbor, North Carolina.

Distribution: Vineyard Sound, Massachusetts, to at least northern Florida (McLaughlin, 1975).

***Pagurus brevidactylus* (Stimpson, 1859)**

Description: Lemaitre et al., 1982:675.

Type-locality: Bathsheba, Barbados.

Distribution: Western Atlantic from Bermuda, northeastern Florida and Bahamas to Brazil; Caribbean; Gulf of Mexico (Lemaitre et al., 1982).

***Pagurus carolinensis* McLaughlin, 1975**

Description: Williams, 1984:212, fig. 150.

Type-locality: Black Rocks, off New River, North Carolina.

Distribution: Newport River (Kellogg, 1971) and Cape Lookout, North Carolina, to Southern Florida (Williams, 1984).

***Pagurus criniticornis* (Dana, 1852)**

Description: Dana, 1852:448.--Lemaitre et al., 1982:684, figs. 1a, b.

Type-locality: Rio de Janeiro, Brazil.

Distribution: Gulf of Mexico; Caribbean; south Atlantic from Brazil to Argentina (Lemaitre et al., 1982).

***Pagurus defensus* (Benedict, 1892)**

Description: Williams, 1984:213, fig. 151.

Type-locality: Gulf of Mexico between delta of Mississippi River and Cedar Keys, Florida, 55 m.

Distribution: Cape Hatteras, North Carolina, to Georgia; Torgugas, Florida, to Alabama (Williams, 1984).

***Pagurus gymnodactylus* Lemaitre, 1982**

Description: Lemaitre, 1982:657.--Lemaitre et al., 1982:687.

Type-locality: 21.75 mi. northeast Cedar Keys Light, Florida.

Distribution: Gulf of Mexico from Mexico to west Florida (Lemaitre et al., 1982).

***Pagurus impressus* (Benedict, 1892)**

Description: Williams, 1984:215, fig. 153.

Type-locality: Boca Ciega Bay, inner shore of Pine Key (mouth of Tampa Bay), Florida (from holotype jar label).

Distribution: Off Diamond Shoals, North Carolina, to near Cape Canaveral, Florida; Florida Bay, near Flamingo, north to vicinity of Pensacola, Florida (Cooley, 1978); Port Aransas, Texas (Williams, 1984).

***Pagurus longicarpus* Say, 1817**

Description: Williams, 1984:216, fig. 154.

Type-locality: "Inhabits bay shore" (east coast

of United States).

Distribution: Minas Basin and Chignecto Bay (Bousfield and Leim 1960) to Hutchinson Island, Florida (Camp et al., 1977); southwestern Florida to coast of Texas (Provenzano 1959; Rouse 1970; Whitten et al., 1950; Williams, 1984).

***Pagurus maclaughlinae* García-Gómez, 1982**

Description: García-Gómez, 1982:647, figs. 1, 2.

Type-locality: Wading Beach, Matheson Hammock, Miami, Florida.

Distribution: From northern Gulf of Mexico to Florida Keys, and from central eastern Florida to Puerto Rico (García-Gómez, 1982).

***Pagurus marshi* Benedict, 1901**

Description: Provenzano, 1959:405, fig. 17.--Lemaitre et al., 1982:680.

Type-locality: Ponce, Puerto Rico.

Distribution: South Florida, through Caribbean to Colombia (Lemaitre et al., 1982).

***Pagurus piercei* Wass, 1963**

Description: Williams, 1984:218, fig. 155.

Type-locality: 39 mi. SE Port Aransas, Texas, 73 m.

Distribution: Texas around the gulf coast of Florida to Georgia.

***Pagurus politus* (Smith, 1882)**

Description: Williams, 1984:219, fig. 156.

Type-locality: Not designated in original description, but syntypes from four localities off New Jersey to Massachusetts were indicated by Smith (1882). The male he illustrated, is in the type collection of USNM (21452), from off Martha's Vineyard, Massachusetts, 40°03'48"N, 70°45'54"W, 130 m, *Fish Hawk* Stn. 922; others in the collection of MCZ, Harvard University, are from United States Fish Commission Stn. 309, 40°11'40"N, 68°22'10"W, 556 m, and Stn. 310, 39°59'16"N, 70°18'30"W, 475.5 m (Williams, 1984).

Distribution: Georges Bank to off Dry Tortugas, Florida (Williams, 1974c).

***Pagurus pollicaris* Say, 1817**

Description: Williams, 1984:220, fig. 157.

Type-locality: (East) "coast of the United States".

Distribution: Grand Manan, New Brunswick,

to northeastern Florida; Key West, Florida, to Texas (Provenzano, 1959, in part).

Pagurus provenzano Forest and De Saint Laurent, 1967

Description: Forest and De Saint Laurent, 1967:118, figs. 72-77, 93, 94.-- Lemaître et al., 1982:672.

Type-locality: *Calypso* Stn. 27, 08°25.5'S, 30°48.5'W.

Distribution: Bermuda southeastern Florida and Bahamas through eastern Caribbean to Brazil, south as far as Uruguay; western Caribbean from Yucatan Peninsula to Colombia (Lemaître et al., 1982).

Pagurus stimpsoni (A. Milne Edwards and Bouvier, 1893)

Description: Lemaître et al., 1982:687, fig. 2.

Type-locality: West Coast of Florida.

Distribution: Western Atlantic from North Carolina to Florida; Gulf of Mexico; Caribbean coast of South America (Lemaître et al., 1982).

Phimochirus holthuisi (Provenzano, 1961)

Description: Williams, 1984:225, fig. 161.

Type-locality: Sand patch on coral rock bottom, 4.5 mi. SE Ram's Head, St. John, Virgin Islands, 15-18 m.

Distribution: Off Oregon Inlet, North Carolina (Herbst et al., 1979) to Alabama(?); through West Indies to Surinam (Williams, 1984).

Phimochirus leurocarpus McLaughlin, 1981

Description: McLaughlin, 1981b:356, figs. 4f, 9b, 10b, 11a-f, 12a-e.

Type-locality: Pillsbury Stn. 736, 10°57'N, 65°52'W.

Distribution: Bermuda, Straits of Florida, Caribbean Sea south of Puerto Rico and north coast of Venezuela (McLaughlin, 1981b).

Phimochirus operculatus (Stimpson, 1859)

Description: McLaughlin, 1981b:336, figs. 4a, 5a, 7a.

Type-locality: Tortugas, Florida.

Distribution: South Florida, Curaçao, Colombia (McLaughlin, 1981b).

Phimochirus randalli (Provenzano, 1961)

Description: Provenzano, 1961:159, fig. 2.-- McLaughlin, 1981b: 340, figs. 4b, 5b, 7b.

Type-locality: Ridge 5 mi. southeast of Lameshur Bay, St. John, Virgin Islands.

Distribution: Bahama Islands; Straits of

Florida, eastern and western Caribbean (McLaughlin, 1981b).

Pylopaguropsis atlantica Wass, 1963

Description: Wass, 1963:153, fig. 10.

Type-locality: Off Surinam, 07°25'N, 54°35'W; 137-146 m; *Oregon* Stn. 2289.

Distribution: Known from the type-locality and Florida (27°47.3'N, 79°57.6'W, 95-99m) (personal communication, D. K. Camp).

Pylopagurus discoidalis (A. Milne Edwards, 1880)

Description: Williams, 1984:226, fig. 162.

Type-locality: Montserrat, 220 m.

Distribution: ENE Oregon Inlet, North Carolina (Provenzano, 1963), through eastern Gulf of Mexico and West Indies to mouth of Amazon River, Brazil (Williams, 1984).

Rhodochirus rosaceus (A. Milne Edwards and Bouvier, 1893)

Description: Williams, 1984:227, fig. 163.

Type-locality: Grenada, 168 m.

Distribution: South of Cape Lookout, North Carolina; southern Florida; Grenada and Surinam (Williams, 1984).

Solenopagurus lineatus (Wass, 1963)

Description: Wass, 1963:139, fig. 3.

Type-locality: 07°25'N 54°35'W; 135-145 m; *Oregon* Stn. 2289.

Distribution: North Carolina to Surinam (Wass, 1963).

Tomopaguropsis problematica (A. Milne Edwards and Bouvier, 1893)

Description: Williams, 1984:228, figs. 164-165.

Type-locality: Near Sand Key (SSW Key West, Florida), 228.6 m.

Distribution: NE Cape Lookout, North Carolina; southern Florida and Bahamas; Barbados; off Honduras (16°39'N, 82°29'W) (Williams, 1984).

Tomopagurus chacei (Wass, 1963)

Description: McLaughlin, 1981a:21, figs. 1h, 7h, 8g.--Wass, 1963: figs. 11a-g.

Type-locality: Off Surinam, *Oregon* Stn. 2289, 07°25'N, 54°35'W.

Distribution: Western Atlantic and Caribbean (McLaughlin, 1981a).

Tomopagurus cokeri (Hay, 1917)

Description: Hay, 1917:73.--McLaughlin,

1981a:13, figs. 1d, 2a, 3a, 4a, 7a, 8a.
 Type-locality: Thirty miles S of Cape Lookout (lightship), North Carolina.
 Distribution: Western Atlantic, Gulf of Mexico and Caribbean (McLaughlin, 1981a).

Tomopagurus cubensis (Wass, 1963)
 Description: Wass, 1963:134, figs. 1a-d.--
 McLaughlin, 1981a:21, figs. 1e, 7e, 8e.
 Type-locality: North of Matanzas Province, Cuba, *Atlantis* Stn. 3480, 23°10'N, 81°28'W.
 Distribution: Western Atlantic and Caribbean; 183-366 m (McLaughlin, 1981a).

Tomopagurus rubropunctatus A. Milne Edwards and Bouvier, 1893
 Description: A. Milne Edwards and Bouvier, 1893:71, pl. 6: figs. 1-6.-- McLaughlin, 1981a:10, figs. 1a, 7d, 8d.
 Type-locality: Barbados, *Blake* Stn. 290.
 Distribution: Western Atlantic and Caribbean Sea (McLaughlin, 1981a).

Tomopagurus wassi McLaughlin, 1981
 Description: McLaughlin, 1981a:14, figs. 1b, 2b, 3b, 4b, 5, 6, 7b, 7c, 8b.
 Type-locality: Off Venezuela, *Oregon* Stn. 1985, 09°41'N, 59°47'W.
 Distribution: Southeastern United States, Straits of Florida, Gulf of Mexico, Caribbean, to northern Brazil (McLaughlin, 1981a).

FAMILY CHIROSTYLIDAE

Uroptychus armatus A. Milne Edwards, 1880
 Description: A. Milne Edwards and Bouvier, 1897: 132, pl. 11: figs. 3, pl. 12: figs. 8, 9.
 Type-locality: *Blake* Stn. 241, 163 fathoms, off Cariacou, Venezuela.
 Distribution: Cariacou, Venezuela; Florida (personal communication, P. A. McLaughlin).

FAMILY GALATHEIDAE

Galathea rostrata A. Milne Edwards, 1880
 Description: Williams, 1984:232, fig. 167.
 Type-locality: 16 mi. north of Jolbos Island (Yucatan Peninsula), at 26 m.
 Distribution: Off Cape Hatteras, North Carolina to southern Florida, northwestern Florida to Mississippi delta, and off Cape Catoche, Yucatan, Mexico (Gore, 1979).

Munida angulata Benedict, 1902
 Description: Benedict, 1902:252, fig. 4.

Distribution: Florida (29°40'N, 80°16'W, 64 m; 29°19.4'N, 80°17.7'W, 45.7 m) (personal communication with D. K. Camp).

Munida affinis A. Milne Edwards, 1880
 Description: A. Milne Edwards, 1880:48.--
 Chace, 1942b:55, fig. 22.
 Type-locality: Off St. Kitts, West Indies.
 Distribution: Known from the type-locality, north and south coast of Cuba (Chace, 1942b). 100 miles south of Panama City, Florida, 183 m (personal communication, D. K. Camp).

Munida forceps A. Milne Edwards, 1880
 Description: A. Milne Edwards, 1880:49.--
 Chace, 1942b:39, fig. 15.
 Type-locality: Off Alacran Reef, north of Yucatan, in 154 m.
 Distribution: North coast of Cuba and throughout the Gulf of Mexico (Pequegnat and Pequegnat, 1970). Florida (122-215 m) (personal communication, P. M. Mikkelsen).

Munida iris iris A. Milne Edwards, 1880
 Description: Williams, 1984:233, fig. 168.
 Type-locality: Off Barbados, 382 m, *Blake* Stn. 274.
 Distribution: SSE of Martha's Vineyard, Massachusetts, through southeastern Gulf of Mexico to near Cozumel Island, Yucatan, and through Caribbean islands to off mouth of Amazon River (Williams, 1984).

Munida irrasa A. Milne Edwards, 1880
 Description: Williams, 1984:234, fig. 169.
 Type-locality: Not designated with certainty; syntypes from 10 localities in the Gulf of Mexico and Caribbean, (MCZ) (Williams, 1984).
 Distribution: Off Cape Lookout, North Carolina, through eastern Gulf of Mexico and Caribbean Sea to 34°14'S, 51°40'W off Uruguay; "600 mi. off St. Davids, Bermuda" (USNM) (Williams, 1984).

Munida longipes A. Milne Edwards, 1880
 Description: Williams, 1984:235, fig. 170.
 Type-locality: Not designated with certainty; syntypes from 7 localities off Cuba and the Lesser Antilles (MCZ).
 Distribution: SE Cape Lookout, North Carolina, through Gulf of Mexico to British Honduras, and through West Indies to Curaçao (Williams, 1984).

Munida miles A. Milne Edwards, 1880

Description: A. Milne Edwards, 1880:51.--Chace, 1942b:36.

Type-locality: Gulf of Mexico; *Blake* Stn. 45; of Havana, Cuba, *Blake* Stn. 53.

Distribution: North coast of Cuba; eastern Gulf of Mexico; the Caribbean off Honduras (Pequegnat and Pequegnat, 1970) and throughout the Lesser Antilles; south as far as Pernambuco, Brazil.

Munida pusilla Benedict, 1902

Description: Williams, 1984:236, fig. 171.

Type-locality: *Albatross* Stn. 2405, Gulf of Mexico (south of Cape San Blas, Florida, 28°45'N, 85°02'W, 55 m).

Distribution: Off Cape Lookout, North Carolina, to Straits of Florida and through eastern Gulf of Mexico to Yucatan; Colombia and Trinidad (Williams, 1984).

Munida santipauli Henderson, 1885

Description: Henderson, 1885:411.--Chace, 1942b:38.

Type-locality: Saint Peter and Saint Paul Rocks (00°56'N, 29°22'W), Atlantic Ocean.

Distribution: In Western Atlantic from off Florida to St. Paul's Rocks and in the eastern Atlantic from the Azores to the African coast in the region of the Canary Islands; off South Africa (Chace, 1942b).

Munida simplex Benedict, 1902

Description: Benedict, 1902:272, fig. 19.

Type-locality: *Albatross* Stn. 2169; depth 140 m.

Distribution: Off Havana, Cuba; Florida.

Munida spinifrons Henderson, 1885

Description: Henderson, 1888:144, pl. 15, fig. 1.

Type-locality: *Challenger* Stn. 113A, off Fernando Noronha; 7-25 fathoms.

Distribution: Southern Florida and the type-locality.

Munida stimpsoni A. Milne Edwards, 1880

Description: A. Milne Edwards and Bouvier, 1897:48, pl. 4: figs. 1-13.

Type-locality: The original material was collected from 20 *Blake* stations throughout the West Indies. A type-locality was not designated although the description was apparently based on material from *Blake* Stn. 143, 150 fathoms, 17°30'N, 69°43.5'E (A. Milne Edwards and Bouvier, 1897).

Distribution: From north coast of Cuba through the West Indian regions to Grenada (Chace, 1942b). 100 miles south of Panama City, Florida, 183 m (personal communication, D. K. Camp).

Munida valida Smith, 1883

Description: Williams, 1984:237, fig. 172, 173.

Type-locality: Off Southern New England *Fish Hawk* Stn. 1112, 39°56'N, 70°35'W, 448 m; Stn. 1124, 40°01'N, 68°54'W, 1171 m.

Distribution: Off southern New England through Gulf of Mexico to Golfo de Morrosquillo, Colombia, and Curaçao (Williams, 1984).

Munidopsis armata (A. Milne Edwards, 1880)

Description: Chace, 1942b:74.--Mayo, 1974:72, figs. 6, 7.

Type-locality: Fredericksted (St. Croix, Virgin Islands), *Blake* Stn. 137, 1144m.

Distribution: Known from the Caribbean and from the Straits of Florida south to British Guiana in the western Atlantic (Mayo, 1974).

Munidopsis platirostris (A. Milne Edwards and Bouvier, 1894)

Description: Chace, 1942b:75.--Mayo, 1974:216, fig. 31.

Type-locality: Off Barbados, 183 m.

Distribution: Known in the western Atlantic from the Straits of Florida, Arrowsmith Bank in the northwest Caribbean, north and south of the Dominican Republic, and in the Lesser Antilles (southeastern Caribbean) from Dominica to Barbados (Mayo, 1974).

Munidopsis polita (Smith, 1883)

Description: Smith, 1883:50, pl. 2: fig. 1, pl. 3: figs. 1-5. --Pequegnat and Pequegnat, 1970:155.

Type-locality: Western North Atlantic (off Martha's Vineyard), *Fish Hawk* Stn.

Distribution: Off the east coast of the United States (off Martha's Vineyard) and in NW Gulf of Mexico. Florida (182-212 m) (personal communication, P. M. Mikkelsen).

Family Porcellanidae***Euceramus praelongus*** Stimpson, 1860

Description: Williams, 1984:239, fig. 174.

Type-locality: Beaufort, North Carolina.

Distribution: Delaware Bay (USNM; Watling and Maurer, 1976) to Aransas area of Texas coast (Williams, 1984).

Megalobrachium poeyi (Guérin-Méneville, 1855)

Description: Benedict, 1901:136, pl. 3: fig. 8.

Type-locality: Cuba.

Distribution: Widely distributed from eastern Central Florida throughout the eastern Caribbean to Brazil, rarely in Panama (Gore and Abele, 1976).

Megalobrachium soriatum (Say, 1818)

Description: Williams, 1984:240, fig. 175.

Type-locality: St. Catherines Island, Georgia.

Distribution: Off Cape Hatteras, North Carolina, to Port Aransas, Texas; West Indies to Barbados; Contoy, Mexico; Bahia Caledonia and Galeta Island, Panama (Williams, 1984).

Neopisosoma angustifrons (Benedict, 1901)

Description: Benedict, 1901:135, pl. 3: fig. 6.

Type-locality: Trinidad.

Distribution: Known from the southwestern Gulf of Mexico, the Lesser Antilles, Trinidad, Cubagua, Islas La Tortuga, Bonaire, Curacao, Panama and Venezuela; littoral (Gore and Abele, 1976). Florida (personal communication, P. A. McLaughlin).

Pachycheles ackleianus A. Milne Edwards, 1880

Description: A. Milne Edwards and Bouvier, 1923:295, pl. 2: fig. 12, pl. 4: fig. 2.

Type-locality: Blake Stn. 11, off west coast of Florida, and Stn. 39, Jolbos Islands.

Distribution: Tampa Bay, Florida; Gulf of Mexico; North of Dry Tortugas; Jolbos Islands, North of Yucatan; Jamaica; St. Thomas; Barbados (Haig, 1956).

Pachycheles monilifer (Dana, 1852)

Description: Dana, 1852:413; 1855, pl. 26: fig. 3.

Type-locality: Rio de Janeiro, Brazil.

Distribution: Outer Hillsboro Reef, Florida; Contoy, Mexico; Isla Cubagua, Venezuela; Mamanguape, Brazil; Pernambuco; Rio de Janeiro; Bahia, Brazil (Haig, 1956).

Pachycheles pilosus (H. Milne Edwards, 1837)

Description: Williams, 1984:241, fig. 176.

Type-locality: Vicinity of Charleston, South Carolina.

Distribution: Charleston, South Carolina; Key West to Sarasota Bay, Florida; through West Indies to Tobago and Aruba (Williams, 1984).

Pachycheles riisei (Stimpson, 1858)

Description: Schmitt, 1935a:188, fig. 48.

Type-locality: St. Thomas.

Distribution: Florida, Key West; Puerto Rico; St. Thomas; Barbados; Ilha Trindade; Maceio, Alagoas, Brazil (Haig, 1956).

Pachycheles rugimanus A. Milne Edwards, 1880

Description: Williams, 1984:242, fig. 177.

Type-locality: Contoy and W of Florida.

Distribution: Off Cape Lookout, North Carolina, through Florida to St. Thomas, Virgin Islands, and Contoy Island, Mexico; Pernambuco, Brazil (Coelho, 1964; Williams, 1984).

Parapetrolisthes tortugensis (Glassell, 1945)

Description: Glassell, 1945:228, fig. 2.

Type-locality: In and around Tortugas, Florida.

Distribution: Dry Tortugas, Florida; off Isla La Tortuga, Venezuela; Virgin Islands.

Petrolisthes armatus (Gibbes, 1850)

Description: Haig, 1960:50, pl. 19: fig. 2.

Type-locality: Florida.

Distribution: Widely distributed from the tropical western coast of Africa, the east central coast of Florida, the Gulf of Mexico and throughout the Caribbean as far south as Santa Catharina, Brazil; in the eastern Pacific from the Gulf of California, Mexico, to Peru (Gore and Abele, 1976).

Petrolisthes galathinus (Bosc, 1802)

Description: Williams, 1984:243, fig. 178.

Type-locality: Unknown.

Distribution: Cape Hatteras, North Carolina, through Gulf of Mexico and Caribbean Sea to Rio de Janeiro, Brazil; Ilha Trindade off Brazil; Pacific Ocean from Isla San Lucas, Costa Rica, to off La Libertad, Ecuador (Williams, 1984).

Petrolisthes jugosus Streets, 1872

Description: Schmitt, 1935a:185.--Gore and Abele, 1976:11.

Type-locality: Saint Martin Islands.

Distribution: Known from the southwestern Gulf of Mexico, the Antilles in the eastern Caribbean, Trinidad, Tobago, and some islands along the northern coast of South America, westward to Panama; Boca Raton, Palm Beach County, Florida (Gore and Abele, 1976).

***Petrolisthes politus* (Gray, 1831)**

Description: Schmitt, 1935a:187.-- Gore and Abele, 1976:12.

Type-locality: Not designated.

Distribution: Florida Keys; Vera Cruz, Mexico; Puerto Rico; West Indies; Panama; Curaçao.

***Polyonyx gibbesi* Haig, 1956**

Description: Williams, 1984:244, fig. 179.

Type-locality: Coast of South Carolina.

Distribution: Woods Hole, Massachusetts, to Uruguay (Coelho and Ramos, 1972).

***Porcellana sayana* (Leach, 1820)**

Description: Williams, 1984:245, fig. 180.

Type-locality: Coast of Georgia and Florida.

Distribution: Cape Hatteras, North Carolina, around Gulf of Mexico and Caribbean Sea to Rio Grande de Sul, Brazil (Coelho and Ramos, 1972).

***Porcellana sigsbeiana* A. Milne Edwards, 1880**

Description: Williams, 1984:246, fig. 181.

Type-locality: Blake Stn. 49, off delta of Mississippi River, 216 m; 36, north of Yucatan, 154 m; 142, Flannegan Passage (Virgin Islands), 49 m.

Distribution: Off Martha's Vineyard, Massachusetts, to southwestern Caribbean Sea off Colombia (Gore, 1970); West Indies to Virgin Islands (Williams, 1984).

***Porcellana stimpsoni* A. Milne Edwards, 1880**

Description: A. Milne Edwards and Bouvier, 1923:292, pl. 1: figs. 4-5.

Type-locality: Woman Key, Florida.

Distribution: Florida, Woman Key.

FAMILY ALBUNEIDAE***Albunea gibbesii* Stimpson, 1859**

Description: Williams, 1984:248, fig. 182.

Type-locality: St. Augustine, Florida.

Distribution: East of Cape Lookout, North Carolina, to Texas; through West Indies to São Sebastiao, São Paulo, Brazil (Williams, 1984).

***Albunea paretii* Guérin-Méneville, 1853**

Description: Williams, 1984:249, fig. 182, 183.

Type-locality: (Uncertain), America.

Distribution: Beaufort Inlet, North Carolina, to Corpus Christi, Texas; through West Indies to Santa Catarina, Brazil (Coelho and Ramos,

1972); West Africa from Cape Verde Islands and Senegal to Ghana (Williams, 1984).

***Lepidopa benedicti* Schmitt, 1935**

Description: Efford, 1971:76, figs. 1a, 23, 3a, 43, p, q, 5a, 63, n, 7a.

Type-locality: The outer beach of Santa Rosa Island, Pensacola, Florida.

Distribution: Florida; Gulf of Mexico; SE of Veracruz, near Mocambo, Mexico.

***Lepidopa websteri* Benedict, 1903**

Description: Williams, 1984:250, fig. 184.

Type-locality: Beach near Fort Macon (Carteret County), North Carolina.

Distribution: Around mouth of Chesapeake Bay (larvae); Drum Inlet, North Carolina, to Sapelo Island, Georgia; Tampa Bay, Florida; Ship Island and Petit Bois Island, Mississippi (Efford, 1971; Sandifer and Van Engle, 1972; Sandifer 1973; Williams, 1984).

***Zygopa michaelis* Holthuis, 1960**

Description: Holthuis, 1960:22, fig. 1, 2.

Type-locality: Sint Michiels Baai, south coast of Curaçao, Netherlands Antilles; sandy-bottom; depth about 4 m.

Distribution: From Curaçao, in southern Caribbean, to southern Florida and eastern Gulf of Mexico.

FAMILY HIPPIDAE***Emerita benedicti* Schmitt, 1935**

Description: Williams, 1984:251, fig. 185.

Type-locality: Tampa Bay, Florida.

Distribution: Charleston County, South Carolina, to Veracruz, Mexico (Efford, 1976).

***Emerita portoricensis* Schmitt, 1935**

Description: Schmitt, 1935a:215, figs. 72a, b.

Type-locality: Mayaguez, Puerto Rico.

Distribution: South and West Florida; Texas; Honduras; Colombia; Puerto Rico; Jamaica; Trinidad (Schmitt, 1935a).

***Emerita talpoida* (Say, 1817)**

Description: Williams, 1984:252, fig. 186.

Type-locality: (East) coast of United States.

Distribution: Harwich [Barnstable County], Massachusetts to Horn Island, Mississippi; Progreso, Yucatan, Mexico (Schmitt, 1935a; Efford, 1976).

***Hippa cubensis* (Saussure, 1857)**

Description: Schmitt, 1935a:217.

Type-locality: Cuba.
 Distribution: Florida to Brazil; West Indies; Ascension Island; Bahamas; West Africa; Puerto Rico, Hucares, St. Thomas (Schmitt, 1935a).

INFRAORDER BRACHYURA

FAMILY DROMIIDAE

Dromia erythropus (George Edwards, 1771)

Description: Rathbun, 1937:31, fig. 11, pl. 6: figs. 1, 2, pl. 8: figs. 1, 2.

Type-locality: Not indicated.

Distribution: Bermuda; Bahamas; Florida Keys and Dry Tortugas; off Louisiana and Texas; north coast of Cuba; Jamaica; Haiti; Puerto Rico; St. Thomas to Barbados; Netherlands Antilles; Pernambuco to São Paulo, Brazil (Powers, 1977).

Dromidia antillensis Stimpson, 1858

Description: Williams, 1984:255, fig. 187.

Type-locality: St. Thomas, Virgin Island, Key Biscayne and Dry Tortugas, Florida.

Distribution: Off Cape Hatteras, North Carolina, through Gulf of Mexico and Caribbean Sea to Rio de Janeiro, Brazil; Bermuda; Saint Helena (Forest, 1974; Williams, 1984).

Hypoconcha arcuata Stimpson, 1858

Description: Williams, 1984:257, fig. 188.

Type-locality: South Carolina sandy shores and St. Thomas (Virgin Island).

Distribution: Off Cape Lookout, North Carolina, to west Florida; St. Thomas, Virgin Island; Surinam (Holthuis, 1959) to Espirito Santo, Brazil (Williams, 1984).

Hypoconcha sabulosa (Herbst, 1799)

Description: Williams, 1984:258, fig. 189.

Type-locality: Listed as "Africa" (probably an error).

Distribution: Off Cape Hatteras, North Carolina, through Gulf of Mexico to Bahia, Brazil (Williams, 1984).

Hypoconcha spinosissima Rathbun, 1933

Description: Williams, 1984:258, fig. 190.

Type-locality: Off Cape Hatteras, North Carolina, 89.6 m.

Distribution: Off Cape Hatteras, North Carolina, to Gulf of Mexico off Mississippi delta and Yucatan; Jamaica (Williams, 1984).

FAMILY HOMOLODROMIIDAE

Dicranodromia ovata A. Milne Edwards, 1880

Description: Rathbun, 1937:60, fig. 15, pl. 13: figs. 3-4.

Type-locality: Barbados, 329 m.

Distribution: East and west coasts of Florida; Florida Keys and Straits; off north coast of Cuba; northwest Caribbean Sea; Guadeloupe; Barbados (Powers, 1977).

FAMILY CYMONOMIDAE

Cymonimus quadratus A. Milne Edwards, 1880

Description: Rathbun, 1937:98, fig. 23, pl. 30: fig. 3, pl. 31: fig. 3.

Type-locality: Havana to Grenada, 320-930 m.

Distribution: Northwest of Dry Tortugas; Cuba; Puerto Rico; Lesser Antilles, from St. Croix to Grenada (Powers, 1977).

Cymopolus agassizi A. Milne Edwards and Bouvier, 1899

Description: Rathbun, 1937:100, pl. 30: fig. 2, pl. 31: fig. 2.

Type-locality: Sand Key, 137 m.

Distribution: Florida Straits; Florida Keys; Puerto Rico (Powers, 1977).

FAMILY CYCLODORIPPIDAE

Clythrocerus granulatus (Rathbun, 1898)

Description: Williams, 1984:259, fig. 191.

Type-locality: Off Trinidad.

Distribution: ESE Cape Lookout, and SE Cape Fear, North Carolina; Honduras; southern Florida through Antilles to Venezuela and Trinidad (Williams, 1984).

Clythrocerus nitidus (A. Milne Edwards, 1880)

Description: Rathbun, 1937:109, figs. 26, 27, pl. 33: figs. 1, 2.

Type-locality: Florida Keys and Grenada.

Distribution: South Carolina to west Florida; Florida Keys; Grenada; 12-479 m.

Clythrocerus stimpsoni Rathbun, 1937

Description: Rathbun, 1937:121, fig. 32, pl. 34: figs. 1, 2.

Type-locality: West coast of Florida; 183 m.

Distribution: Known only from the type specimen.

Tymolus antennaria (A. Milne Edwards, 1880)

Description: Rathbun, 1937:104, fig. 24, pl.

32: figs. 1, 2.

Type-locality: Twenty stations, ranging from Havana to Barbados, 158 to 517 m.

Distribution: West coast of Cuba; north coast of Yucatan (Gulf); north coast of Cuba; Puerto Rico; Lesser Antilles, from Dominica to Grenada (Powers, 1977). Florida (26°48'N, 84°37'W, 190.5-210 m) (personal communication, D. K. Camp).

FAMILY HOMOLIDAE

Homola barbata (Fabricius, 1793)

Description: Williams, 1984:261, fig. 193.

Type-locality: Bay of Naples.

Distribution: Off southeastern Massachusetts to Rio de Janeiro, Brazil; eastern Atlantic Ocean from Portugal and Azores to Cape Verde Islands and Angola; South Africa; Mediterranean Sea (Williams, 1984).

FAMILY LATREILLIDAE

Latreillia manningi Williams, 1982

Description: Williams, 1984:262, fig. 194.

Type-locality: American Shoal Light, Florida, about 10 mi. N by W, 192-201 m.

Distribution: Nantucket Shoals off Massachusetts to off Havana, Cuba; Venezuela; Ascension Island; Frost (1936) reported a megalopa off Newfoundland (Williams, 1984). Florida (64 m) (personal communication, P. M. Mikkelsen).

FAMILY RANINIDAE

Lyreidus nitidus (A. Milne Edwards, 1880)

Description: A. Milne Edwards, 1880:34.--Goeke, 1980:149.

Type-locality: Grenada, British West Indies, 259, 288m.

Distribution: Martha's Vineyard; Gulf of Mexico; Greater Antilles; off Tortugas, Florida; off Surinam, NE of Paramaribo; north coast of Venezuela, off Puerto Cabello.

Ranilia constricta (A. Milne Edwards, 1880)

Description: Williams, 1984:265, fig. 196.

Type-locality: Near Sombrero (Florida?), 86 m. (see Rathbun, 1937).

Distribution: SE of Cape Fear, North Carolina, 33°42'N, 76°39.5'W, 140 m; Palm Beach, Florida, to Florida Straits and Yucatan Channel; Cuba; off Barbados; Ascension Island; eastern Atlantic from Sierra Leone and Annobon Island (Manning and Holthuis, 1981).

Ranilia muricata H. Milne Edwards, 1837

Description: Williams, 1984:266, fig. 197.

Type-locality: Unknown.

Distribution: North Carolina; Bahamas; Florida Straits; southern to northwestern Florida; Swan Island (Caribbean), Colombia.

Raninoides loevis (Latreille, 1825)

Description: Williams, 1984:267, fig. 198.

Type-locality: Unknown.

Distribution: S Cape Hatteras, 35°03.2'N, 75°35.1'W; around Gulf of Mexico and southern Caribbean Sea, including Leeward Islands, to Bahia, Brazil (Williams, 1984).

Raninoides louisianensis Rathbun, 1933

Description: Rathbun, 1937:12, figs. 6, 7, pl. 1: figs. 5, 6.

Type-locality: East of Mississippi delta, 122 m.

Distribution: Gulf of Mexico, from the Mississippi delta to Campeche Banks (Powers, 1977). South of Appalachicola Bay, Florida (personal communication, D. K. Camp).

Symethis variolosa (Fabricius, 1793)

Description: Williams, 1984:264, fig. 195.

Type-locality: "In oceano Indico."

Distribution: SE Cape Lookout, North Carolina, 34°10'N, 76°10'W, through western Gulf of Mexico (Goeke, 1980) to Bahia, Brazil; Fernando de Noronha.

FAMILY DORIPPIDAE

Ethusa mascarone americana A. Milne Edwards, 1880

Description: Williams, 1984:269, fig. 199.

Type-localities: West Florida, 23.7 m, and West Florida, 26°16'N, 36.6 m (A. Milne Edwards, 1880).

Distribution: S of Cape Lookout, North Carolina (34°06'N, 76°15'W) to Gulf of Mexico and West Indies; Maranhao to Bahia, Brazil; Golfo de California; Taboga Island, Panama (Williams, 1984).

Ethusa microphthalmia Smith, 1881

Description: Williams, 1984:269, fig. 200.

Type-locality: Off Martha's Vineyard, Massachusetts, 260.6 m, Stn. 878, *Fish Hawk*, 39°55'00"N, 70°54'15"W.

Distribution: Off Martha's Vineyard, Massachusetts, to Cuba and around Gulf of Mexico (Pequegnat et al., 1971).

***Ethusa tenuipes* Rathbun, 1897**

Description: Williams, 1984:270, fig. 201.

Type-locality: Off Key West, 91.5m.

Distribution: ESE Cape Lookout, North Carolina (94-77 m); East Florida to Gulf of Mexico E of Mississippi River delta; Cuba (Chace, 1940b; Williams, 1984).

***Ethusa truncata* A. Milne Edwards and Bouvier, 1899**

Description: Rathbun, 1937:85, pl. 28: figs. 1-2.

Type-locality: Gulf of Mexico (Not Antilles) 217-218 m.

Distribution: Off west coast of Florida; off Mississippi delta and Louisiana; northwest of Trinidad (Powers, 1977).

FAMILY CALAPPIDAE***Acanthocarpus alexandri* Stimpson, 1871**

Description: Williams, 1984:271, fig. 202.

Type-locality: Off Quicksands, Florida Keys, 135 m.

Distribution: George Bank off Massachusetts to west coast of Florida; Puerto Rico to Grenadines; Rio de Janeiro, Brazil (Coelho and Ramos, 1972).

***Acanthocarpus bispinosus* A. Milne Edwards, 1880**

Description: Rathbun, 1937:224, pl. 68: figs. 1-3.

Type-locality: Reefs of the Grenadines, 256 m.

Distribution: Off West and northwest coasts of Florida; Dry Tortugas; Grenadines, Windward Islands (Powers, 1977).

***Calappa angusta* A. Milne Edwards, 1880**

Description: Williams, 1984:273, fig. 203.

Type-locality: Barbados.

Distribution: Off Cape Lookout, North Carolina, through eastern and southwestern Gulf of Mexico, to Venezuela (Turkay, 1968) and Grenada (Williams, 1984).

***Calappa flammea* (Herbst, 1794)**

Description: Williams, 1984:273, figs. 204-205.

Type-locality: America.

Distribution: Woods Hole region, Massachusetts, to Florida Keys; Gulf coast of United States and Mexico; Bahamas; Bermuda (Williams, 1984).

***Calappa gallus* (Herbst, 1803)**

Description: Rathbun, 1937:214, pl. 65:

figs. 2, 3.

Type-locality: "East Indies."

Distribution: Bermuda; Bahamas; Florida Keys and Dry Tortugas; northwest Cuba; Jamaica; Puerto Rico; St. Croix to Barbados; Off Campeche snapper banks (Gulf of Mexico); Panama (Caribbean) to Venezuela; Netherlands Antilles; Ceara to Bahia, Brazil; St. Helena Island (South Atlantic); off Western Africa, from Senegal to Angola; South Africa; Red Sea; Reunion and Seychelles, in Persian Gulf; off India and Maldives; Philippines; Formosa; Japan; Marshall Inlands; Samoa; Hawaiian Islands (Powers, 1977).

***Calappa ocellata* Holthuis, 1958**

Description: Williams, 1984:275, figs. 206-207.

Type-locality: Klein Bonaire, Dutch West Indies.

Distribution: Cape Hatteras, North Carolina, to Rio de Janeiro, Brazil; Bermuda (Williams, 1984).

***Calappa sulcata* Rathbun, 1898**

Description: Williams, 1984:276, figs. 208-209.

Type-locality: Off Louisiana 29°24'30"N, 88°01'00"W, 64 m.

Distribution: Cape Hatteras, North Carolina, through Gulf of Mexico to Sergipe, Brazil (Williams, 1984).

***Cycloes bairdii* Stimpson, 1860**

Description: Williams, 1984:278, fig. 210.

Type-locality: Cape St. Lucas, Mexico.

Distribution: Bermuda; ESE Cape Lookout, North Carolina, to Espírito Santo, Brazil, in west Atlantic; tip of Baja California to Ecuador and Galapagos Islands, including Clarion, Socorro and Cocos Islands, in eastern Pacific (Williams, 1984).

***Hepatus epheliticus* (Linnaeus, 1763)**

Description: Williams, 1984:279, fig. 211.

Type-locality: Carolina.

Distribution: Chesapeake Bay to western Bay of Campeche, Mexico (Rickner, 1977); Cuba; Jamaica; Dominican Republic (Williams, 1984).

***Hepatus pudibundus* (Herbst, 1785)**

Description: Williams, 1984:280, figs. 212, 213.

Type-locality: Martinique.

Distribution: Georgia to State of São Paulo, Brazil, (Coelho and Ramos, 1972).

***Osachila antillensis* Rathbun, 1898**

Description: Rathbun, 1937:251, pl. 77: fig. 2.
Type-locality: Off Havana, 209m.

Distribution: North coast of Cuba; St. Croix, Virgin Islands; Montserrat; Dominica; Barbados; Grenada (Powers, 1977). Florida (personal communication, P. A. McLaughlin).

***Osachila semilevis* Rathbun, 1916**

Description: Williams, 1984:281, fig. 214.
Type-locality: Gulf of Mexico, 48 m.

Distribution: Off Beaufort, North Carolina, to northwest Florida (Williams, 1984).

***Osachila tuberosa* Stimpson, 1871**

Description: Williams, 1984:282, fig. 215.

Type-locality: Five stations among the south Florida reefs.

Distribution: Off Cape Hatteras, North Carolina, to northwest Florida and Yucatan Channel (Pequegnat 1970; Springer and Bullis, 1956).

FAMILY LEUCOSIIDAE***Callidactylus asper* Stimpson, 1871**

Description: Williams, 1984:289, fig. 224.

Type-locality: Three stations off Florida keys, 29-69 m.

Distribution: S of Cape Lookout, North Carolina (Williams et al., 1968), through SE Gulf of Mexico to Panama and southeastward to Alagoas, Brazil (Coelho and Ramos, 1972).

***Ebalia cariosa* (Stimpson, 1860)**

Description: Williams, 1984:284, fig. 216.

Type-locality: Beaufort, North Carolina.

Distribution: Beaufort, North Carolina, to west Florida; western Gulf of Mexico (Rickner, 1977); Jamaica; northeastern South America to São Paulo, Brazil (Williams, 1984).

***Ebalia stimpsonii* A. Milne Edwards, 1880**

Description: Williams, 1984:284, fig. 217.

Type-locality: Barbados, 13.72 to 91.45 m.

Distribution: SE Cape Lookout, North Carolina; west Florida to Barbados; off mouth of Amazon River, Brazil (Williams, 1984).

***Iliacantha intermedia* Miers, 1886**

Description: Williams, 1984:290, fig. 225.

Type-locality: Bahia, Brazil.

Distribution: Off Beaufort, North Carolina, to NW Florida; St. Thomas, Virgin Islands; Venezuela; Ceara and Bahia, Brazil (Williams, 1984).

***Iliacantha liodactylus* Rathbun, 1898**

Description: Rathbun, 1937:186, text-fig. 41, pl. 55: figs. 1-2.

Type-locality: North of Trinidad, West Indies.

Distribution: West coast of Florida; ?Haiti; Puerto Rico; St. John, Virgin Islands; Trinidad; Alagoas to Bahia, Brazil (Powers, 1977).

***Iliacantha sparsa* Stimpson, 1871**

Description: Rathbun, 1937:190, pl. 56: figs. 1-2.

Type-locality: West of Tortugas, 55 m.

Distribution: Northwest of the Dry Tortugas; off north and southeast coasts of Puerto Rico; Barbados; Maranhao to Bahia, Brazil (Powers, 1977).

***Iliacantha subglobosa* Stimpson, 1871**

Description: Williams, 1984:290, fig. 226.

Type-locality: Three stations in Florida reefs, 73-146 m.

Distribution: Off Cape Hatteras, North Carolina, to northwest Florida; through eastern Gulf of Mexico and Caribbean Sea south to Alagoas, Brazil (Williams, 1984).

***Lithadia cadaverosa* Stimpson, 1871**

Description: Rathbun, 1937:137, pl. 38: figs. 3-6.

Type-locality: West of Tortugas, 64 m, and off Conch Reef, Florida, 72 m.

Distribution: Bahamas; northwest coast of Florida and northeast portion of Gulf of Mexico (Powers, 1977).

***Lithadia granulosa* A. Milne Edwards, 1880**

Description: Rathbun, 1937:140, fig. 36.

Type-locality: Off St. Croix Island, 210 m.

Distribution: Known from type specimen and southern Florida (personal communication, P. A. McLaughlin).

***Myropsis quinquespinosa* Stimpson, 1871**

Description: Williams, 1984:287, fig. 222.

Type-locality: Tennessee Reef, Florida Keys.

Distribution: South of Martha's Vineyard, through Gulf of Mexico and Caribbean Sea to Surinam (Williams, 1984).

***Persephona crinita* Rathbun, 1931**

Description: Rathbun, 1937:163, pl. 43: figs. 2, 3, pl. 44: figs. 1-3.

Type-locality: Horn Island Pass, Mississippi, 5 m.

Distribution: Northwest Florida to Texas;

Trinidad; Ilha São Sebastiao, Brazil (Powers, 1977).

Persephona mediterranea (Herbst, 1794)

Description: Williams, 1984:288, fig. 223.

Type-locality: Erroneously, Mediterranean Sea.

Distribution: New Jersey through Gulf of Mexico and Caribbean Sea to Santa Catarina, Brazil (Williams, 1984).

Speloeophorus elevatus Rathbun, 1898

Description: Rathbun, 1937:145, pl. 39: figs. 7-9.

Type-locality: Off Key West.

Distribution: Florida Keys; Jamaica; off Cape St. Roque and from Maranhao to Alagoas, Brazil (Powers, 1977).

Speloeophorus nodosus (Bell, 1855)

Description: Williams, 1984:285, figs. 218-219.

Type-locality: Unknown.

Distribution: Florida; West Indies (Williams, 1984).

Speloeophorus pontifer (Stimpson, 1871)

Description: Williams, 1984:286, figs. 220-221.

Type-locality: Barbados.

Distribution: Southeast of Cape Lookout and off Beaufort, North Carolina, to west Florida; West Indies to Barbados (Williams, 1984).

Uhlias limbatus Stimpson, 1871

Description: Rathbun, 1937:150, pl. 36: figs. 3-5.

Type-locality: St. Thomas.

Distribution: West of Key West, Florida; north coast of Cuba; Jamaica; Haiti; St. Thomas, Virgin Islands (Powers, 1977).

FAMILY MAJIDAE

Acanthonyx petiverii H. Milne Edwards, 1834

Description: Rathbun, 1925:142, fig. 52, pl.

44, pl. 222: figs. 1-6.

Type-locality: Antilles.

Distribution: Bahamas; southeast and northwest Florida; Texas; Cuba; Jamaica; Puerto Rico; Virgin Islands; Netherlands Antilles; Panama (Caribbean) to Rio de Janeiro, Brazil. Along the Pacific coast, from Baja California to Caldera, Chile; Galapagos Islands (Powers, 1977).

Achaeopsis thomsoni (Norman, 1873)

Description: Rathbun, 1925:29, text-fig. 7, pl. 10.

Type-locality: Deep water between the Faroes and Scotland.

Distribution: Western Atlantic from Nantucket Shoals to Grenada. Eastern Atlantic from Faroe Islands to Cape Verde; Mediterranean, Gough Island (South Atlantic); Agulhas Bank, near cape of Good Hope. Indian Ocean. West and South Pacific Ocean. South Australia (Rathbun, 1925).

Aepinus septemspinosus (A. Milne Edwards, 1879)

Description: Williams, 1984:292, fig. 227, 241c.

Type-locality: Florida Strait, 24°55'N, 83°25'W, 67.7 m.

Distribution: S Cape Lookout, North Carolina, 34°10'N, 76°10'W; SW Cape San Blas, Florida, and Bahama Banks to Bahia, Brazil (Williams, 1984).

Anasimus latus Rathbun, 1894

Description: Williams, 1984:293, fig. 228, 241n.

Type-locality: Gulf of Mexico, east of delta of Mississippi River, 29°14'30" N, 88°09'30" W, 124.4 m.

Distribution: Off Cape Lookout, North Carolina, through Gulf of Mexico (Felder, 1973) to Amapa, Brazil (Coelho and Ramos, 1972).

Anomalothir furcillatus (Stimpson, 1871)

Description: Williams, 1984:294, fig. 229, 241a.

Type-locality: Off "The Samboes" (southern Florida), 225 m.

Distribution: Off Cape Lookout, North Carolina, through eastern Gulf of Mexico and West Indies to Grenada (Williams, 1984).

Arachnopsis filipes Stimpson, 1871

Description: Williams, 1984:295, fig. 230, 241d.

Type-locality: Off Conch, Carysfort and French reefs, Florida.

Distribution: SE Capes Hatteras and Lookout, North Carolina; Gulf of Mexico off NW Florida; through West Indies to off Rio Grande do Norte, Brazil.

Batrachonotus fragosus Stimpson, 1871

Description: Williams, 1984:296, 297, fig. 231, 241e.

Type-locality: South of Tortugas, (Florida), 24°36'40"N, 80°02'20"W, 29.3 m.

Distribution: Cape Hatteras, North Carolina, to southern and western Florida; West Indies to Barbados (Williams, 1984).

Chorinus heros (Herbst, 1790)

Description: Rathbun, 1925:305, fig. 101, pl. 107, pl. 246: figs. 3-5; 1933:20, fig. 21.

Type-locality: "Der Ocean."

Distribution: Bermuda; Florida Keys and Dry Tortugas; Cuba; Caribbean coast of Yucatan, Mexico; Jamaica; Hispaniola; Puerto Rico; St. Croix; Barbados; Ceara to Bahia, Brazil (Powers, 1977).

Coelocerus spinosus A. Milne Edwards, 1875

Description: Williams, 1984:315, figs. 251, 259i.

Type-locality: Off Florida, 34.75 m.

Distribution: Off Cape Fear, North Carolina, to near Cape Canaveral, Florida; W Florida to E of Mississippi River delta (Williams, 1984).

Collodes leptocheles Rathbun, 1894

Description: Rathbun, 1925:117, fig. 42, pl. 38: figs. 5-6.

Type-locality: Gulf of Mexico, 124-309 m.

Distribution: All quadrants of Gulf of Mexico except southeast, off coasts of Florida, Alabama, and Texas; off Vera Cruz, Mexico (Powers, 1977).

Collodes nudus Stimpson, 1871

Description: Rathbun, 1925:110.

Type-locality: Off Carysfort Reef, 73 m.

Distribution: Known only from the male holotype off Carysfort Reef, Florida, 25°13'40"N, 80°10'45"W; 73 m; sand.

Collodes obesus A. Milne Edwards, 1878

Description: Rathbun, 1925:109, pl. 36: figs. 3, 4, pl. 217: figs. 2-5.

Type-locality: Near Sombrero, Florida Strait, 99 m.

Distribution: Known only from the type-locality.

Collodes robustus Smith, 1883

Description: Rathbun, 1925:114, text-fig. 36-41, pl. 29.--Williams, 1984:297, fig. 241g.

Type-locality: Twenty-one stations between off Martha's Vineyard to off Chesapeake Bay, 1-2 to 285 m.

Distribution: North of Cape Cod, 42°12'N, 70°13'W, to southeast of Cape Lookout, North Carolina (Williams, 1984). Florida (27°37'N, 83°58'W, 73 m) [personal communication, D. K. Camp].

Collodes trispinosus Stimpson, 1871

Description: Williams, 1984:297, fig. 232, 241f.

Type-locality: Off the Quicksands, Carysfort Reef, and French Reef (Florida), 62.1 to 91 m.

Distribution: Near Cape Hatteras, North Carolina, to south and west Florida near Apalachicola (Williams, 1984).

Epialtus bituberculatus H. Milne Edwards, 1834

Description: Rathbun, 1925:148, figs. 53a, 54, pl. 45: figs. 3, 4.

Type-locality: Chile.

Distribution: East coast of Florida; Key West, Florida; Puerto Rico; Panama (Caribbean) to Colombia; Ceara to Pernambuco, Brazil (Powers, 1977).

Epialtus dilatatus A. Milne Edwards, 1878

Description: Williams, 1984:313, figs. 249, 259d.

Type-locality: St. Thomas.

Distribution: Off Beaufort Inlet and New River, North Carolina; southwest Florida; Yucatan; Bahamas to St. Thomas (Williams, 1984).

Epialtus dilatatus forma elongata Rathbun, 1923

Description: Rathbun, 1925:154, fig. 53k, pl. 48.

Type-locality: Off Duck Key, Florida.

Distribution: Florida Keys; south, west, and northwest coasts of Florida.

Epialtus kingsleyi Rathbun, 1923

Description: Rathbun, 1925:152, pl. 45: fig. 1.

Type-locality: Florida.

Distribution: Florida.

Epialtus longirostris Stimpson, 1860

Description: Rathbun, 1925:151, figs. 53g, 56.

Type-locality: Ensenada Honda, Culebra Island.

Distribution: Key West and west coast of Florida; Cuba; Jamaica; St. Thomas, Virgin Islands; northeast Brazil (Powers, 1977).

Euprognatha gracilipes A. Milne Edwards, 1878

Description: Rathbun, 1925:101, pl. 34: figs. 3, 4.

Type-locality: 23°32'N, 88°05'W, 174 m.

Distribution: Florida Keys; north coast of

Yucatan; off north coast of Cuba; Puerto Rico; St. Croix, Virgin Islands; Barbados; Amapa to São Paulo, Brazil (Powers, 1977).

Euprognatha rastellifera Stimpson, 1871
Description: Williams, 1984:298, figs. 233, 241b.

Type-locality: Southwest of Martha's Vineyard, Massachusetts, 40°00'N, 70°57'W, 155 m.

Distribution: Off Georges Bank (40°35'N, 67°37'W) to São Paulo, Brazil (Coelho and Ramos, 1972).

Hemus cristulipes A. Milne Edwards, 1875
Description: Williams, 1984:325, fig. 261.
Type-locality: Near Contoy (Yucatan), at the entrance to the Gulf of Mexico, 21.9 to 32.9 m.
Distribution: Off Cape Lookout, North Carolina and South Carolina; northwest of Gulf of Mexico and Yucatan, through West Indies to Pernambuco, Brazil (Powers, 1977; Herbst et al., 1979).

Inachoides forceps A. Milne Edwards, 1879
Description: Williams, 1984:299, figs. 234, 240i.

Type-locality: Guiana and Desterro, Brazil.
Distribution: SE Cape Lookout, North Carolina; west coast of Florida to Desterro (= Florianopolis), Brazil (Williams, 1984).

Leptopisa setirostris (Stimpson, 1871)
Description: Rathbun, 1925:375, pl. 134: figs. 1-3; pl. 253: fig. 2.
Type-locality: Florida Keys.
Distribution: From Miami to northern Brazil; Puerto Rico, Vieques, Culebra, St. Thomas (Rathbun, 1933).

Libinia dubia H. Milne Edwards, 1834
Description: Williams, 1984:316, figs. 252, 259g.
Type-locality: "Côtes des États-Unis."
Distribution: Cape Cod, Massachusetts, to southern Texas; Bahamas and Cuba (Williams, 1984).

Libinia emarginata Leach, 1815
Description: Williams, 1984:318, figs. 253, 259h.
Type-locality: Unknown.
Distribution: Windsor, Nova Scotia, to western Gulf of Mexico (Williams, 1984).

Libinia erinacea (A. Milne Edwards, 1879)
Description: Rathbun, 1925:321, pl. 10.
Type-locality: 24°44'N, 83°26'W, between Florida and Cuba, 69 m.
Distribution: Florida Keys; southeast to northwest Florida; north coast of Cuba (Powers, 1977).

Macrocoeloma camptocerum (Stimpson, 1871)
Description: Williams, 1984:326, figs. 262, 275m.
Type-locality: Near Key West (Florida), 3.7 to 9.2 m.
Distribution: Beaufort Harbor, North Carolina, around southern Florida to Alligator Harbor, Florida (Williams, 1984).

Macrocoeloma diplacanthum (Stimpson, 1860)
Description: Rathbun, 1925:478, pl. 169: fig. 1, pl. 269: fig. 1-3.
Type-locality: St. Thomas.
Distribution: Key West, Florida; Cuba; Jamaica; Puerto Rico; Virgin Islands; Guadeloupe; Curaçao, Netherlands Antilles; Old Providence Island (Caribbean) (Powers, 1977).

Macrocoeloma eutheca (Stimpson, 1871)
Description: Williams, 1984:327, figs. 263, 275k.
Type-locality: Off French Reef (Florida), 27.4 m., and west of Tortugas, 67.7 m.
Distribution: SE of Cape Lookout, North Carolina; off NW Florida through Bahama Banks and West Indies; Panama (Williams, 1984).

Macrocoeloma laevigatum (Stimpson, 1860)
Description: Rathbun, 1925:483, fig. 136, pl. 169: figs. 2, 3.
Type-locality: St. Thomas.
Distribution: Florida Keys; north coast of Cuba; Jamaica; St. Thomas, Virgin Islands; Guadeloupe; Piaui to Alagoas, Brazil (Powers, 1977).

Macrocoeloma septemspinosum (Stimpson, 1871)
Description: Rathbun, 1925:477, pl. 173, figs. 2-3.
Type-locality: West of Tortugas, 65 m.
Distribution: South Carolina; Bahamas; Florida Keys; northeast quadrant of Gulf; Ceara to Rio Grande do Norte, Brazil (Powers, 1977).

Macrocoeloma subparallelum (Stimpson, 1860)

Description: Rathbun, 1925:480, pl. 172.

Type-locality: St. Thomas.

Distribution: North coast of Cuba; Jamaica; Haiti; Puerto Rico; St. Thomas, Virgin Islands; Guadeloupe; Barbados; Old Providence Island (Caribbean); Rio Grande do Norte, Pernambuco; Brazil (Powers, 1977).

Macrocoeloma trispinosum trispinosum (Latreille, 1825)

Description: Rathbun, 1925:466, fig. 132, pl. 166; fig. 1, pl. 167.

Type-locality: "Nouvelle Hollande" (an error).

Distribution: North Carolina; Bermuda; south Florida to northwest Florida; off Louisiana and Texas; Gulf and Caribbean coasts of Yucatan, Mexico; Cuba; Jamaica; Puerto Rico; St. Thomas to St. Lucia; Curaçao, Netherlands Antilles; Piaui to Bahia, Brazil (Powers, 1977).

Macrocoeloma trispinosum nodipes (Desbonne, 1867)

Description: Williams, 1984:328, figs. 264, 275l.

Type-locality: "Nouvelle Holland" (?) (error).

Distribution: Beaufort, North Carolina, to Alligator Harbor, Florida; Yucatan; through West Indies to Bahia, Brazil (Williams, 1984).

Macrocoeloma trispinosum, Variety

Description: Rathbun, 1925:468, pl. 168: fig. 1.

Type-locality: None designated.

Distribution: From North Carolina to Gulf of Mexico, including Florida; Yucatan; West Indies and Caribbean Sea (Rathbun, 1925).

Metoporphaphis calcarata (Say, 1818)

Description: Williams, 1984:300, figs. 235, 240h.

Type-locality: Bay of Charleston, South Carolina.

Distribution: Off Cape Hatteras, North Carolina, through Gulf of Mexico and Caribbean Sea to Rio de Janeiro, Brazil (Williams, 1984).

Microphrys antillensis Rathbun, 1920

Description: Williams, 1984:329, figs. 265, 275h.

Type-locality: Off Montego Bay Point, Jamaica.

Distribution: Near Capes Hatteras and

Lookout, North Carolina, to Cape Fear, North Carolina; Cuba; Jamaica; Puerto Rico; Pernambuco, Brazil (Williams, 1984).

Microphrys bicornutus (Latreille, 1825)

Description: Williams, 1984:330, figs. 266, 275g.

Type-locality: "Nouvelle Hollande".

Distribution: Near Beaufort, North Carolina, through Gulf of Mexico (Ray, 1974) to Florianopolis, Santa Catarina, Brazil; Bermuda (Williams, 1984).

Mithrax acuticornis Stimpson, 1870

Description: Williams, 1984:332, figs. 267, 275a.

Type-locality: Off the Quicksands (Florida), 62.6 m; west of the Tortugas, 67.7 m and 76.8 m.

Distribution: Off Cape Lookout, North Carolina; west Florida and Yucatan Channel through West Indies to Espirito Santo, Brazil (Williams, 1984).

Mithrax caribbaeus Rathbun, 1900

Description: Rathbun, 1925:409, plates 148, 149.

Type-locality: St. Thomas.

Distribution: West Indies to South America, Puerto Rico, St. Thomas, St. Croix. Airport Lagoon, Key West, Dry Tortugas, 29-33 m (personal communication, D. K. Camp).

Mithrax cinctimanus (Stimpson, 1860)

Description: Rathbun, 1925:438, pl. 158.

Type-locality: Tortugas and St. Thomas.

Distribution: Bahamas and Florida Keys to West Indies and Curaçao (Rathbun, 1925).

Mithrax cornutus Saussure, 1857

Description: Rathbun, 1925:386, pl. 137: figs. 3-4, pl. 256.

Type-locality: Antilles.

Distribution: Bermuda; Florida Straits; north coast of Cuba; between Jamaica and Haiti; Dominica; Martinique; off Bahia, Brazil (Powers, 1977).

Mithrax coryphe (Herbst, 1801)

Description: Rathbun, 1925:426, pl. 153.

Type-locality: Not Known.

Distribution: Miami to southern Brazil; Puerto Rico, Culebra, St. Thomas, St. Croix (Rathbun, 1933).

Mithrax forceps (A. Milne Edwards, 1875)

Description: Williams, 1984:337, figs. 272, 275f.

Type-locality: Guiana.

Distribution: From Cape Hatteras, North Carolina, through Gulf of Mexico to Rio de Janeiro, Brazil; Bermuda (Williams, 1984).

Mithrax hemphilli Rathbun, 1892

Description: Rathbun, 1925:395, pl. 139, pl. 259: fig. 2.

Type-locality: Indian Key, Florida.

Distribution: Florida Keys to Rio de Janeiro, Brazil; Culebra (Rathbun, 1933).

Mithrax hispidus (Herbst, 1790)

Description: Williams, 1984:333, figs. 268, 275d.

Type-locality: Unknown.

Distribution: Delaware Bay (Say, 1818), off Charleston Harbor, South Carolina, and Georgia (Gibbes, 1850). Northwestern Gulf of Mexico; Bahamas and Florida Keys through West Indies to São Paulo, Brazil; Bermuda (Williams, 1984).

Mithrax holderi Stimpson, 1871

Description: Rathbun, 1925:392, pl. 138: figs. 1, 2, pl. 257: fig. 2.

Type-locality: Tortugas, 13 m.

Distribution: Florida Keys and Dry Tortugas; north and south coasts of Cuba; Jamaica; Puerto Rico; Virgin Islands (Powers, 1977).

Mithrax pilosus Rathbun, 1892

Description: Rathbun, 1925:394, pl. 138: fig. 3, pl. 258.

Type-locality: Abaco, Bahamas.

Distribution: Bahamas; Florida Keys and Dry Tortugas; Vera Cruz, Mexico; Cuba; Puerto Rico; St. Thomas; Virgin Islands to Barbados; Caribbean coast of Panama; Venezuela (Powers, 1977).

Mithrax pleuracanthus Stimpson, 1871

Description: Williams, 1984:334, figs. 269, 275e.

Type-locality: Key West, 3.6-9.1 m, Tortugas (Florida), 9.1-11 m; St. Thomas.

Distribution: Beaufort, North Carolina, to Pensacola, Florida, western Gulf of Mexico to Yucatan Channel off Cape Catoche, Mexico; West Indies to Venezuela; Bermuda (Williams, 1984).

Mithrax ruber (Stimpson, 1871)

Description: Rathbun, 1925:432, pl. 15 '.

Type-locality: Cruz del Padre, Cuba.

Distribution: Cuba to Curaçao and Barbados; Puerto Rico, St. Thomas, Water Island (Rathbun, 1933)

Mithrax sculptus (Lamarck, 1818)

Description: Rathbun, 1925:422, figs. 125, 126, pl. 152.

Type-locality: Unknown.

Distribution: From Miami to Brazil; Puerto Rico; Vieques; Culebra; St. Thomas; Water Island.

Mithrax spinosissimus (Lamarck, 1818)

Description: Williams, 1984:335, figs. 270, 275b.

Type-locality: "Ile-de-France," Locality erroneous.

Distribution: North (?) and South Carolina to Nicaragua, and through West Indies to Barbados and Venezuela (Williams, 1984).

Mithrax tortugae Rathbun, 1920

Description: Rathbun, 1925:417, pl. 147, fig. 2.

Type-locality: Tortugas.

Distribution: Bahamas; Florida Keys; Curaçao.

Mithrax verrucosus H. Milne Edwards, 1832

Description: Williams, 1984:336, figs. 271, 275c.

Type-locality: Robert Bay, Martinique.

Distribution: Charleston, South Carolina; Campeche Banks; through West Indies to Fernando Noronha Island, Brazil (Williams, 1984).

Mocosoa crebripunctata Stimpson, 1871

Description: Rathbun, 1925:159, fig. 59, pl. 49: figs. 3-4.

Type-locality: Off French Reef, Florida, 27 m.

Distribution: Florida Straits; off Cape San Blas, northwest Florida; Maranhao to Espirito Santo, Brazil.

Nibilia antilocapra (Stimpson, 1871)

Description: Williams, 1984:320, figs. 254, 259f.

Type-locality: Florida, off Carysfort Reef, 95 and 109.7 m; and off Alligator Reef, 251.8 m.

Distribution: Off Cape Hatteras, North Carolina, to Gulf of Mexico just east of Mississippi River delta and Gulf of Campeche;

Windward Islands, West Indies, off Guyana (Williams, 1984).

Oplopisa spinipes A. Milne Edwards, 1879
Description: Rathbun, 1925:228, pl. 232: figs. 1, 2.

Type-locality: Straits of Florida, 185 m.
Distribution: Known only from the type-locality.

Pelia mutica (Gibbes, 1850)

Description: Williams, 1984:321, figs. 255, 259a.

Type-locality: Charleston Harbor, off White Point Battery, South Carolina.

Distribution: Buzzards Bay and Vineyard Sound, Massachusetts, to off Port Mansfield, Willacy County, Texas (Felder, 1973); Cuba, Puerto Rico, and St. Thomas, West Indies (Williams, 1984).

Picroceroides tubularis Miers, 1886

Description: Rathbun, 1925:354, fig. 115, pl. 126, pl. 254: figs. 2-5.

Type-locality: Fernando Noronha and Bahia, in shallow water.

Distribution: Bahamas; southeast Florida; north and south coasts of Cuba; between Jamaica and Haiti: St. Thomas, Virgin Islands; Maranhao to Espirito Santo, Brazil (Powers, 1977).

Pitho aculeata (Gibbes, 1850)

Description: Rathbun, 1925:357, fig. 116c, pl. 127, pl. 251: fig. 1.

Type-locality: Key West, and "Florida".

Distribution: Bahamas; Florida Keys and Dry Tortugas; west coast of Florida; north coast of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Guadeloupe; Old Province Island (Caribbean); Netherlands Antilles (Powers, 1977).

Pitho anisodon (Von Martens, 1872)

Description: Rathbun, 1925:368, figs. 116b, 117d, 118, pl. 131, pl. 251: fig. 2.

Type-locality: Cuba.

Distribution: Bahamas; south, west and northwest coast of Florida; Florida Keys; north coast of Cuba; Jamaica; Puerto Rico; Guadeloupe; Curaçao, Netherlands Antilles (Powers, 1977).

Pitho lherminieri (Schramm, 1867)

Description: Williams, 1984:311, figs. 246, 259a.

Type-locality: Guadeloupe.

Distribution: Off Beaufort Inlet, North Carolina, to west Florida; Veracruz, Mexico; West Indies to Islet of São Paulo, Brazil (Williams, 1984).

Pitho laevigata (A. Milne Edwards, 1875)

Description: Rathbun, 1925:372, pl. 132: figs. 3-4, pl. 133: fig. 3, pl. 250: figs. 11-13.

Type-locality: Antilles.

Distribution: West and northwest coasts of Florida; Antilles, location unspecified; Colombia; Trinidad (Powers, 1977).

Pitho mirabilis (Herbst, 1794)

Description: Rathbun, 1925:366, figs. 116d, 117c, pl. 128: fig. 3; pl. 129: fig. 3; pl. 253: figs. 1.

Type-locality: Unknown.

Distribution: Bahamas and Florida Keys; Guadeloupe; Puerto Rico (Powers, 1977).

Pitho quadridentata (Miers, 1879)

Description: Rathbun, 1925:369, pl. 132: fig. 2, pl. 133: fig. 2; pl. 250: fig. 10.

Type-locality: West Indies.

Distribution: Jamaica; Puerto Rico; Content Keys, Monroe County, Florida, 5-6 m (personal communication, D. K. Camp).

Podochela curvirostris (A. Milne Edwards, 1879)

Description: Rathbun, 1925:58, pls. 19, 210.

Type-locality: Barbados, 180 m, and near Havana, 229 m.

Distribution: Florida Straits; north coast of Cuba; Caribbean coast of Yucatan; Montserrat; Barbados; Grenadines (Powers, 1977).

Podochela gracilipes Stimpson, 1871

Description: Williams, 1984:301, fig. 236, 241j.

Type-locality: West of Tortugas, off Pacific and Carysfort Reefs (Florida), 66 to 110 m.

Distribution: Off Cape Lookout, North Carolina, through Gulf of Mexico and Caribbean Sea to Santa Catarina, Brazil (Williams, 1984).

Podochela lamelligera (Stimpson, 1871)

Description: Rathbun, 1925:52, pl. 20: figs. 1-2.

Type-locality: Off Tennessee Reef, Florida Keys, 38 m.

Distribution: Southeast Florida; off Key West, Florida; off northwest Florida (Powers, 1977).

Podochela macrodera Stimpson, 1860

Description: Rathbun, 1925:44, fig. 11, pl. 16.

Type-locality: St. Thomas and Key Biscayne, Florida.

Distribution: Bahamas; Florida Keys; west coast of Florida; off Caribbean coast of Yucatan; Cuba; Puerto Rico; Virgin Islands; Guadalupe; Curaçao, Netherlands Antilles; Brazil (Powers, 1977).

Podochela riisei Stimpson, 1860

Description: Williams, 1984:302, figs. 237, 241k.

Type-locality: Island of St. Thomas (West Indies).

Distribution: North Carolina to Campeche, Mexico; through West Indies to Trinidad; Rio de Janeiro, Brazil; Bermuda (Williams, 1984).

Podochela sidneyi Rathbun, 1924

Description: Williams, 1984:302, figs. 238, 241l.

Type-locality: Off Cape Hatteras, North Carolina, 90 m.

Distribution: Off Cape Hatteras, North Carolina, to Veracruz (Ray, 1974); northwestern Cuba; Yucatan Channel (Williams, 1984).

Pyromaia arachna Rathbun, 1924

Description: Rathbun, 1925:131, pls. 42-43.

Type-locality: Gulf of Mexico, SW of Cape San Blas, Florida; 309 m.

Distribution: Off South Carolina; off west coast of Florida to off east coast of Mexico, throughout the Gulf of Mexico.

Pyromaia cuspidata Stimpson, 1871

Description: Williams, 1984:303, figs. 239, 241m.

Type-locality: Off Sand Key, 150 m; Alligator Reef, 170 m; the Samboes, 170 and 221 m; southwest of Sand Key, 229 m (Florida).

Distribution: Off Cape Lookout, North Carolina, to west Florida; Cuba and Yucatan Channel to off Nicaragua 14°31'N, 80°41'W (Williams, 1984).

Rochinia crassa (A. Milne Edwards, 1879)

Description: Williams, 1984:322, figs. 256, 260a.

Type-locality: Between Cuba and Florida, 24°15'N, 82°13'W.

Distribution: Nantucket Shoals,

Massachusetts, to Gulf of Mexico off southern Texas; northern Cuba; west of Cabo de la Vela, Colombia; off French Guiana (Williams, 1984).

Rochinia hystrix (Stimpson, 1871)

Description: Rathbun, 1925:214, pls. 70, 71.

Type-locality: Off Sand Key, Florida, 24°16'N, 81°42'W, 252 m.

Distribution: Off Key West (Rathbun, 1925); off Cuba (Chace, 1940b); Gulf of Mexico.

Rochinia tanneri (Smith, 1883)

Description: Williams, 1984:323, figs. 257, 260b.

Type-locality: Off Delaware Bay.

Distribution: Off Martha's Vineyard, Massachusetts, to Straits of Florida (Williams, 1984).

Rochinia umbonata (Stimpson, 1871)

Description: Williams, 1984:323, figs. 258, 260c.

Type-locality: Off Sand Key, Florida.

Distribution: Southeast of Cape Lookout, North Carolina, through eastern and northern Gulf of Mexico to northeast of Nicaragua; through West Indies to St. Vincent (Williams, 1984).

Sphenocarcinus corrosus A. Milne Edwards, 1875

Description: Williams, 1984:314, figs. 250, 259c.

Type-locality: Off Barbados, 180 m.

Distribution: Off Cape Lookout, North Carolina; Gulf of Mexico (Goeke and Shaw, 1980) to Barbados (Williams, 1984).

Stenocionops furcata coelata (A. Milne Edwards, 1878)

Description: Williams, 1984:338, figs. 273, 275i.

Type-locality: Ten mi. from Jolbos Islands (Yucatan), and near Havana (Cuba), 320 m.

Distribution: Shelly reefs off Beaufort, North Carolina, to northwest Florida and Alabama; Yucatan Channel; West Indies to Barbados, (Williams, 1984).

Stenocionops furcata furcata (Olivier, 1791)

Description: Rathbun, 1925:449, text-fig. 131,

pls. 160, 161.

Type-locality: Unknown.

Distribution: Georgia; Florida (location unspecified); ?Gulf of Mexico; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Dominica; Barbados; French Guiana; Paraiba to Rio de Janeiro, Brazil; South Africa (Powers, 1977).

Stenocionops spinimana (Rathbun, 1892)

Description: Williams, 1984:339, figs. 274, 275j.

Type-locality: Off Cape Lookout, North Carolina, 227m.

Distribution: Off Cape Hatteras, North Carolina, to Florida Straits and Gulf of Mexico off Mobile Bay, Alabama, and east of Chandeleur Island, off Mississippi (Franks et al., 1972; Williams, 1984).

Stenocionops spinosissima (Saussure, 1857)

Description: Rathbun, 1925:445, pl. 165: fig. 2, pl. 264: figs. 374, pl. 265.

Type-locality: Guadeloupe.

Distribution: North Carolina; south and southwest Florida; off Texas and east coast of Mexico; north coast of Cuba; Haiti; Guadeloupe; Dominica; Rio de Janeiro and Fernando de Noronha, Brazil (Powers, 1977).

Stenorhynchus seticornis (Herbst, 1788)

Description: Williams, 1984:304, fig. 240, 241o.

Type-locality: Guadeloupe (Holthuis 1959).

Distribution: North Carolina to Santa Catarina, Brazil; Bermuda (Williams, 1984).

Stilbomastax margaritifera (Monod, 1939)

Description: Monod, 1939:561, figs. 6-9.-- Williams et al., 1977:887.

Type-locality: Basse-Terre, Guadeloupe, 15-20 m.

Distribution: southeast of Cape San Blas to southeast Florida; Guadeloupe (Williams et al., 1977).

Thoe puella Stimpson, 1860

Description: Rathbun, 1925:348, figs. 111, 112, pl. 125: figs. 1, 2.

Type-locality: Tortugas, Florida.

Distribution: Florida Keys and Dry Tortugas; Jamaica; Puerto Rico; St. Thomas; Guadeloupe; Curacao (Powers, 1977).

Tyche emarginata White, 1847

Description: Williams, 1984:312, figs. 247-

248, 259b.

Type-locality: West Indies.

Distribution: Off Beaufort Inlet, North Carolina; through Bahamas to west coast of Florida (Williams, 1984).

FAMILY PARTHENOPIDAE

Cryptopodia concava Stimpson, 1871

Description: Williams, 1984:346, figs. 281, 286a.

Type-locality: Off Conch Reef (Florida), 62.2 m.

Distribution: Southeast of Cape Lookout, North Carolina; central east Florida; Cape San Blas, Florida, to St. Thomas; Ceara to Bahia, Brazil (Williams, 1984).

Heterocrypta granulata (Gibbes, 1850)

Description: Williams, 1984:347, figs. 282, 286b.

Type-locality: Near Kiawah Island, Sullivan's Island, and White Point Shoal, Charleston Harbor, South Carolina.

Distribution: Nantucket Sound, Massachusetts, around peninsular Florida to southern Texas; through West Indies to Trinidad; Ceara to Bahia, Brazil (Williams, 1984).

Leiolambrus nitidus Rathbun, 1901

Description: Rathbun, 1925:545, pl. 199; pl. 281: fig. 1.

Type-locality: Mayaguez Harbor, Puerto Rico, 22-33 m.

Distribution: Gulf of Mexico, from off Alabama to south Texas; Jamaica; Puerto Rico; French Guiana (Powers, 1977).

Mesorhoea sexspinosa Stimpson, 1871

Description: Williams, 1984:348, figs. 283, 286c.

Type-locality: Four mi. southwest of Loggerhead Key, Florida, 20 m.

Distribution: Southeast of Cape Lookout, North Carolina; off northeast Florida, to Flanagan Passage, Virgin Islands (Williams, 1984).

Parthenope agona (Stimpson, 1871)

Description: Williams, 1984:342, figs. 276, 280a.

Type-locality: Off the Marquesas, Carysfort Reef, and Conch Reef, 73 and 89.6 m (southern Florida).

Distribution: Off Capes Hatteras and Lookout, North Carolina, and central eastern Florida;

Gulf of Mexico and Pensacola, Florida, to near Ft. Myers; through Florida Straits, West Indies and Caribbean Sea to Surinam (Williams, 1984).

Parthenope fraterculus (Stimpson, 1871)

Description: Williams, 1984:343, figs. 277, 280b.

Type-locality: Off Sand Key, Caryfort and Conch Reefs, west of Tortugas, 47.6-124.4 m (southern Florida).

Distribution: Off Cape Fear, North Carolina; central eastern Florida southward; Gulf of Mexico, off Cape San Blas, Florida, to Florida Straits; off Cape Catoche, Yucatan, Mexico; through West Indies to Mouth of Amazon River (Williams, 1984).

Parthenope granulata (Kingsley, 1879)

Description: Williams, 1984:344, figs. 278, 280c.

Type-locality: Tortugas, Florida.

Distribution: Off the three North Carolina Capes southward around Florida to Louisiana; Bermuda; Bahia Honda, Cuba(?); St. Thomas, Virgin Islands (Gore, 1977).

Parthenope pourtalesii (Stimpson, 1871)

Description: Williams, 1984:345, figs. 279, 280d.

Type-locality: Off Conch Reef, French Reef, and American Shoal (southern Florida) 73-214 m.

Distribution: Off Martha's Vineyard Massachusetts; New Jersey southward; Gulf of Mexico through West Indies to Grenada.

Parthenope serrata (H. Milne Edwards, 1834)

Description: Gore and Scotto, 1979:57, figs. 24 e-h, 25 B, 26.

Type-locality: "L' Océan Indien" by original designation; erroneous locality for the West Indies. Veracruz, Mexico by selection of male lectotype (Gore, 1977).

Distribution: Bermuda; Ft. Pierce, Florida, around the Gulf of Mexico; Central America; Cuba; Lesser Antilles; islands off the northern coast of South America, southward to Bahia, Brazil (Gore and Scotto, 1979).

Solenolambrus decemspinosus Rathbun, 1894

Description: Rathbun, 1925:540, pl. 194: figs. 1, 2.

Type-locality: Off Cape San Blas, Florida, 28°44'N, 85°16'W, 110 m.

Distribution: Northeastern Gulf of Mexico and off San Juan, Puerto Rico.

Solenolambrus typicus Stimpson, 1871

Description: Williams, 1984:349, figs. 285, 286e.

Type-locality: Off the Samboes and off Alligator Reef (southern Florida), 146.3 to 201.2 m.

Distribution: SE Cape Lookout; western Gulf of Mexico off Corpus Christi, Texas, and N of Yucatan; Swan Island and Nicaragua Shelf; southern Florida through West Indies to Surinam and Brazil (Gore and Scotto, 1979).

Solenolambrus tenellus Stimpson, 1871

Description: Williams, 1984:348, figs. 284, 286d.

Type-locality: Off Carysfort, Conch, and French Reefs, 64-89.6 m (southern Florida).

Distribution: Off Cape Lookout, North Carolina; central east Florida southward; Gulf of Mexico, near Cape St. George, Florida, to Florida Keys; Bahamas; Barbados (Williams, 1984).

Tutankhamen cristatipes (A. Milne Edwards, 1880)

Description: Rathbun, 1925:530, pl. 277: figs. 3-5.

Type-locality: St. Vincent, Lesser Antilles.

Distribution: Pourtales Plateau, Florida Straits; St. Vincent, Lesser Antilles.

FAMILY ATELECYCLIDAE

Trichopeltarion nobile A. Milne Edwards, 1880

Description: Rathbun, 1930:168, pl. 73.--Pequegnat, 1970:184, figs. 6-4, 6-5.

Type-locality: Off St. Lucia, 276 m.

Distribution: Off northwest Florida and Mississippi; east coast of Mexico; Bay of Campeche; off St. Lucia (Powers, 1977).

FAMILY CANCRIDAE

Cancer borealis Stimpson, 1859

Description: Williams, 1984:351, fig. 287.

Type-locality: Nova Scotia to Cape Cod.

Distribution: Nova Scotia to south of Tortugas, Florida; Verrill (1908) judged that a Bermuda record was probably mislabelled (Williams, 1984).

***Cancer irroratus* Say, 1817**

Description: Williams, 1984:353, fig. 288.
 Type-locality: "Inhabits the ocean" (Atlantic coast of the United States).
 Distribution: Labrador to off Miami, Florida (Williams, 1984).

FAMILY GERYONIDAE

***Geryon fenneri* Manning and Holthuis, 1984**
 Description: Manning and Holthuis, 1984:666, figs. 1, 2a, b, 3a-c, 4a, b.
 Type-locality: Off Fernandina, Florida, Albatross Stn. 2669.
 Distribution: Around coasts of Florida.

FAMILY PORTUNIDAE

***Arenaeus cribrarius* (Lamarck, 1818)**
 Description: Williams, 1984:362, fig. 292.
 Type-locality: Brazil.
 Distribution: Vineyard Sound, Massachusetts, to Santa Catarina, Brazil (Williams, 1984).

***Bathynectes longispina* Stimpson, 1871**
 Description: Rathbun, 1930:28, pls. 9, 10.-- Manning and Holthuis, 1981:80.
 Type-locality: Off Sand Key, Key West, and American Shoal, all in the Florida Straits, 183-275 m.
 Distribution: Off Martha's Vineyard, Massachusetts, to Gulf Stream in Florida Straits.

***Benthochason schmitti* Rathbun, 1931**
 Description: Rathbun, 1931:125, pls. 1, 2.-- Pequegnat, 1970:187, fig. 6-6.
 Type-locality: South of Loggerhead Key, Tortugas, Florida, 329 m.
 Distribution: Off Dry Tortugas; deep waters off Mississippi to Texas; off southern Gulf coast of Mexico; recently found off New England (Powers, 1977).

***Callinectes bocourti* A. Milne Edwards, 1879**
 Description: Williams, 1984:365, figs. 293f, 294.
 Type-locality: Mullins River, 20 mi. south of Belize, (British) Honduras.
 Distribution: Jamaica and Belize to Santa Catarina, Brazil; Florida, Mississippi, North Carolina, United States of America (Williams, 1974; Perschbacher and Schwartz, 1979; Williams and Williams, 1981).

***Callinectes danae* Smith, 1869**
 Description: Williams, 1984:367, figs. 293d,

295.

Type-locality: Recife [=Pernambuco, Estado de Pernambuco], Brazil.
 Distribution: Bermuda; New Hanover County, North Carolina, near Cape Fear, rare (Perschbacher and Schwartz, 1979); southern Florida and eastern side of Yucatan Peninsula to Estado de Santa Catarina, Brazil.

***Callinectes exasperatus* (Gerstaecker, 1856)**
 Description: Williams, 1984:369, figs. 293e, 296.
 Type-locality: Puerto Cabello, Venezuela.
 Distribution: Duval County, east of Jacksonville, Florida (rarely) to Santa Catarina, Brazil; Veracruz, Mexico; Bermuda; also reported from extreme southern Texas (Williams, 1984).

***Callinectes larvatus* Ordway, 1863**
 Description: Williams, 1984:371, figs. 293a, 297.
 Type-locality: Key West, Florida; Tortugas; Bahama Islands; Haiti.
 Distribution: Beaufort, North Carolina, through Caribbean Sea to south central Brazil off São Paulo; Bermuda. North Carolina records rare (Williams, 1974; Perschbacher and Schwartz, 1979). Florida (intertidal) (personal communication, P. M. Mikkelsen).

***Callinectes ornatus* Ordway, 1863**
 Description: Williams, 1984:373, figs. 293c, 298.
 Type-locality: Charleston, South Carolina; Gonaives, Haiti; Cumana, Venezuela; Tortugas and Bahamas also listed in original description.
 Distribution: Bermuda; Virginia, North and South Carolina through southern Florida; northwestern Yucatan to Estado de São Paulo, Brazil (Williams, 1984).

***Callinectes sapidus* Rathbun, 1896**
 Description: Williams, 1984:376, figs. 293g, 299.
 Type-locality: East coast of United States.
 Distribution: Occasionally Nova Scotia, Maine, and northern Massachusetts, to northern Argentina, Bermuda, and the Antilles; Resund, Denmark; the Netherlands and adjacent North Sea; northwest and southwest France; Golfo di Genova; northern Adriatic; Aegean, western Black, and eastern Mediterranean Sea; Lake Hamana-ko, central Japan (Williams, 1984).

Callinectes similis Williams, 1966

Description: Williams, 1984:383, figs. 293b, 300.

Type-locality: Off beach between St. Johns River jetties and Jacksonville Beach, Florida.

Distribution: Off Delaware Bay to Key West, Florida; northwestern Florida around Gulf of Mexico to off Campeche, Yucatan; also Isla de Providencia, Colombia; reported from northern Jamaica (Norse, 1978; Williams, 1984).

Cronius ruber (Lamarck, 1818)

Description: Williams, 1984:385, fig. 301.

Type-locality: Brazil.

Distribution: Vicinity of Little Egg Inlet, New Jersey (Milstein et. al., 1977); Rehoboth Bay, Delaware; Virginia (rare, Van Engel and Sandifer, 1972); South Carolina to Santa Catarina, Brazil; Baja California to Peru; Clipperton, Galapagos Island; West Africa from Mauritania to Angola; Cape Verde, Principe, São Tomé and Annobon Islands (Williams, 1984).

Cronius tumidulus (Stimpson, 1871)

Description: Rathbun, 1930:142, pl. 64.

Type-locality: West of Tortugas, 68 m and off Conch Reef, 73 m.

Distribution: Bermuda; Bahamas; Florida Keys and Dry Tortugas; west coast of Florida; north and south coasts of Cuba; Jamaica; Puerto Rico; Virgin Islands; Netherlands Antilles; Old Province Island (Caribbean); Ceara to Bahia, Brazil (Powers, 1977).

Ovalipes floridanus Hay and Shore, 1918

Description: Turkay, 1971:139, fig. 3.

Type-locality: Pensacola, Florida.

Distribution: Southwest Florida to south Texas.

Ovalipes stephensoni Williams, 1976

Description: Williams, 1984:361, fig. 291.

Type-locality: South of Beaufort Inlet, North Carolina, 31°11'N, 76°42'W, 35 m.

Distribution: Off Accomack County, Virginia, 37°31'N, to near Biscayne Bay, Florida.

Portunus anceps (Saussure, 1858)

Description: Williams, 1984:387, fig. 302.

Type-locality: Cuba.

Distribution: Cape Hatteras, North Carolina (Park, 1978), to Bahia, Brazil; Bermuda (Williams, 1984).

Portunus binoculus Holthuis, 1969

Description: Holthuis, 1969:409, fig. 1.

Type-locality: Straits of Florida.

Distribution: Bahamas; Florida Straits; north coast of Cuba; east of Yucatan, in Caribbean Sea; off Caribbean coasts of Panama and Colombia (Powers, 1977).

Portunus depressifrons (Stimpson, 1859)

Description: Williams, 1984:387, fig. 303.

Type-locality: South Carolina and Florida Keys.

Distribution: Fort Macon, North Carolina (Coues, 1871; Kingsley, 1878-79), through northwest Florida to Bay of Campeche and Caribbean Sea; Bermuda (Williams, 1984).

Portunus floridanus Rathbun, 1930

Description: Williams, 1984:388, fig. 304.

Type-locality: Off Key West, Florida, 24°25'45"N, 81°48'00"W.

Distribution: East Cape Lookout, North Carolina, to Honduras and Nicaragua, through West Indies and northern South America to Surinam (Williams, 1984).

Portunus gibbesii (Stimpson, 1859)

Description: Williams, 1984:389, fig. 305.

Type-locality: South Carolina and St. Augustine, Florida.

Distribution: Southern Massachusetts through Gulf of Mexico along coast to French Guiana, but reported absent from the Antilles (Park, 1978; Williams, 1984).

Portunus ordwayi (Stimpson, 1860)

Description: Williams, 1984:390, fig. 306.

Type-locality: Key Biscayne and Tortugas, Florida; St. Thomas (Virgin Islands).

Distribution: Vineyard Sound, Massachusetts; North Carolina through Gulf of Mexico, West Indies and Caribbean Sea to near Rio de Janeiro, Brazil (Park, 1978); Bermuda; Fernando de Noronha (Williams, 1984).

Portunus sayi (Gibbes, 1850)

Description: Williams, 1984:391, fig. 307.

Type-locality: South Carolina.

Distribution: North Atlantic Ocean from Nova Scotia through Gulf of Mexico to the Guianas; Bermuda; mid-Atlantic Ocean; Canary Islands and Morocco. The only record from Brazil is that of Gerstaecker for his *Lupea pudica* (=sayi), and modern collections have not confirmed this (Williams, 1984).

Portunus sebae (H. Milne Edwards, 1834)
Description: Rathbun, 1930:79, plates 34, 35.
Type-locality: Brazil.
Distribution: Bermuda; Florida Keys and Straits; Dry Tortugas; south coast of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Dominica; Netherlands Antilles (Powers, 1977).

Portunus spinicarpus (Stimpson, 1871)
Description: Williams, 1984:392, fig. 308.
Type-locality: Straits of Florida south of Dry Tortugas, 24°23'N, 82°57'W to 24°24'N, 82°56'W, (Holthuis, 1969, restricted).
Distribution: East southeast Oregon Inlet, North Carolina, 35°42'00"N, 74°54'30"W (Musick and McEachren, 1972) to Santa Catarina, Brazil (Williams, 1984).

Portunus spinimanus Latreille, 1819
Description: Williams, 1984:393, fig. 309.
Type-locality: American waters, common in Brazil.
Distribution: New Jersey through Gulf of Mexico and West Indies to Santa Catarina, Brazil; Bermuda (Williams, 1984).

Portunus ventralis (A. Milne Edwards, 1879)
Description: Rathbun, 1930:43, pl. 13: figs. 1, 2.
Type-locality: Guadeloupe.
Distribution: Georgia to east coast of Florida; Dry Tortugas; ?Texas; north and west coasts of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Barbados; Rio Grande do Norte to Rio de Janeiro, Brazil (Powers, 1977).

Portunus vocans (A. Milne Edwards, 1878)
Description: Rathbun, 1930:10, figs. 8, 9, pl. 25.
Type-locality: Cape Verde Islands.
Distribution: North coast of Cuba; between Jamaica and Haiti; Ascension Island, in the south Atlantic Ocean (Powers, 1977). Florida (personal communication, P. A. McLaughlin).

FAMILY GONEPLACIDAE

Chacellus filiformis Guinot, 1969
Description: Guinot, 1969:722, figs. 135-136, pl. 5: fig. 4.
Type-locality: Gulf of Mexico.
Distribution: Between Bahamas and east coast of Florida; off northwest Florida (Powers, 1977).

Eucratopsis crassimanus (Dana, 1852)
Description: Rathbun, 1918:52, fig. 22, pl. 12: fig. 3, pl. 159: figs. 1-2.--Guinot, 1969:258, figs. 6, 10, 25.
Type-locality: Rio de Janeiro.
Distribution: Florida Keys; south and west coasts of Florida; Yucatan; Jamaica; Bahia to Rio de Janeiro, Brazil (Powers, 1977).

Euphrosynoplax clausa Guinot, 1969
Description: Guinot, 1969:720, figs. 127, 139, pl. 4: fig. 3.--Pequegnat, 1970:194.
Type-locality: Florida, Tortugas.
Distribution: Dry Tortugas; off Alabama and Mississippi; Campeche, Yucatan, (91 to 210 m) (Powers, 1977).

Euryplax nitida Stimpson, 1859
Description: Williams, 1984:432, fig. 343.
Type-locality: Florida Keys.
Distribution: Off Beaufort, North Carolina, to Heald Bank, Texas; West Indies to St. Thomas; Bermuda; specimen from "Bresil, Dertero" [sic] (=Florianopolis?) figured by Guinot, 1969b (Williams, 1984).

Frevillea barbata A. Milne Edwards, 1880
Description: Rathbun, 1918:26, pl. 4: figs. 1, 3, pl. 5.--Guinot, 1969:513, pl. 2: fig. 2.
Type-locality: 23°13'N; 89°16'W, 154 m. Stn. 36, Blake.
Distribution: West coast of Florida; Yucatan (Gulf); north coast of Cuba; off Grenada, (55 to 168 m) (Powers, 1977).

Frevillea hirsuta (Borradaile, 1916)
Description: Williams, 1984:432, fig. 344.
Type-locality: Off Rio de Janeiro, 72 m.
Distribution: North Carolina to Rio de Janeiro, Brazil (Williams, 1984).

Glyptoplax smithii A. Milne Edwards, 1880
Description: Williams, 1984:434, fig. 346.
Type-locality: Reefs west of Florida, 23.8 m.
Distribution: From Cape Hatteras, North Carolina, to Gulf of Mexico and Yucatan Channel (Williams, 1984).

Goneplax sigsbei (A. Milne Edwards, 1880)
Description: Williams, 1984:433, fig. 345.
Type-locality: Grenada.
Distribution: East Cape Fear, North Carolina, 33°56'N, 76°26'W, to 33°55.3' N, 76°28.8'W, 130-120 m, Eastward Stn. 3213; Grenada,

11°27'N, 62°11'W, and 11°25'00"N,
62°04'15"W (Williams et al., 1968).

Nanoplax xanthiformis (A. Milne Edwards, 1881)

Description: Williams, 1984:436, fig. 348.
Type-locality: Off Grenada, 168.3 m.
Distribution: Cape Hatteras, North Carolina; through Gulf of Mexico and West Indies to Cabo Frio, Rio de Janeiro, Brazil (Williams, 1984).

Neopilumnoplax americana (Rathbun, 1898)

Description: Rathbun, 1918:21, figs. 5-6.--
Guinot, 1969:689, figs. 83-84.
Type-locality: Off Georgia, 792 m.
Distribution: Off North Carolina and Georgia; Florida Keys and Straits; north coast of Cuba; Guadeloupe; Espirito Santo, Brazil; Arabian Sea (Powers, 1977).

Panoplax depressa Stimpson, 1871

Description: Williams, 1984:435, fig. 347.
Type-locality: East and Middle Keys, Tortugas, (Florida), 9.1 to 12.8 m.
Distribution: Southeast of Cape Lookout, North Carolina; off Jacksonville and Cape San Blas, Florida, through West Indies to Barbados (Williams, 1984).

Pilumnoplax elata (A. Milne Edwards, 1880)

Description: Guinot, 1969:688.
Type-locality: West Florida, 23.4 m.
Distribution: Only from the type-locality.

Pseudorhombila quadridentata (Latreille, 1828)

Description: Hernandez, 1982:1, figs. 1e, 1d, 2c, 3c, 4c, 5c, 6c.
Type-locality: Unknown (Guinot, 1969).
Distribution: Specimens are known from southern Florida (Tortugas and northwest of New Grounds Shoal Light) Louisiana (west Delta lease area) and south of Lobos Islands, Mexico; Puerto Rico (North of Arecibo).

Sotoplax robertsi Guinot, 1984

Description: Guinot, 1984:92, figs. 1-3.
Type-locality: Gulf of Mexico, middle shelf region off Apalachicola Bay, lat. 28°30' long. 84°58', *Tursiops*, cruise T-7109, Stn. 4, 54 m.
Distribution: Only from the type locality.

Speocarcinus lobatus Guinot, 1969

Description: Guinot, 1969:710, figs. 124-125,

pl. 4: fig. 2.--Felder 1973:70, pl. 10, fig. 3.

Type-locality: Sabine Pass, Texas.

Distribution: Dry Tortugas; off Louisiana and Texas (Powers, 1977).

Thalassaplax angusta Guinot, 1969

Description: Guinot, 1969:717; figs. 131-132, pl. 4: fig. 2.--Pequegnat, 1970:192.
Type-locality: Southwest of Cape San Blas, Florida, *Albatross*, Stn. 2402.
Distribution: East coast of Florida; off northwest Florida, Alabama and Mississippi; off east coast of Mexico; off Campeche, Yucatan (Powers, 1977).

Trapezioplax tridentata (A. Milne Edwards, 1880)

Description: Guinot, 1969:713, figs. 128-129, 142.
Type-locality: Barbados, 13.5-90 m.
Distribution: Florida Keys and Dry Tortugas; west coast of Florida; Barbados (Powers, 1977).

FAMILY XANTHIDAE

Actaea acantha (H. Milne Edwards, 1834)

Description: Rathbun, 1930:261, pl. 105: fig. 5, pl. 106: fig. 1, 2.
Type-locality: Unknown.
Distribution: Bahamas; Florida Keys and Dry Tortugas; northwest coast of Cuba; Jamaica; Haiti; Puerto Rico; Guadeloupe; St. Bartholomew; Fernando de Noronha, Brazil (Powers, 1977).

Actaea bifrons Rathbun, 1898

Description: Rathbun, 1930:255, fig. 41, pl. 104: figs. 3-6.
Type-locality: Colon, Panama.
Distribution: Key West, Florida; Puerto Rico; Virgin Islands; St. Bartholomew; Barbados; Curaçao; Colon, Panama (Powers, 1977).

Allactaea lithostrota Williams, 1974

Description: Williams, 1984:397, figs. 311, 331a.
Type-locality: Southeast Cape Lookout, North Carolina, 33°43'N, 76°40.2'W, 90 m to 33°42.7'N, 76°40.2'W, 110m, *Eastward* Stn. 1087.
Distribution: Near edge of continental shelf southeast of Cape Lookout, North Carolina; Florida Straits; off Cape Catoche, Yucatan; off Venezuela and Surinam; Bermuda (Markham and McDermott, 1981; Williams, 1984).

Banareia palmeri (Rathbun, 1894)

Description: Rathbun, 1930:260, pl. 106, fig. 3-6.

Type-locality: Rodriguez Creek, Florida.

Distribution: Bahamas; east coast of Florida; Florida Keys; north coast of Cuba; Haiti; Virgin Islands; Curaçao (Powers, 1977).

Carpilius corallinus (Herbst, 1783)

Description: Rathbun, 1930:240, pls. 97-99.

Type-locality: Unknown.

Distribution: Bermuda; Bahamas; West Flower Garden Bank, off Texas; north coast of Cuba; Jamaica; Puerto Rico; Virgin Islands; Guadeloupe; Dominica; Curaçao; Old Providence Island (Caribbean); Pernambuco and Ceara, Brazil (Powers, 1977). Monroe, County, Florida, lobster trap (personal communication, D. K. Camp).

Carpoporus papulosus Stimpson, 1871

Description: Williams, 1984:399, figs. 313, 331c.

Type-locality: Southwest of Tortugas and off Carysfort Reef, (Florida).

Distribution: Between Capes Hatteras and Lookout, North Carolina; Gulf of Mexico off Mobile Bay southeastward; Cape Catoche, Yucatan (Williams, 1984).

Cataleptodius floridanus (Gibbes, 1850)

Description: Rathbun, 1930:297, pl. 137: figs. 1, 2, pl. 138: fig. 1-- Guinot, 1968:706, figs. 20, 23, 29.

Type-locality: Key West, Florida.

Distribution: Bermuda; Bahamas; Florida Keys and Dry Tortugas; northwest coast of Florida; north coast of Cuba; Jamaica; Puerto Rico; Virgin Islands; Antigua; Barbados; Curaçao; Panama to Colombia (Caribbean coasts; Abolhos Islands to São Paulo, Brazil) (Powers, 1977).

Chlorodiella longimana (H. Milne Edwards, 1834)

Description: Rathbun, 1930:462, pl. 186.

Type-locality: Puerto Rico.

Distribution: Florida to Curaçao and Barbados; West Africa. Puerto Rico, Culebra, St. Thomas, St. Croix (Rathbun, 1933).

Domecia acanthophora acanthophora

(Desbonne and Schramm, 1867)

Description: Williams, 1984:417, figs. 330, 331q.

Type-locality: Guadeloupe.

Distribution: Bermuda; Cape Lookout Shoals, North Carolina, NW Gulf of Mexico through West Indies and Caribbean Sea to Alagoas, Brazil (Williams, 1984). Florida (5-6 m) (personal communication, P. M. Mikkelsen).

Eriphia gonagra (Fabricius, 1781)

Description: Williams, 1984:419, figs. 332, 333a-c.

Type-locality: Jamaica.

Distribution: North Carolina to Patagonia; Bermuda (Williams, 1984).

Etisus maculatus (Stimpson, 1860)

Description: Rathbun, 1930:295, pl. 136.-- Guinot, 1969:234.

Type-locality: Tortugas, Florida.

Distribution: Florida Keys and Dry Tortugas; Bahamas; north coast of Cuba; Puerto Rico; Virgin Islands (Powers, 1977).

Eurypanopeus abbreviatus (Stimpson, 1860)

Description: Williams, 1984:407, figs. 322, 331i.

Type-locality: Barbados, British West Indies.

Distribution: South Carolina, through West Indies and Gulf of Mexico to Santa Catarina, Brazil (Williams, 1984).

Eurypanopeus depressus (Smith, 1869)

Description: Williams, 1984:408, figs. 323, 331j.

Type-locality: New Haven, Connecticut.

Distribution: Massachusetts Bay through Florida to southern Texas; Dutch West Indies; Uruguay; Bermuda (Williams, 1984).

Eurypanopeus dissimilis (Benedict and

Rathbun, 1891)

Description: Rathbun, 1930:411, fig. 66, pl. 173: figs. 1-2.

Type-locality: Trinidad.

Distribution: West coast of Florida; north coast of Cuba; Jamaica; Nicaragua; Trinidad; Brazil (Powers, 1977).

Eurypanopeus turgidus (Rathbun, 1930)

Description: Rathbun, 1930:364, pl. 166.

Type-locality: Chandeleur Islands, Louisiana.

Distribution: Northwest coast of Florida to Texas (Powers, 1977).

Eurytium limosum (Say, 1818)

Description: Williams, 1984:416, figs. 329,

331p.

Type-locality: "Inhabits shores of the Northern States".

Distribution: South Carolina to Louisiana through West Indies and Caribbean Sea to São Paulo, Brazil; Bermuda (Williams, 1984).

Glyptoxanthus erosus (Stimpson, 1859)

Description: Williams, 1984:398, figs. 312, 331b.

Type-locality: Florida.

Distribution: Cape Lookout, North Carolina, southward; off Grand Isle, Louisiana, southeastward; Yucatan; through West Indies to Guadeloupe (Williams, 1984).

Heteractaea ceratopus (Stimpson, 1860)

Description: Rathbun, 1930:530, pl. 212: figs. 5-8, pl. 213.--Guinot, 1968:721, figs. 50, 56.

Type-locality: Key Biscayne, Florida.

Distribution: Bahamas; east coast of Florida Keys and Dry Tortugas; north coast of Cuba; Curaçao; Trinidad; Barbados (Powers, 1977).

Hexapanopeus angustifrons (Benedict and Rathbun, 1891)

Description: Williams, 1984:415, figs. 327, 331n.

Type-locality: Long Island Sound.

Distribution: Vineyard Sound, Massachusetts, to Port Aransas, Texas; Bahamas; Jamaica (Williams, 1984).

Hexapanopeus caribbaeus (Stimpson, 1871)

Description: Rathbun, 1930:399, pl. 171: figs. 3-5.

Type-locality: St. Thomas.

Distribution: West Indies to state of Santa Catarina, Brazil; Puerto Rico, St. Thomas, (Rathbun, 1933). Florida (intertidal to 10 m) (personal communication, P. M. Mikkelsen).

Hexapanopeus hemphillii (Benedict and Rathbun, 1891)

Description: Rathbun, 1930:400, pl. 171: figs. 1, 2, 6.

Type-locality: Indian Key, Florida.

Distribution: Florida and West Indies; Puerto Rico; St. Thomas (Rathbun, 1933).

Hexapanopeus lobipes (A. Milne Edwards, 1880)

Description: Rathbun, 1930:329, fig. 50, pl. 155: figs. 3-5.--Menzies, 1948:23.

Type-locality: South of Florida, 24°43'N, 83°25'W, 68 m.

Distribution: Bahamas; off Key West, in Florida Straits; northwest of Dry Tortugas (Powers, 1977).

Hexapanopeus paulensis Rathbun, 1930

Description: Williams, 1984:416, figs. 328, 331o.

Type-locality: Santos, São Paulo, Brazil.

Distribution: South Carolina, through Gulf of Mexico to Uruguay (Milstein, et al., 1976).

Hexapanopeus quinquedentatus Rathbun, 1901

Description: Rathbun, 1930:402, fig. 62.

Type-locality: Mayaguez, Puerto Rico.

Distribution: Northwest Florida; Puerto Rico.

Leptodius parvulus (Fabricius, 1793)

Description: Rathbun, 1930:305, pl. 141: figs. 1-3; 1933:58, fig. 50.

Type-locality: Islands of South America.

Distribution: Bermuda; Bahamas; Florida Keys; Jamaica; Haiti; Puerto Rico; Barbados; Curaçao; Fernando de Noronha, Brazil (Powers, 1977).

Lobopilumnus agassizii (Stimpson, 1871)

Description: Williams, 1984:429, figs. 340g, 341.

Type-locality: Typical form: East and Middle Keys, Tortugas, Florida.

Distribution: North Carolina; eastern Gulf of Mexico; Yucatan; Cuba; Venezuela and Trinidad; Bermuda (Williams, 1984).

Melybia thalassia Stimpson, 1871

Description: Williams, 1984:430, fig. 342.

Type-locality: Off French Reef, 27.4 m, and west of Tortugas (southern Florida) 64-76.8 m.

Distribution: About 30 mi. south southeast Cape Lookout, North Carolina (34°11'N, 76°09'W); southwest of Mississippi River delta, through West Indies to Bahia, Brazil (Williams, 1984).

Menippe mercenaria (Say, 1818)

Description: Williams, 1984:420, figs. 333d, e, 334.

Type-locality: "The Southern States".

Distribution: Cape Lookout, North Carolina, to Yucatan, Mexico; Bahamas; Cuba; Jamaica (Williams, 1984).

***Menippe nodifrons* Stimpson, 1859**

Description: Rathbun, 1930:479, pl. 198: fig. 3; pl. 199.

Type-locality: Indian River, Florida.

Distribution: East coast of Florida; ?Louisiana; north and south coasts of Cuba; Jamaica; Virgin Islands; Trinidad; Caribbean coasts of Panama and Colombia; Paraiba to São Francisco do Sul, Brazil; Gabon, West Africa (Powers, 1977).

***Micropanope barbadensis* (Rathbun, 1921)**

Description: Rathbun, 1930:446, fig. 72.

Type-locality: Barbados.

Distribution: Dry Tortugas; Barbados.

***Micropanope lobifrons* A. Milne Edwards, 1880**

Description: Rathbun, 1930:429, pl. 178: figs. 4-6.

Type-locality: Off Montserrat, 161 m.

Distribution: South Florida, in Gulf Stream; Dry Tortugas; off northwest Florida; off north coast of Cuba; Puerto Rico; Virgin Islands; Santa Cruz Island (Caribbean); Grenada; Barbados; Colon, Panama (Powers, 1977).

***Micropanope nuttingi* (Rathbun, 1898)**

Description: Williams, 1984:404, figs. 318, 331g.

Type-locality: Bahama Banks.

Distribution: Cape Hatteras, North Carolina, through Gulf of Mexico and West Indies to Bahia, Brazil (Williams, 1984).

***Micropanope pusilla* A. Milne Edwards, 1880**

Description: Rathbun, 1930:431, pl. 179: figs. 7, 8.

Type-locality: Off west coast of Florida, 31 m.

Distribution: Dry Tortugas; northwest of Key West; west and northwest coasts of Florida; Alabama; north coast of Cuba; Jamaica; Puerto Rico; Virgin Islands (Powers, 1977).

***Micropanope sculptipes* Stimpson, 1871**

Description: Williams, 1984:405, fig. 319.

Type-locality: Seven hauls in Florida Keys, 27.4 to 124 m.

Distribution: SE Cape Lookout, North Carolina, to Port Aransas, Texas; West Indies to Barbados.

***Micropanope spinipes* A. Milne Edwards, 1880**

Description: Rathbun, 1930:443, fig. 71, pl.

181: figs. 1, 2.-- Pequegnat and Ray, 1974:238, figs. 18-22.

Type-locality: Abrolhos Islands, Brazil, 55 m.

Distribution: Bermuda; Bahamas; Florida Keys; West Flower Garden Bank, off Texas; Curacao; Alagoas and off the Abrolhos Islands, Brazil (Powers, 1977).

***Micropanope urinator* (A. Milne Edwards, 1881)**

Description: Williams, 1984:405, fig. 320.

Type-locality: Near Santa Cruz (St. Croix), West Indies, 448 m.

Distribution: Off Capes Hatteras and Lookout, North Carolina; Florida Keys to St. Croix, West Indies (Williams, 1984).

***Neopanope packardii* (Kingsley, 1879)**

Description: Abele, 1972b:269, figs. 1B, 3A.

Type-locality: Key West, Florida.

Distribution: Southeast and south Florida; Bahamas; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; Louisiana; north coast of Cuba (Powers, 1977).

***Neopanope sayi* (Smith, 1869)**

Description: Williams, 1984:409, figs. 324, 331k.

Type-locality: New Haven, Connecticut, and Cape Cod, Massachusetts.

Distribution: Miscou Harbor, mouth of Chaleur Bay, New Brunswick, Prince Edward Island, and Cape Breton Island, Nova Scotia, Canada (Bousfield 1956, Bousfield and Laubitz, 1972), to Florida Keys (Abele, 1972b).

***Neopanope texana* (Stimpson, 1859)**

Description: Abele, 1972b:266, figs. 1A, 2A, 2E, 3B, 3C.--Felder, 1973:68, pl. 9: fig. 19.

Type-locality: St. Joseph's Island, Texas.

Distribution: West coast of Florida (south as far as Charlotte County) to south Texas (Powers, 1977).

***Panopeus americanus* Saussure, 1857**

Description: Rathbun, 1930:357, pl. 164: figs. 3, 4, 6.

Type-locality: Guadeloupe.

Distribution: Bahamas; Florida Keys; west coast of Florida; north coast of Cuba; Jamaica; Dominican Republic; Puerto Rico; St. Thomas, Virgin Islands; Guadeloupe; Trinidad; Caribbean coast of Colombia; Rio Parahyba do Norte to Santa Catarina, Brazil (Powers, 1977).

Panopeus bermudensis Benedict and Rathbun, 1891

Description: Rathbun, 1930:360, fig. 56, pl. 165.

Type-locality: Bermuda.

Distribution: Bermuda; Bahamas; west coast of Florida; ?Texas; north coast of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Trinidad; Old Providence Island (Caribbean); Colombia to Santa Catarina, Brazil. In the eastern pacific, from Magdalena Bay, Mexico to Peru (Powers, 1977).

Panopeus harttii Smith, 1869

Description: Rathbun, 1930:355, pl. 164: figs. 1, 2, 5.

Type-locality: Abrolhos Reefs, Brazil.

Distribution: Florida Keys to State to São Paulo, Brazil, Puerto Rico, St. Thomas (Rathbun, 1933).

Panopeus herbstii H. Milne Edwards, 1834

Description: Williams, 1983:866, fig. 3.

Type-locality: "Inhabits oyster beds, & found on oysters (*O. virginica*) in our markets" [by implication the eastern United States] (Say, 1817:58). Holthuis's (1979) selection of the specimen figured by Say (1817, pl. 4, fig. 3) as the lectotype for *P. herbstii* restricts the nominal species to the common mud crab occurring on oyster bars of the eastern United States.

Distribution: Shallow intertidal and subtidal waters of the eastern United States from Boston Harbor, Massachusetts, to Indian River County, southeastern Florida (Williams, 1983).

Panopeus lacustris Desbonne, 1867

Description: Williams, 1983:868, fig. 11.

Type-locality: The lagoons of Guadeloupe, hiding under rocks.

Distribution: Shallow and subtidal waters from Bermuda and extreme southern Florida, through the West Indies, and along the continental margin of the Caribbean Sea and South America to Cabo Frio, Brazil. The species has been introduced in Hawaii, and, according to a report by Edmonson (1962), apparently has been known on the California coast for a number of years (Williams, 1983).

Panopeus obesus Smith, 1869

Description: Williams, 1983:873, figs. 6, 7.

Type-locality: Egmont Key (mouth of Tampa

Bay) Florida (restricted by Williams, 1983).

Distribution: Marsh edge, shallow intertidal, and subtidal waters of the Carolinian Province from environs of Beaufort, North Carolina to Georgia (and perhaps northeastern Florida), and from Sarasota County, Florida, to Louisiana; Texas and northeastern Mexico (Williams, 1983).

Panopeus occidentalis Saussure, 1857

Description: Williams, 1984:413, figs. 326, 331 m.

Type-locality: Guadeloupe.

Distribution: North Carolina to State of Santa Catarina, Brazil; Bermuda (Williams, 1984). Florida (intertidal) (personal communication, P. M. Mikkelsen).

Panopeus rugosus A. Milne Edwards, 1880

Description: Rathbun, 1930:353, pl. 162, 163.

Type-locality: Bahia. (Brazil).

Distribution: Florida Keys and Dry Tortugas; west and northwest coasts of Florida; north coast of Cuba; Haiti; Virgin Islands; Puerto Rico; Honduras to Nicaragua; Curaçao; Bahia to Santa Catarina, Brazil (Powers, 1977).

Panopeus simpsoni Rathbun, 1930

Description: Williams, 1983:875, fig. 8.

Type-locality: Saint George Sound, Apalachicola, Florida.

Distribution: Shallow intertidal and subtidal waters of the northern Gulf of Mexico: Key West, Florida; Lee County, Florida to Corpus Christi, Texas (Williams, 1983).

Paractaea rufopunctata nodosa (Stimpson, 1860)

Description: Williams, 1984:397, fig. 310.

Type-locality: Tortugas, Florida.

Distribution: Southeast Cape Lookout, North Carolina (34°12.2'N, 76°08'W, 90 m, to 34°12.27'N, 76°08'W, 50 m; 33°55.5'N, 76°28.4'W); off Mississippi River delta through West Indies to Rio de Janeiro, Brazil; Ascension Island (Williams, 1984).

Paraliomera dispar (Stimpson, 1871)

Description: Rathbun, 1930:244, fig. 38, pl. 101: figs. 4, 5.

Type-locality: Cruz del Padre, Cuba.

Distribution: Florida Keys to north coast of South America; Bermudas, Puerto Rico (Rathbun, 1933).

Paraliomera longimana (A. Milne Edwards, 1865)

Description: Rathbun, 1930:243, pl. 101: figs. 1-3.

Type-locality: Guadeloupe.

Distribution: Florida Keys and Dry Tortugas; Veracruz, Mexico; Puerto Rico; Virgin Islands; Barbados; Curaçao (Powers, 1977).

Pilumnoides nudifrons (Stimpson, 1871)

Description: Rathbun, 1930:538, pl. 218: figs. 1-2.

Type-locality: Off Sombrero Key, 203-229 m.

Distribution: Florida Straits and Keys; Barbados.

Pilumnus caribaeus Desbonne and Schramm, 1867

Description: Rathbun, 1930:491, pl. 200: figs. 3, 4.

Type-locality: Guadeloupe.

Distribution: Bahamas; Florida Keys; north coast of Cuba; Jamaica; Puerto Rico; Vieques and Culebra; Virgin Islands; Guadeloupe; Curaçao; Bahia to São Paulo, Brazil (Powers, 1977).

Pilumnus dasypodus Kingsley, 1879

Description: Williams, 1984:425, figs. 335, 340a.

Type-locality: Key West, Florida.

Distribution: Off Cape Hatteras, North Carolina, through Gulf of Mexico, Caribbean Sea and West Indies to Santa Catarina, Brazil (Williams, 1984).

Pilumnus floridanus Stimpson, 1871

Description: Williams, 1984:426, figs. 336, 340b.

Type-locality: Tortugas, (Florida).

Distribution: Off Cape Lookout, North Carolina, through Gulf of Mexico, and Yucatan Channel, to Honduras; through West Indies to Bahia, Brazil (Williams, 1984).

Pilumnus gemmatus Stimpson, 1860

Description: Rathbun, 1930:513, pl. 207: figs. 1-3.

Type-locality: St. Thomas and Tortugas.

Distribution: Dry Tortugas; Culebra; Virgin Islands; Curaçao (Powers, 1977).

Pilumnus holosericus Rathbun, 1898

Description: Rathbun, 1930:519, fig. 81, pl. 207: figs. 8, 9.

Type-locality: St. Thomas, Virgin Islands.

Distribution: Bahamas; Dry Tortugas; Puerto Rico; Virgin Islands; Trinidad; Curaçao (Powers, 1977).

Pilumnus lacteus Stimpson, 1871

Description: Williams, 1984:426, figs. 337, 340c.

Type-locality: Cruz del Padre, Cuba, and Key West, Florida.

Distribution: Near Beaufort, North Carolina, to Florida; Cuba (Williams, 1984).

Pilumnus longleyi Rathbun, 1930

Description: Rathbun, 1930:502, pl. 202: figs. 4-5.

Type-locality: South end of Loggerhead Key, Tortugas, Florida.

Distribution: Bahamas; Florida Keys and Dry Tortugas (Powers, 1977).

Pilumnus marshi Rathbun, 1901

Description: Rathbun, 1930:499, fig. 80.

Type-locality: St. Thomas, 37-55 m.

Distribution: Tortugas, Florida; St. Thomas, St. Croix.

Pilumnus nudimanus Rathbun, 1900

Description: Rathbun, 1930:523, fig. 82.

Type-locality: Arroyo, Puerto Rico.

Distribution: Known from the type-locality. Dry Tortugas, Florida, 29-33 m (personal communication, D. K. Camp).

Pilumnus pannosus Rathbun, 1896

Description: Williams, 1984:427, figs. 338, 340d.

Type-locality: Key West, Florida.

Distribution: Bogue Sound off Beaufort, North Carolina, to Port Aransas, Texas; West Indies to Virgin Islands (Williams, 1984).

Pilumnus sayi Rathbun, 1897

Description: Williams, 1984:428, figs. 339, 340e.

Type-locality: Georgia and east Florida.

Distribution: North Carolina through Gulf of Mexico and West Indies to Curaçao (Williams, 1984).

Pilumnus spinosissimus Rathbun, 1898

Description: Rathbun, 1930:494, fig. 79, pl. 200: figs. 7-8.

Type-locality: Off Key West, 10 m.

Distribution: Florida Keys and Dry Tortugas.

Platyactaea setigera (H. Milne Edwards, 1834)

Description: Rathbun, 1930:251, pl. 103.--Guinot, 1967:561, fig. 36.

Type-locality: Antilles.

Distribution: Bermuda; Bahamas; Florida Keys and Dry Tortugas; north coast of Cuba; Jamaica; Puerto Rico; Virgin Islands; Antigua; Barbados; Trinidad; Curaçao; Caribbean coast of Colombia (Powers, 1977).

Platypodiella spectabilis (Herbst, 1794)

Description: Rathbun, 1930:247, fig. 39, pl. 102: fig. 4.--Guinot, 1967:562.--Felder, 1973:65, pl. 9: fig. 10.

Type-locality: Unknown.

Distribution: Bermuda; Bahamas; Florida Keys; Texas; Veracruz, Mexico; Jamaica; Puerto Rico; Virgin Islands; Guadeloupe; Martinique; Barbados; Curaçao; Fernando de Noronha, Brazil (Powers, 1977).

Pseudomedeus agassizii (A. Milne Edwards, 1880)

Description: Williams, 1984:400, figs. 314, 331d.

Type-locality: Florida Reefs, 21.9-32.9 m.

Distribution: Cape Hatteras, North Carolina, to southern Texas (Williams, 1984).

Pseudomedeus distinctus (Rathbun, 1898)

Description: Williams, 1984:400, figs. 315, 331e.

Type-locality: Gulf of Mexico, northwest Dry Tortugas, 25°33'N, 84°21'W, 184.7 m.

Distribution: Off Cape Hatteras, North Carolina, 34°57'N, 75°19'W, through Straits of Florida to northwest of Dry Tortugas; Puerto Rico; Barbados (Williams, 1984).

Rhithropanopeus harrisii (Gould, 1841)

Description: Williams, 1984:401, figs. 316, 317, 331f.

Type-locality: Cambridge Marshes and Charles River, Massachusetts.

Distribution: The original range of this species is presumed to be in fresh to estuarine waters from the southwestern Gulf of St. Lawrence, Canada, to Veracruz, Mexico. The species has been introduced on the west coast of the United States and in parts of Europe (Williams, 1984).

Tetraxanthus bidentatus (A. Milne Edwards, 1880)

Description: Rathbun, 1930:459, pl. 185 (As

T. rugosus).--Chace, 1939:52.

Type-locality: Grenada, 168 m.

Distribution: Florida Keys; north and south coasts of Cuba; Grenada (Powers, 1977).

Tetraxanthus rathbunae Chace, 1939

Description: Williams, 1984:406, fig. 321.

Type-locality: Old Bahama Channel due north Punta Caldera, Camaguey Province, Cuba, 22°44'N, 78°41'W, 274-329 m.

Distribution: Off Cape Lookout, North Carolina, to Rio de Janeiro, Brazil (Coelho and Ramos, 1972), including Gulf of Mexico (Pequegnat, 1970; Williams, 1984).

Xantho denticulata White, 1847

Description: Monod, 1956:280, figs. 335-339.--Forest and Guinot, 1961:60, fig. 51.

Type-locality: West Indies.

Distribution: Bermuda; Bahamas; Florida Keys and Dry Tortugas; northwest Florida; Jamaica; Puerto Rico; Virgin Islands; Antigua; Barbados; Colon, Panama; Curaçao; Trinidad; Pernambuco to Abrolhos Islands, Brazil, Gulf of Guinea, west coast of Africa (Powers, 1977).

FAMILY GECARCINIDAE

Cardisoma guanhumi Latreille, 1825

Description: Rathbun, 1918:341, fig. 155, plates 106, 107.--Felder, 1973:79, pl. 12: figs. 1, 4.

Type-locality: Brazil.

Distribution: Bermuda; Bahamas; southeast Florida; Florida Keys; Louisiana and south Texas; eastern Mexico to Colombia; north and south coasts of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands to Barbados; Trinidad; Netherlands Antilles; Colombia to São Paulo, Brazil (Powers, 1977).

Gecarcinus lateralis (Fremenville, 1835)

Description: Rathbun, 1918:355, fig. 161, pls. 119-120.--Turkay, 1973:974, fig. 2.

Type-locality: Martinique, Guadeloupe, Marie Galante, Desirde and Iles do Saintes.

Distribution: Bermuda; Bahamas; southeast Florida; Florida Keys; south Texas to north coast of Yucatan; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; St. Thomas, Virgin Islands to Barbados; Netherlands Antilles; Honduras to Costa Rica; Caribbean coast of Colombia to Surinam (Powers, 1977).

***Gecarcinus ruricola* (Linnaeus, 1758)**

Description: Rathbun, 1918:352, text-fig. 160, pls. 117-118.--Chace and Hobbs, 1969:200, figs. 66, 67 h-j.

Type-locality: America.

Distribution: Bahamas; southeast Florida; north and south coasts of Cuba; Cayman Islands; Jamaica; Navassa Island (Caribbean); Hispaniola; Puerto Rico; St. Croix to Barbados; Curaçao; Old Providence and Swan Islands (Caribbean) (Powers, 1977).

FAMILY GRAPSIDAE***Araatus pisonii* (H. Milne Edwards, 1837)**

Description: Rathbun, 1918:323, pl. 96.--Chace and Hobbs, 1969:172, figs. 54, 58a.

Type-locality: Antilles.

Distribution: Bahamas; southeast to southwest Florida; north and south coasts of Cuba; New Province Island (Atlantic); Jamaica; Puerto Rico; Virgin Islands to Guadeloupe; Netherlands Antilles; Belize; Rio Parahyba do Norte to São Paulo, Brazil; in eastern Pacific, Nicaragua to Peru (Powers, 1977).

***Cyclograpsus integer* H. Milne Edwards, 1837**

Description: Rathbun, 1918:326, pl. 97: figs. 1, 2.--Chace and Hobbs, 1969:173, figs. 55, 58b-d.

Type-locality: Brazil.

Distribution: Bermuda; Bahamas; south Florida; Florida Keys; Texas; Cuba; Jamaica; Hispaniola; Puerto Rico; St. Croix; Dominica; Islas Los Roques and Caribbean coast of Colombia; Ceara to Pernambuco, Brazil; eastern Atlantic, from Senegal to Zaire (Powers, 1977).

***Euchirograpsus americanus* A. Milne Edwards, 1880**

Description: Williams, 1984:461, fig. 370.

Type-locality: Barbados, 126.2 m, Blake Stn. 278.

Distribution: Off Oregon Inlet, North Carolina, Florida through West Indies, and Colombia to Venezuela (Williams, 1984).

***Euchirograpsus antillensis* Turkay, 1975**

Description: Turkay, 1975:112, figs 4-5, 16a, 19, 25.

Type-locality: Cuba, Havana, Playa Baracoa, 23°04'30"N, 82°34'00"W, 414 m.

Distribution: Off Havana, Cuba; Arrowsmith

Banks, between Cuba and Yucatan; south of Florida Keys; Bahamas (Powers, 1977).

***Geograpsus lividus* (H. Milne Edwards, 1837)**

Description: Rathbun, 1918:232, pl. 55.--

Chace and Hobbs, 1969:157, figs. 48, 52a-c.

Type-locality: Antilles.

Distribution: Bermuda; Florida Keys, north and south coasts of Cuba; Jamaica; Puerto Rico; Virgin Islands to Barbados; Netherlands Antilles to Trinidad; Old Providence Island (Caribbean); Caribbean coast of Colombia to São Paulo, Brazil; eastern Atlantic, from Senegal to Angola; Cape Verde Islands; eastern Pacific, from southern part of Baja California to northern Chile; Clipperton Island; Galapagos Islands; Hawaiian Islands (Powers, 1977).

***Goniopsis cruentata* (Latreille, 1802)**

Description: Rathbun, 1918:237, fig. 136, pl. 57.--Chace and Hobbs, 1969:160, figs. 49, 52d-f.

Type-locality: Islands of South America.

Distribution: Bermuda; Bahamas; northwest Florida (rare); Tampico, Mexico; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; Virgin Islands to Barbados; Netherlands Antilles; Belize; Old Providence Islands (Caribbean); Surinam to Rio de Janeiro, Brazil; eastern Atlantic, from Senegal to northern Angola (Powers, 1977).

***Grapsus grapsus* (Linnaeus, 1758)**

Description: Rathbun, 1918:227, fig. 135, pls. 53, 54.--Chace and Hobbs 1969:163, figs. 50, 52 g-i.

Type-locality: America and Ascension Island.

Distribution: Bermuda; Bahamas; southeast and south Florida; Texas; north and south coasts of Cuba; Jamaica; Puerto Rico; Hispaniola; Virgin Islands to Barbados; Netherlands Antilles to Trinidad; Old Providence Island and Swan Island (Caribbean); Colombia to northern Brazil; eastern Atlantic, and from Portugal to Angola; Cape Verde Islands and Azores; St. Helena Island; Ascension Island; eastern Pacific from central Baja California to central Chile; Galapagos Islands; Clipperton Island (Powers, 1977).

***Pachygrapsus gracilis* (Saussure, 1858)**

Description: Rathbun, 1918:249, pl. 60: fig. 3, pl. 61: fig. 1.--Chace and Hobbs, 1969:167, figs. 51, 52j.

Type-locality: St. Thomas.

Distribution: Bermuda; Bahamas; south Florida; Texas; north and south coasts of Cuba; Jamaica; Puerto Rico; Virgin Islands; Caribbean coast of Columbia; Pernambuco to Bahia, Brazil; eastern Atlantic, from Senegal to Zaire (Powers, 1977)

Pachygrapsus transversus (Gibbes, 1850)

Description: Williams, 1984:459, fig. 368.

Type-locality: Key West, (Florida).

Distribution: Cape Lookout, North Carolina, to Montevideo, Uruguay; Bermuda; Mediterranean Sea to northern Angola; eastern Pacific from California to Peru, Galapagos Islands (Williams, 1984).

Percnon gibbesi (H. Milne Edwards, 1853)

Description: Williams, 1984:462, fig. 371.

Type-locality: Antilles.

Distribution: Fort Macon, North Carolina; southern Florida and Bahamas to Brazil; Bermuda; Azores to Angola; Cape San Lucas, Baja California, to Chile; Galapagos Islands (Williams, 1984).

Plagusia depressa (Fabricius, 1775)

Description: Williams, 1984:463, fig. 372.

Type-locality: "In mari mediterraneo" (erroneous).

Distribution: Beaufort, North Carolina, through Gulf of Mexico and West Indies to Pernambuco, Brazil; Bermuda; Azores; Madeira; Morocco to northern Angola; St. Helena Island (Chace 1966; Williams, 1984).

Planes minutus (Linnaeus, 1758)

Description: Chace, 1951:67, figs. 1a, 2a, d, g, j, k, l, 3a-h.

Type-locality: "Habitat in Palgi *Fuco natante*, supra aquam saepius cursitans".

Distribution: From off eastern North America (south of Newfoundland) through the eastern coast of America, Florida to Bahamas; West Indies.

Platychoirograpsus spectabilis De Man, 1896

Description: Monod, 1956:426, figs. 584-588.

Type-locality: Gabon.

Distribution: Gulf coast of Mexico; west coast of Florida (Powers, 1977); Gabon.

Sesarma benedicti Rathbun, 1897

Description: Rathbun, 1918:316, pl. 93.-- Abele, 1973:379, figs. 1A, 1G.

Type-locality: Surinam.

Distribution: Key West, Florida; Guyana and Surinam; Brazil (Powers, 1977).

Sesarma cinereum (Bosc, 1802)

Description: Williams, 1984:465, fig. 373.

Type-locality: "La Caroline."

Distribution: Magothy River, Chesapeake Bay, Maryland, to Palm Beach, east Florida; Collier County, west Florida, to Veracruz Mexico (Abele, 1973). Older records from the West Indies and elsewhere are erroneous (Williams, 1984).

Sesarma curacaoense De Man, 1892

Description: Rathbun, 1918:293, fig. 147, pl. 78: figs. 1, 2, pl. 160: fig. 3.--Abele, 1973:380, figs. 1C, 1F.

Type-locality: Curaçao.

Distribution: Key West, Florida; south and southwest Florida; north coast of Cuba; Jamaica; Puerto Rico; Curaçao; Bahia, Brazil (Powers, 1977).

Sesarma miersii Rathbun, 1897

Description: Abele, 1972a:166, figs. 1B, 1C, 2B, 2C; 1973:380, fig. 1I.

Type-locality: Bahamas.

Distribution: Bahamas; Key West, Florida; south coast of Cuba; Swan Island (Caribbean); Dominica (Powers, 1977).

Sesarma reticulatum (Say, 1817)

Description: Williams, 1984:466, fig. 374.

Type-locality: Muddy salt marshes (east coast of United States).

Distribution: Woods Hole, Massachusetts, to Volusia County, east Florida; Sarasota, west Florida, to Calhoun County, Texas (Abele, 1973).

Sesarma ricordi H. Milne Edwards, 1853

Description: Chace and Hobbs, 1969:183, fig. 62k.--Abele, 1973:378, fig. 1J.

Type-locality: Haiti.

Distribution: Bermuda; Bahamas; southeast Florida; Florida Keys; west coast of Florida; north coast of Yucatan; Cuba; Jamaica; Hispaniola; Puerto Rico; Virgin Islands to Trinidad; Curaçao; Old Providence Island (Caribbean); Yucatan to Surinam (Powers, 1977).

FAMILY PINNOTHERIDAE

Dissodactylus borradailei Rathbun, 1918
Description: Rathbun, 1918:121, fig. 68, pl. 27: figs. 5-8.

Type-locality: Miami, Florida; 55 m.
Distribution: Off southeast and southwest coasts of Florida; Jamaica (Powers, 1977).

Dissodactylus crinitichelis Moreira, 1901
Description: Williams, 1984:438, fig. 350.
Type-locality: Estado de Rio Grande do Sul, Brazil.
Distribution: Southeast of Cape Lookout, North Carolina off northwest Florida; Caribbean Sea and South America to Rio de la Plata, Argentina (Coelho and Ramos, 1972).

Dissodactylus mellitae (Rathbun, 1900)
Description: Williams, 1984:439, fig. 351.
Type-locality: Pensacola, Florida, on *Mellita quinquiesperforata*.
Distribution: Western part of Vineyard Sound, Massachusetts, to Charleston, South Carolina; Hutchinson Island, east Florida (Camp et al., 1977); western Florida; off Galveston, Texas (Rogers 1968; Williams, 1984).

Dissodactylus primitivus Bouvier, 1917
Description: Milne Edwards and Bouvier, 1923:346, fig. 8, pl. 8: figs. 3, 4, pl. 9: fig. 1.
Type-locality: West of Tortugas, Florida.
Distribution: Known only from the type-locality.

Dissodactylus rugatus Bouvier, 1917
Description: A. Milne Edwards and Bouvier, 1923:238, fig. 9, pl. 8: figs. 5, 6, pl. 9: figs. 2.
Type-locality: Dominique.
Distribution: East coast of Florida; Dominica.

Dissodactylus stebbingi Rathbun, 1918
Description: Rathbun, 1918:123, fig. 69, pl. 28: figs. 1, 2.
Type-locality: Sarasota Bay, Florida.
Distribution: Virginia; west and northwest coasts of Florida (Powers, 1977).

Fabia byssomiae (Say, 1818)
Description: Rathbun, 1918:105, fig. 56, pl. 24: figs. 6, 8.
Type-locality: Inhabits the *Byssomia distorta* (southern Atlantic coast of United States).
Distribution: West coast of Florida; northwest coast of Cuba (Powers, 1977).

Fabia tellinae Cobb, 1973

Description: Cobb, 1973:70, figs. 1-2.
Type-locality: Gulf of Mexico off NW Florida, 30°13'N 85°53'W, 12.2 m.
Distribution: Off northwest Florida to Alabama.

Orthotheres strombi (Rathbun, 1905)
Description: Rathbun, 1918:90, fig. 45, pl. 20: figs. 1, 2.
Type-locality: Clearwater Harbor, Florida.
Distribution: West and northwest coasts of Florida (Powers, 1977).

Parapinnixa bouvieri Rathbun, 1918
Description: Williams, 1984:447, fig. 357.
Type-locality: Off Cape Catoche, Yucatan (Mexico), 22°08'30"N, 86°53'30"W, 45.7 m, Albatross Stn. 2362.
Distribution: Off Charleston, South Carolina; south of Tortugas, Florida; Puerto Rico; and the type-locality (Williams, 1984).

Parapinnixa hendersoni Rathbun, 1918
Description: Williams, 1984:448, fig. 358.
Type-locality: Los Arroyos, Cuba.
Distribution: Southeast Cape Lookout, North Carolina, 34°29'N, 76°13'W, 33 m; 34°34'N, 75°50'W, 64 m; off Tampa Bay, Florida, through West Indies to Curaçao; Maranhao to Bahia, Brazil (Coelho and Ramos, 1972).

Pinnaxodes floridensis Wells and Wells, 1961
Description: Williams, 1984:449, fig. 359.
Type-locality: Outer beach near Fort Walton Beach, Florida.
Distribution: Off North Carolina to Georgia; northwest Florida (Williams, 1984).

Pinnixa chacei Wass, 1955
Description: Wass, 1955:160, figs. 5-9.
Type-locality: Gulf Beach, Alligator Point, Franklin County, Florida.
Distribution: Northwest Florida; Louisiana and Texas (Powers, 1977).

Pinnixa chaetoptera Stimpson, 1860
Description: Williams, 1984:451, fig. 360.
Type-locality: Charleston Harbor, South Carolina, on muddy or clayey shores in tubes of *Chaetopterus variopedatus*.
Distribution: Wellfleet, Massachusetts, to Rio Grande do Sul, Brazil (Williams, 1984).

***Pinnixa cristata* Rathbun, 1900**

Description: Williams, 1984:453, fig. 361.
Type-locality: Beaufort, North Carolina.
Distribution: Beaufort, North Carolina, to Edisto Island, South Carolina; Grande Isle, Louisiana, to Long Lake, Blackjack Peninsula, Aransas County, Texas (Hedgpeth, 1950; Williams, 1984).

***Pinnixa cylindrica* (Say, 1818)**

Description: Williams, 1984:453, fig. 362.
Type-locality: Jekyll Island, Georgia.
Distribution: North Falmouth, Massachusetts, to Pensacola, Florida (Cooley 1978), including Dry Tortugas (Williams, 1984).

***Pinnixa floridana* Rathbun, 1918**

Description: Williams, 1984:454, fig. 363.
Type-locality: Marco, Florida, also Sarasota Bay.
Distribution: Southeast off Cape Lookout, North Carolina; Hutchinson Island, east central Florida (Camp et al., 1977); west coast of Florida (Williams, 1984).

***Pinnixa leptosynaptae* Wass, 1968**

Description: Wass, 1968:137, figs. 1-6.
Type-locality: Bald Point at the entrance to Ochlockonee Bay, Franklin County, Florida.
Distribution: West coast of Florida (Powers, 1977).

***Pinnixa lunzi* Glassell, 1937**

Description: Glassell, 1937:3, figs. 1-8.-- Williams, 1984:455, figs. 364-365.
Type-locality: Isle of Palms (about 15 mi. NE of Charleston), South Carolina.
Distribution: Off Delmarva Peninsula, Virginia, North and South Carolina, Georgia; off Mississippi River delta and Seven and One-Half Fathom Reef off Texas near 26°51'N, 96°18'W (Williams, 1984).

***Pinnixa pearsei* Wass, 1955**

Description: Wass, 1955:164, figs. 10-13.
Type-locality: Indian Pass, Apalachicola, Florida.
Distribution: Northwest Florida (Powers, 1977).

***Pinnixa retinens* Rathbun, 1918**

Description: Williams, 1984:456, fig. 366.
Type-locality: Chesapeake Bay, off Poplar Island, Maryland, 36.6 m., soft bottom.
Distribution: Delaware Bay (Watling and

Maurer, 1976); Little River Inlet, South Carolina, Alligator Harbor, Florida; Aransas area of Texas coast (Williams, 1984).

***Pinnixa sayana* Stimpson, 1860**

Description: Williams, 1984:457, fig. 367.
Type-locality: Mouth of Beaufort Harbor, North Carolina, 10.97 m, sandy mud.
Distribution: Vineyard Sound, Massachusetts, to Beaufort, North Carolina; Hutchinson Island, east central Florida (Camp et al., 1977), Sarasota Bay, Florida, to Grand Isle, Louisiana; Amapá, Pará, Pernambuco, São Paulo, Brazil (Williams, 1984).

***Pinnotheres hemphilli* Rathbun, 1918**

Description: Rathbun, 1918:99, fig. 51, pl. 23.
Type-locality: Cedar Keys, Florida.
Distribution: Cedar Keys, Florida.

***Pinnotheres maculatus* Say, 1818**

Description: Williams, 1984:441, Fig. 353.
Type-locality: Given as "Inhabits the muricated *Pinna* of our coast."
Distribution: Off Martha's Vineyard, Massachusetts, to Golfo San Matias, Argentina (Fenucci, 1975).

***Pinnotheres moseri* Rathbun, 1918**

Description: Rathbun, 1918:94, text-fig. 47, pl. 21: figs. 3, 4, fig. 47.
Type-locality: Port Royal, Jamaica.
Distribution: West coast of Florida; Jamaica.

***Pinnotheres ostreum* Say, 1817**

Description: Williams, 1984:444, figs. 354-356.
Type-locality: "United States" (see Schmitt, et al., 1973); these authors mentioned six probable syntypes from the United States and Virginia in the British Museum (Natural History) and that, according to DeKay (1844), Say's specimen was from New Jersey.
Distribution: Salem, Massachusetts, to Santa Catarina, Brazil.

***Pinnotheres shoemakeri* Rathbun, 1918**

Description: Rathbun, 1918:95, fig. 48, pl. 22: figs. 1-4.
Type-locality: St. Thomas.
Distribution: West coast of Florida; St. Thomas, Virgin Islands.

FAMILY OCYPODIDAE

Ocypode quadrata (Fabricius, 1787)

Description: Williams, 1984:468, fig. 375.

Type-locality: Jamaica.

Distribution: Block Island, Rhode Island, to Santa Catarina, Brazil (megalopae have been taken at Woods Hole); Bermuda; Fernando de Noronha (Williams, 1984).

Uca burgersi Holthuis, 1967

Description: Crane, 1975:168, figs. 26F, 31H, 54G, 66F, 100, pl. 24E-H, map 12.

Type-locality: Plantage Knip, Westpunt, Curaçao, Netherlands Antilles.

Distribution: Bahamas; east coast of Florida; northeast (Gulf) coast of Yucatan; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; St. Thomas, Virgin Islands to Trinidad; Curaçao; east coast of Yucatan to Guatemala; Caribbean coast of Panama; Venezuela to Rio de Janeiro, Brazil (Powers, 1977).

Uca leptodactyla Rathbun, 1898

Description: Crane, 1975:304, figs. 37M, 56F, 60N-O, 69K-L, 101, map 17, pl. 41A-D.

Type-locality: Near Fort Montague, Nassau, New Province, Bahamas.

Distribution: West coast of Florida (not recently); east coast of Yucatan; north coast of Cuba; Jamaica; Puerto Rico; St. Croix; Curaçao; Venezuela to Santa Catarina, Brazil (Powers, 1977).

Uca longisignalis Salmon and Atsides, 1968

Description: Salmon and Atsides, 1968:279, figs. 1-4, 6, 7.

Type-locality: Ocean Springs, Mississippi.

Distribution: Northwest Florida to south Texas (Powers, 1977).

Uca minax (Le Conte, 1855)

Description: Williams, 1984:473, fig. 377a.

Type-locality: Beesleys Point, New Jersey.

Distribution: Buzzards Bay (Wareham and southwestern Cape Cod), Massachusetts, to northeast Florida, and from the area of Yankeetown, northwest Florida, to Louisiana, and on to Matagorda Bay, Texas (Williams, 1984).

Uca panacea Novak and Salmon, 1974

Description: Novak and Salmon, 1974:316, figs. 1-8.

Type-locality: Panacea, Florida.

Distribution: Northwest Florida to south Texas.

Uca pugilator (Bosc, 1802)

Description: Williams, 1984:475, figs. 376, 377c.

Type-locality: "Caroline."

Distribution: Cape Cod, Massachusetts; (rare on the north shore) southward around the tip of Peninsular Florida to near Pensacola (Heard, 1982); Old Providence Island, Bahamas, and Santo Domingo (Crane, 1975; Williams, 1984).

Uca pugnax (Smith, 1870)

Description: Williams, 1984:478, fig. 377b.

Type-locality: New Haven, (Connecticut).

Distribution: Provincetown, Massachusetts, to Daytona Beach, Florida (Williams, 1984).

Uca rapax (Smith, 1870)

Description: Crane, 1975:190, figs. 52c-D, 54F, 67C, 86, 91E-F, 100, pls. 27A-D, 45C-F, map 14.

Type-locality: Atlantic coast of Panama: Aspinwall.

Distribution: Bahamas; east coast of Florida; Florida Keys; southwest coast of Florida; northeast coast of Mexico to northeast Yucatan; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; St. Thomas, Virgin Islands to Trinidad and Tobago; Netherlands Antilles; east coast of Yucatan to Guatemala; Caribbean coast of Panama to Santa Catarina, Brazil (Powers, 1977).

Uca speciosa (Ives, 1891)

Description: Crane, 1975:236, figs. 68G, 101, map 15, pl. 31 A-D.

Type-locality: Port of Silam, Yucatan.

Distribution: Southeast Florida; Florida Keys; west and northwest coasts of Florida; northeast Yucatan and northwest Cuba (Powers, 1977).

Uca spinicarpa Rathbun, 1900

Description: Rathbun, 1918:411, pl. 148.-- Crane, 1975:239, figs. 68k, 101, pl. 31E-H, map 15.

Type-locality: Galveston.

Distribution: Alabama to northeastern coast of Mexico (Powers, 1977). Pensacola area, Florida (personal communication, P. M. Mikkelsen).

Uca thayeri Rathbun, 1900

Description: Crane, 1975:112, figs. 46K, 56E, 60H-I, 73A-B, 81I, 82I, 99, map 11, pl. 17.

Type-locality: Rio Parahyba do Norte at Cabedello, Brazil.

Distribution: East and southwest coasts of Florida; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; Guadeloupe; Trinidad; Tobago; Guatemala and Panama (Caribbean coasts) Venezuela to São Paulo, Brazil.

Uca vocator (Herbst, 1804)

Description: Crane, 1975:27, figs. 66D, 100, pl. 23E-G, pl. 24A-D, map 13.

Type-locality: "Amerika" (restricted by neotype selection of Holthuis, 1959, to Bank of Suriname River at Leosberg, Surinam).
Distribution: southern Florida; Tampico, Mexico; Belize to Guyana; Puerto Rico; Santo Domingo; Guadeloupe; Dominica; Trinidad and Tobago; Paraíba to Pernambuco, Brazil; ?Santa Catarina, Brazil.

Ucides cordatus (Linnaeus, 1763)

Description: Rathbun, 1918:347, fig. 158, pls. 110-113, pl. 159: figs. 3, 4.

Type-locality: America.

Distribution: Bahamas; southeast Florida; northeast Mexico to Panama; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; St. Thomas, Virgin Islands to Grenada; Colombia to Santa Catarina, Brazil.

FAMILY PALICIDAE***Palicus affinis*** A. Milne Edwards and Bouvier, 1899

Description: Rathbun, 1918:196, fig. 121, pl. 46, pl. 47: fig. 3.

Type-locality: Santa Cruz, 210 m.

Distribution: Southeast and west coasts of Florida; Dry Tortugas; Virgin Islands; Barbados; Guianas to Espirito Santo, Brazil (Powers, 1977).

Palicus alternatus Rathbun, 1897

Description: Williams, 1984:482, fig. 378.

Type-locality: 29°11'30"N, 85°29'00"W, 47.6 m. (south of Cape San Blas, Florida).

Distribution: Cape Hatteras to SE Cape Fear, North Carolina; Gulf of Mexico along west coast of Florida from Cape San Blas to Key West (Williams, 1984).

Palicus cristatipes (A. Milne Edwards, 1880)

Description: Rathbun, 1918:186, fig. 110.

Type-locality: Grenada, 166 m.

Distribution: Off Grenada, Windward Islands. Florida (personal communication, P. A. McLaughlin).

Palicus cursor (A. Milne Edwards, 1880)

Description: Rathbun, 1918:215, figs. 130-131, pl. 52: figs. 1, 2.

Type-locality: Sand Key, Havana, St. Kitts, Dominique, Barbados, 252-448 m.

Distribution: North Carolina; Florida Keys; west and northwest coasts of Florida; north coast of Cuba; St. Christopher; Dominica; Barbados (Powers, 1977).

Palicus dentatus A. Milne Edwards, 1880

Description: Rathbun, 1918:202, fig. 124.

Type-locality: Charlotte Harbor, 91 m, and Barbados, 110-176 m.

Distribution: Florida Keys; west coast of Florida; off Alabama; off Barbados (Powers, 1977).

Palicus faxoni Rathbun, 1897

Description: Williams, 1984:483, fig. 379.

Type-locality: Off Cape Hatteras, North Carolina, 89.6 m.

Distribution: Off Cape Hatteras, North Carolina, to near Cape Canaveral, Florida; off Yucatan, Mexico; near Quita Sueno Banks; southwest St. Christopher; off Cabo Frio, Rio de Janeiro (Williams, 1984).

Palicus floridana (Rathbun, 1918)

Description: Rathbun, 1918:220, pl. 41: figs. 3, 4.

Type-locality: Off Sand Key, Florida; 216 m.

Distribution: Known only from the type-locality.

Palicus gracilis (Smith, 1883)

Description: Rathbun, 1918:218, text-fig. 132, pl. 50, pl. 51: fig. 1.

Type-locality: Martha's Vineyard, Massachusetts, 260 m.

Distribution: Off Massachusetts; east coast of Florida; northwest Florida; Louisiana to central east coast of Mexico; north coast of Cuba; Curaçao (Powers, 1977).

Palicus obesus (A. Milne Edwards, 1880)

Description: Rathbun, 1918:205, fig. 125, pl. 49.

Type-locality: 23°13'N, 89°16'W, 154 m.

Distribution: Off northwest Florida and Mississippi; Campeche, Mexico (Powers, 1977).

Palicus sica (A. Milne Edwards, 1880)

Description: Williams, 1984:483, fig. 380.

Type-locality: Barbados, 150 m, *Blake Stn.* 293.

Distribution: Off Charleston, South Carolina, to northeast Cape Canaveral, Florida; west coast of Florida through West Indies to Barbados and Grenada (Williams, 1984).

FAMILY CRYPTOCHIRIDAE

Pseudocryptochirus corallicola (Verrill, 1908)

Description: Shaw and Hopkins, 1977:178, figs. 2b, 3b.

Type-locality: Dominica.

Distribution: Known only from Western

Atlantic: Dominica Island on *Mussa*; Bermuda Islands on *Mussa*, *Meandra* (= *Manicina*) and *Dichocognia*; Dry Tortugas, Florida on *Meandra* (= *Manicina*) *areolata* and *Meandrina*; Florida Middle Ground on *Scolymia lacera*, multiple polyp *Scolymia*, and *Manicina areolata* (Shaw and Hopkins, 1977).

Pseudocryptochirus hypostegus Shaw and Hopkins, 1977

Description: Shaw and Hopkins, 1977:179, figs. 1, 2a, 3a.

Type-locality: Florida Middle Ground about 137 km west of Tarpon Springs, Florida 28°30'49"N, 84°20'30"W, 27 m, from *Agaricia fragilis*.

Distribution: Known only from the eastern Gulf of Mexico on the Florida Middle Ground, in 25-30 meters, on *Agaricia fragilis* (Shaw and Hopkins, 1977).

ADDENDUM

The records cited below came to our attention or were published after this volume was completed.

FAMILY PALAEMONIDAE

Neopontonides chacei Heard, 1986

Description: Heard, 1986:472, figs. 1A, 2, 3, 4B-D.

Type-locality: Reef south of Marigot Bay, St. Lucia Island.

Distribution: Florida Keys south to Carrie Bow Cay, Belize, on the gorgonian

Pseudopterogorgia americana.

Periclimenaeus bredini Chace, 1972

Description: Chace, 1972:26, fig. 5.

Type-locality: Isla Mujeres off the Yucatan Peninsula.

Distribution: Known from the type-locality and the Florida Middle Grounds (Dardeau, 1984) where it was collected from a sponge.

Pontonia mexicana Guérin-Méneville, 1855

Description: Holthuis, 1951b:130, pl. 41.

Type-locality: Mexico (Holthuis, 1951b).

Distribution: Bahamas and Dry Tortugas; east coast of Mexico; West Indies (Chace, 1972).

Pseudopontonides principis (Crales, 1980)

Description: Crales, 1980:68.--Heard, 1986:481, Figs. 5A-F.

Type-locality: Awa di Oostpunt, Curacao.

Distribution: Northeastern Gulf of Mexico; Puerto Rico; Bonaire and Curacao. On antipatharians.

FAMILY ALPHEIDAE

Alpheus bahamensis Rankin, 1898

Description: Zimmer, 1913:405, figs. U¹-Z¹ (as *A. hippothoe* var. *edamensis*?).--Chace, 1972:58.

Type-locality: New Providence Island, Bahamas.

Distribution: Bermudas; Dry Tortugas; Yucatan Peninsula; West Indies (Chace, 1972).

Fenneralpheus chacei Felder and Manning, 1986

Description: Felder and Manning, 1986:498, figs. 1-3.

Type-locality: Fort Pierce Inlet, St. Lucie

Country, Florida.

Distribution: Fort Pierce and Key West, Florida (Felder and Manning, 1986).

Salmonaeus cavicolus Felder and Manning, 1986

Description: Felder and Manning, 1986:503, figs. 4-6.

Type-locality: South side of Fort Pierce Inlet, St. Lucie County, Florida.

Distribution: Known only from the type-locality.

Synalpheus scaphoceris Coutière, 1910

Description: Dardeau, 1986:491, figs. 1-3.

Type-locality: Dry Tortugas, Florida.

Distribution: Gulf of Mexico: Isla de Lobos, West Flower Garden Bank, East Flower Garden Bank, Florida Middle Grounds, off Sanibel Island, Dry Tortugas; Caribbean: Peurto Rico and Curacao; Brazil (Dardeau, 1986).

FAMILY GONEPLACIDAE

Chasmocarcinus chacei Felder and Rabalais, 1986

Description: Felder and Rabalais, 1986:548, figs. 1, 2a-g, 3a-h.

Type-locality: Vicinity of Flower Garden Banks, northwestern Gulf of Mexico (27°53.97'N, 93°34.79'W, 126 m).

Distribution: Gulf of Mexico from the Texas coast to Dry Tortugas, Florida; possibly off Anguilla (Felder and Rabalais, 1986).

Speocarcinus carolinensis Stimpson, 1859

Description: Williams, 1984:437, fig. 349.--Felder and Rabalais, 1986:572, figs. 11d-f, 12.

Type-locality: Charleston Harbor, South Carolina.

Distribution: South of Cape Hatteras, North Carolina, through the West Indies to Amapá, Brazil (Williams, 1984). Dry Tortugas and the Margueas Keys, Florida (Felder and Rabalais, 1986).

Key to families of Florida decapods

[Based on Rathbun, 1918, 1930, 1937, Barnard, 1950,
Holthuis, 1955, and Williams, 1984]

1. General form shrimplike, usually compressed; pleuron of second somite never overlapping that of first somite; first 3 pairs of pereopods usually chelate (except in some Sergestoidea), third pair never unusually robust 2

General form shrimplike, lobsterlike, or crablike; if shrimplike with pleuron of second abdominal somite overlapping that of first somite and third pair of pereopods not chelate or unusually enlarged (except Stenopodidae, with pleuron of second abdominal somite not overlapping that of first somite; third pereopods chelate, stronger than preceding) 8
2. (1) Fourth and fifth pereopods well developed..... 3

Fourth and fifth pereopods reduced or absent..... 7
3. (2) Postorbital spine present..... **Solenoceridae** (page 97)

Postorbital spine absent..... 4
4. (3) Integument rigid, of stony appearance; cervical groove very faint or absent.....

..... **Sicyoniidae** (page 109)

Integument more or less flexible, not stony and rigid in appearance; cervical groove present and easily discerned 5
5. (4) Eyestalks without tubercles on their mesial (inner) borders; epipods absent behind third pereopods **Penaeidae** (page 82)

Eyestalks each with tubercle on its mesial (inner) border; epipods on all coxae from second maxillipeds through fourth pereopods 6
6. (5) Distal, filamentous portion of upper antennular flagellum extensively developed.....

..... **Benthescymidae** (page 79)

Distal, filamentous portion of upper antennular flagellum not extensively developed. **Aristeidae** (page 79)
7. (2) Anterior region of cephalothorax not greatly elongate; gills present.....

..... **Sergestidae** (page 115)

Anterior region of cephalothorax greatly elongate; gills absent.....

..... **Luciferidae** (page 125)
8. (1) Form shrimplike; usually with body compressed..... 9

Form lobsterlike or crablike..... 24

9. (8) Pleuron of second abdominal somite not overlapping that of first somite; third pereopods chelate, stronger than preceding **Stenopodidae** (page 281)
- Pleuron of second abdominal somite overlapping that of first somite; third pair of pereopods never chelate 10
10. (9) First pair of pereopods chelate or simple..... 11
- First pair of pereopods subchelate..... 23
11. (10) Fingers of all four chelae slender, their cutting edges pectinate.....
- **Pasiphaeidae** (page 137)
- Cutting edges of fingers of chelae not all pectinate..... 12
12. (11) Carpi of second pair of pereopods entire; first pair of pereopods always with well-developed chelae 13
- Carpi of second pair of pereopods usually subdivided into two or more segments; if not, first pair of pereopods not chelate 19
13. (12) First pair of pereopods stronger and heavier though often shorter than second..... 14
- First pair of pereopods usually more slender than, rarely subequal to, second..... 16
14. (13) Ends of fingers of first two pairs of pereopods not dark colored; ultimate segment of second maxilliped placed at end of penultimate segment; exopod of first maxilliped without flagellum **Bresiliidae** (page 141)
- Ends of fingers of first two pairs of pereopods dark colored; ultimate segment of second maxilliped applied as strip alongside of penultimate segment; exopod of first maxilliped with distinct flagellum 15
15. (14) Rostrum immovable; exopods on pereopods..... **Eugonatonotidae** (page 145)
- Rostrum movable; no exopods on pereopods..... **Rhynchocinetidae** (page 145)
16. (13) Pereopods usually with exopods; if not, fingers of chelae with terminal brushes of long hairs 17
- Pereopods without exopods; chelae without terminal brushes of long hairs..... 18
17. (16) Mandible without palp; fingers of chelae usually with conspicuous terminal brushes of hairs; last three pairs of pereopods not conspicuously lengthened; pereopods with or without exopods **Atyidae** (page 127)
- Mandible with palp; fingers of chelae without terminal brushes of hairs; pereopod with exopods (last three pairs of pereopods not conspicuously lengthened; carpi of these pereopods distinctly shorter than propodi) **Oplophoridae** (page 131)

18. (16) Mandible usually with incisor process; if not, third maxilliped not expanded or leaf-like **Palaemonidae** (page 152)
- Mandible without incisor process; third maxillipeds expanded and leaf-like.....
..... **Gnathophyllidae** (page 149)
19. (12) Chelae of first pair of pereopods microscopically small or absent (mandible bifid, with palp; rostrum laterally compressed, distinctly dentate)
..... **Pandalidae** (page 262)
- Chelae of first pair of pereopods distinct, at least on one side..... 20
20. (19) First pair of pereopods both chelate; rostrum dentate or unarmed, not with single subdistal dorsal tooth 21
- Usually right first pereopod chelate, the other ending in simple claw-like dactyl; if both chelate, rostrum with subdistal dorsal tooth **Processidae** (page 254)
21. (20) Ends of fingers of first pair of chelae usually dark colored; first pair of chelipeds short and rather heavy but not swollen; eyes free, never extremely elongate
..... **Hippolytidae** (page 230)
- Ends of fingers of first pair of chelae not dark colored; eyes either extremely long or partly or wholly covered by carapace 22
22. (21) Eyes extremely elongate, reaching almost to end of antennular peduncle; cornea small; first pair of pereopods shorter than and about as robust as second
..... **Ogyrididae** (page 251)
- Eyes usually partly or wholly covered by carapace, never very elongate; first pair of pereopods distinctly stronger than second, often unequal and swollen
..... **Alpheidae** (page 194)
23. (10) Carpi of second pair of pereopods multi-articulate.....
..... **Glyphocrangonidae** (page 277)
- Carpi of second pair of pereopods not subdivided..... **Crangonidae** (page 271)
24. (8) Body lobsterlike and strongly calcified; abdomen with pleura well developed; first three pairs of pereopods either all chelate or none chelate 25
- Body crablike or lobsterlike, sometimes weakly calcified in part; pleura often reduced or absent; first three pairs of pereopods never alike; first, second, or first and second pereopods chelate or subchelate 28
25. (24) First three pairs of pereopods chelate, first largest; uropods well developed, lateral ramus (uropodal exopod) transversely divided .. **Nephropidae** (page 285)
- First three pairs of pereopods never chelate; uropods well developed, lateral ramus without transverse division 26

26. (25) Carapace with small rostrum; first pereopods much larger than others (body tubular; antennae cylindrical, shorter than body) **Synaxiidae** (page 323)
- Carapace without rostrum; first pereopods not enlarged except in *Justitia* 27
27. (26) Carapace subcylindrical; antennal flagella long, strong, and spiny..... **Palinuridae** (page 313)
- Carapace more or less flattened dorsoventrally, lateral margins sharp; antennae short, flagella replaced by plates with dentate or lobulate margins **Scyllaridae** (page 316)
28. (24) Either lobsterlike or crablike; abdomen extended, bent upon itself, or flexed beneath thorax; last thoracic sternite free; uropods present; carapace not fused with epistome; first, second, or first 2 pairs of pereopods chelate or subchelate 29
- Crablike; abdomen permanently flexed beneath carapace; last thoracic sternite fused with preceding; uropods rarely present, never biramous; carapace fused with epistome; first pair of pereopods chelate or subchelate 40
29. (28) Second to fourth pereopods with dactyli conspicuously curved and flattened; abdomen much reduced in size and flexed beneath thorax 30
- Second to fourth pereopods with dactyli not conspicuously curved and flattened; abdomen well developed but may be flexed beneath thorax 31
30. (29) First pair of pereopods subchelate; carapace depressed... **Albuneidae** (page 427)
- First pair of pereopods simple; carapace subcylindrical..... **Hippidae** (page 433)
31. (29) Abdomen usually asymmetrical (rarely secondarily straightened), usually membranous and with uropods adapted for holding body in hollow objects; rarely leathery, unprotected, and bent under thorax 32
- Abdomen symmetrical and obviously segmented; uropods well developed for swimming, never for holding body in hollow objects 35
32. (31) Third maxillipeds approximated at base; chelipeds subequal, or left much larger than right, rarely with right slightly larger than left 33
- Third maxillipeds widely separated at base by sternum; right cheliped usually much larger than left, left never larger than right, occasionally subequal 34
33. (32) Ventral antennular flagellum ending in filament..... **Diogenidae** (page 330)
- Ventral antennular flagellum ending bluntly..... **Coenobitidae** (page 327)

34. (32) Carapace firm anteriorly, more or less membranous posteriorly; rostrum obsolete or nearly so; fourth pereopods unlike third **Paguridae** (page 359)
- Carapace firm all over, spiny in many species; rostrum more or less spiniform; fourth pereopods like third **Lithodidae** (page 355)
35. (31) Body subcylindrical; first two pairs of pereopods chelate or subchelate (first only in Upogebiidae); abdomen extended 36
- Body slightly depressed; only first pereopods chelate; abdomen bent under thorax. 38
36. (35) No *linea thalassinica*; both movable and fixed antennal thorns present; first pereopods strongly chelate and conspicuously hairy **Axiidae** (page 289)
- Linea thalassinica* present; fixed antennal thorn absent; first pereopods chelate or subchelate but not conspicuously hairy 37
37. (36) First pereopods chelate; rostrum inconspicuous or absent. **Callianassidae** (page 293)
- First pereopods subchelate; rostrum well developed, dorsally flattened, spiny, and hairy **Upogebiidae** (page 309)
38. (35) Form somewhat lobsterlike; rostrum extended, well developed; abdomen loosely flexed beneath posterior thorax; third maxilliped pediform 39
- Form crablike (*Euceramus* elongate) with abdomen completely folded under thorax; rostrum short and broad or wanting; third maxilliped flattened, operculiform **Porcellanidae** (page 410)
39. (38) Antennal peduncle composed of four movable segments; telson subdivided into two or more plates, not folded sharply against itself, without lateral indentation ... **Galatheidae** (page 397)
- Antennal peduncle with five segments, third segment not being fused with second; telson never subdivided into two or more plates, folded sharply against itself, with lateral indentation **Chirostylidae** (page 393)
40. (28) Mouth-frame (buccal cavity) triangular..... 41
- Mouth-frame (buccal cavity) more or less quadrate..... 46

41. (40) Posterior thoracic sternites narrow, keel-like (bases of 2nd-4th pereopods close together); last pair of pereopods dorsal in position; female genital openings coxal (body elongate in dorsal view, subcylindrical; pereopods adapted for burrowing; orbits hidden ventrolaterally if present; *linea homolica* absent) **Raninidae** (page 455)
- Posterior thoracic sternites broad (bases of walking legs far apart); last pair of pereopods normal in position, or last two pairs dorsal; female genital openings sternal (except in *Cyclodorippidae*) 42
42. (41) Carapace subquadrilateral or subcircular, short, leaving the first 2 or 3 abdominal segments exposed; last two pairs of pereopods dorsal in position, ending in hook-like movable fingers 43
- Carapace of usual crablike shape 45
43. (42) Third maxilliped leaving all anterior part of buccal cavity uncovered **Dorippidae** (page 461)
- Third maxilliped greatly elongate and not leaving any appreciable portion of buccal cavity uncovered 44
44. (43) Third maxilliped with flagellum **Cymonomidae** (page 443)
- Third maxilliped without flagellum **Cyclodorippidae** (page 447)
45. (42) Afferent opening to each gill chamber in front of base of cheliped **Calappidae** (page 465)
- Afferent opening to each gill chamber at base of outer (third) maxilliped **Leucosiidae** (page 479)
46. (40) Last pair of pereopods abnormal, dorsal; female openings coxal; first abdominal limbs of female present; gills usually many 47
- Last pair of pereopods normal, rarely reduced, not dorsal, except in *Palicus* (*Palicidae*) and *Retropluma*; female openings sternal; first abdominal limbs of female wanting; gills few 50
47. (46) Sternum of female with longitudinal grooves; vestiges of sixth abdominal limbs usually present; eyes usually completely sheltered by orbits when retracted 48
- Sternum of female without longitudinal grooves; no vestiges of sixth abdominal limbs; eyes incompletely or not at all sheltered by orbits when withdrawn against body 49

48. (47) Vestiges of sixth abdominal limbs present (except in *Hypoconcha*, where also no mastigobranchs are present); carapace usually not longer than broad, with well-marked side edge [mastigobranchs on first pereopods (chelipeds) only or none] ...
..... **Dromiidae** (page 437)
- No vestige of sixth abdominal limbs; carapace longer than broad, with ill-marked side edge; first three pereopods with mastigobranchs, fourth and fifth small, subdorsal, and prehensile **Homolodromiidae** (page 443)
49. (47) Body rectangular; basal article of eyestalk not much longer than terminal article....
..... **Homolidae** (page 451)
- Body pyriform; basal article of eyestalk much longer than terminal article.....
..... **Latreilliidae** (page 451)
50. (46) Forepart of body narrow, usually forming distinct rostrum; body more or less triangular; orbits generally incomplete 51
- Forepart of body broad; rostrum usually reduced or wanting; body oval, round, or square; orbits nearly always well enclosed 52
51. (50) Chelipeds not much larger than other pereopods; hooked hairs almost always present; second segment of antenna well developed, usually fused with epistome and front **Majidae** (page 493)
- Chelipeds very much larger than other pereopods; hooked hairs almost always absent; second segment of antenna small, short, and not fused with epistome or front **Parthenopidae** (page 558)
52. (50) Merus of third maxilliped small, bearing terminally carpus of nearly its own width; ischium very broad; body somewhat oblong; antennule not retractile into sockets; parasitic on corals **Cryptochiridae** (page 727)
- Carpus of third maxilliped articulate at or near antero-lateral angle of merus; body usually rounded or transversely oval; male openings nearly always coxal; right chela larger than left in many species 53
- Carpus of third maxilliped not articulating at or near inner angle of merus; body usually square or squarish; male openings sternal except in *Retropluma*, where duct passes along a sternal groove to coxopodite; right chela almost never larger than left (except Palicidae) 57
53. (52) Pereopods more or less distinctly adapted for swimming; usually a small lobe on inner angle of endopod in first maxillipeds; first antenna fold slanting or transverse **Portunidae** (page 572)
- Pereopods not adapted for swimming, or if so modified, then male genital duct opening sternally or running in sternal groove; inner lobe on endopod in first maxillipeds wanting 54

54. (53) Antennule folds lengthwise..... 55
 Antennule folds slanting or transversely..... 56

55. (54) Carapace subcircular; antennal flagella either long and hairy or wanting.....
 **Atelecyclidae** (page 569)
 Carapace broadly oval or hexagonal; antennal flagella present, short, not hairy.....
 **Cancridae** (page 569)

[The following three families are not sharply separated.]

56. (54) Carapace usually transversely oval or transversely hexagonal (xanthoid); male openings coxal; male abdomen greatly narrowed in segments 4-7; tending to occur in shallow water **Xanthidae** (page 603)

Carapace subquadrate to xanthoid; male openings coxal with genital duct lying in groove between sternites 7 and 8 or sternal; male abdomen somewhat more triangular than above; part of sternite 8 visible from above at level of second abdominal segment but variable in size (some species in above family share this character); tending to occur in deeper water near edge of continental shelf
 **Goneplacidae** (page 591)

Carapace hexagonal to trapezoidal in shape, with anterolateral margins generally armed with three to five teeth, with front bearing four short teeth; orbits and eyes well developed; antennules transverse or transversely oblique; basal antennal article movable and not reaching front of carapace; genital openings in male coxal; pereopods long and compressed **Geryonidae** (page 569)

57. (52) Small, usually commensal crabs, with very small eyes and orbits; body usually more or less rounded **Pinnotheridae** (page 681)

Free-living crabs, with eyes not especially reduced and usually square body.... 58

58. (57) Last pair of pereopods dorsally placed and weaker than others; interantennular septum very thin; no distinct epistome; exopod of third maxilliped not hidden
 **Palicidae** (page 718)

Last pair of pereopods not dorsally placed or markedly weaker than others; interantennular septum not very thin 59

59. (58) Gap of greater or less size between third maxillipeds; front very or moderately broad 60

Third maxillipeds almost or quite close to mouth; front moderately or very narrow.
 **Ocypodidae** (page 707)

60. (59) Sides of body either straight or very slightly arched; shape squarish; front broad..
 **Grapsidae** (page 665)

Sides of body strongly arched; shape transversely oval; front narrow.....
 **Gecarcinidae**(page 661)

Keys to Species of Florida Decapods

Suborder Dendrobranchiata

Family Aristeidae

Key to genera and species
[Based on Roberts and Pequegnat, 1970]

Hepatic spine absent (epipod on fourth pereopod large; podobranch on third pereopod large; rostrum tridentate) *Plesiopenaeus edwardsianus*

Hepatic spine present (podobranch on third pereopod and epipod on fourth pereopod well developed) *Aristaeomorpha foliacea*

Family Benthescymidae

Genus *Betheogennema* Burkenroad, 1936
[based on Roberts and Pequegnat, 1970]

Podobranchs present on first maxilliped through third pereopod; telson with more than single pair of movable lateral spinules but without posteriomedian point *B. intermedia*

Aristaeomorpha foliacea

a. adult male lateral view

(after Pérez Farfante, 1978)

Plesiopenaeus edwardsianus

b. lateral view

c. male petasma

(after Crosnier and Forest, 1973)




Bentheogennema intermedia

female:

d. anterior region, lateral view

e. telson, dorsal view

(after Crosnier and Forest, 1973)



Family Penaeidae

Key to genera and species
[Adapted from Pérez Farfante, 1978]

1. Rostrum toothed on dorsal margin, usually also on ventral margin; pleurobranch present on last thoracic somite 2
 Rostrum toothed on dorsal margin only; no pleurobranch on last thoracic somite... 3
2. (1) Carapace hairy.....*Funchalia villosa*
 Carapace smooth.....*Penaeus*
3. (1) Telson tridentate, with fixed spine on each side of tip; mesial border of first segment of antennular peduncle bearing spine (parapenaeid spine) 4
 Telson usually without fixed spines; no spine on mesial border of first segment of antennular peduncle 6
4. (3) Carapace with longitudinal and transverse sutures.....*Parapenaeus*
 Carapace without longitudinal or transverse sutures..... 5
5. (4) Male with symmetrical petasma; single arthrobranch on last thoracic somite, no trace of second arthrobranch *Penaeopsis serrata*
 Male with asymmetrical petasma; 2 arthrobranchs present on last thoracic somite, one of them well developed, other vestigial *Metapenaeopsis*
6. (3) Dactyli of fourth and fifth pairs of pereopods elongate and subdivided.....
 *Xiphopenaeus kroyeri*
 Dactyli of fourth and fifth pairs of pereopods of normal shape and undivided..... 7
7. (6) Carapace without longitudinal sutures.....*Trachypeneopsis mobilispinis*
 Carapace with longitudinal sutures (upper antennular flagella shorter than carapace and not much longer than lower flagella; fourth and fifth pairs of pereopods about as heavy as 3 anterior pairs; exopod of fifth pair of pereopods well developed)
 *Trachypenaeus*

Genus *Metapenaeopsis* Bouvier, 1905

Key to species

[Adapted from Pérez Farfante, 1971]

1. Thelycum with median plate bearing horseshoe-shaped marginal strip and coiled lateral strips; petasma with distoventral projection cleft by deep sinus into 2 long, subequal lobes *M. smithi*
- Thelycum with median plate lacking marginal and coiled strips; petasma with distoventral projection simple, forming one single lobe or cleft by shallow sinus into 2 short, subequal, or into 2 unequal lobes 2
2. (1) Thelycum with anterior part of median plate convex, bearing 2 large pits; petasma with distoventral projection mittenlike in outline, large left lobe extending distally far beyond small right lobule *M. gerardoi*
- Thelycum with anterior part of median plate long, half or more as long as median plate; petasma with distoventral projection cleft into 2 unequal lobes, right lobe noticeably larger than left *M. goodei*

Genus *Parapenaeus* Smith, 1886

Key to species

[Adapted from Roberts and Pequegnat, 1970]

- Branchiostegal spine present behind anterior margin of carapace; rostral teeth usually seven; epigastric tooth and hepatic spine not as far behind orbital margin as in *P. americanus* *P. politus*
- Branchiostegal spine on anterior margin of carapace; rostral teeth usually six; epigastric tooth and hepatic spine farther behind orbital margin than in *P. politus*
..... *P. americanus*

Genus *Penaeus* Fabricius, 1798

Key to species of adults and subadults
[Adapted from Williams, 1984]

1. Lateral rostral grooves reaching only slightly beyond posterior rostral tooth (non-grooved shrimps) *P. setiferus*
- Lateral rostral grooves reaching nearly to posterior margin of carapace (grooved shrimps) 2
2. (1) Petasma with distomesial projection long; distal fold expanded mesially forming large spined lobe; ventral costa with apex free; thelycum with anteromesial corners of lateral plates extended to cover posterior process of median protuberance *P. brasiliensis*
- Petasma with distomesial projection relatively short; distal fold not forming lobe; ventral costa with apex attached to adjacent membraneous part; thelycum with anteromesial corners not extended, exposing posterior process of median protuberance 3
3. (2) Petasma armed with minute spines on ventral costa along terminal part of free border; thelycum with anteromesial corners of lateral plates slightly divergent, posterior process of median protuberance with undivided median carina *P. duorarum*
- Petasma unarmed on ventral costa along terminal part of free border; thelycum with anteromesial corners of lateral plates widely divergent, posterior process of median carina bifurcate anteriorly *P. aztecus*

Key to species of juveniles between 17 and 47 mm total length
[Adapted from Williams, 1984]

1. Lateral rostral grooves reaching only slightly beyond posterior rostral tooth; rostrum long and slightly upturned at tip in individuals exceeding 22 mm total length; ground color light gray, sometimes with greenish cast in shrimp taken from beds of vegetation; chromatophores (widely spaced except on spines, ridges, and uropods) colored slate-blue and brown; uropods with reddish-brown to brown areas distally *P. setiferus*
- Lateral rostral grooves reaching almost to posterior margin of carapace (shallow in 17-mm individuals); rostrum relatively short; color gray to light brown, sometimes with greenish cast in shrimp taken from beds of vegetation; chromatophores numerous and closely spaced, often in bands or patches 2
2. (1) Rostrum with toothed dorsal margin straight; tip attenuate and straight *P. brasiliensis*
- Rostrum with toothed dorsal margin slightly arched over eye; tip short or attenuate and slightly upturned 3

3. (2) Rostrum usually not upturned at tip and not extremely attenuate; chromatophore: slate-blue and brown; usually with conspicuously pigmented lateral spot at juncture of third and fourth abdominal somites; uropods with uniform sprinkling of chromatophores, degree of transparency uniform throughout (color more dense in older individuals) *P. duorarum*

Rostrum usually slightly upturned and attenuate at tip; chromatophores brown and olive-green; usually lacking lateral spot at juncture of third and fourth abdominal somites; uropods with reddish-brown to brown areas distally *P. aztecus*

Genus *Trachypenaeus* Alcock, 1901

Key to species
[Adapted from Chace, 1972]

Thelycum pubescent, lips of transverse groove strongly biconvex; male with sternal elevation between coxae of fifth pereopods goblet-shaped, constricted posteriorly ...
..... *T. constrictus*

Thelycum naked, lips of transverse groove subhorizontal; male with sternal elevation between coxae of fifth pereopods triangular, sloping regularly to posterior apex *T. similis*

Metapenaeopsis smithi

a. petasma, ventral view (male)

b. thelycum (female)

(after Pérez Farfante, 1971)

Metapenaeopsis gerardoi

c. petasma, ventral view (allotype male)

d. thelycum (holotype female)

(after Pérez Farfante, 1971)





Metapenaeopsis goodei

e. lateral view (female)

f. petasma, ventral view (male)

g. thelycum (female)

(after Pérez Farfante, 1971)



e

f



g

Parapenaeus politus

- a. anterior region, lateral view
(after Williams, 1965a)

Parapenaeus americanus

- b. lateral view (female)
(after Rathbun, 1901)



Penaeus setiferus

- a. lateral view (female)
 - b. petasma (male)
 - c. thelycum (female)
- (after Pérez Farfante, 1978)

Penaeus brasiliensis


- d. lateral view (male)
 - e. thelycum (female)
 - f. petasma (male)
- (after Pérez Farfante, 1978)

Penaeus duorarum

- g. anterior region, lateral view (female)
 - h. thelycum (female)
 - i. petasma (male)
- (after Pérez Farfante, 1978)

Penaeus aztecus

- j. lateral view (female)
 - k. petasma (male)
 - l. thelycum (female)
- (after Pérez Farfante, 1978)




Trachypenaeus constrictus

- a. lateral view (female)
 - b. thoracic sternites between fourth pereopods and fifth pereopods
 - c. petasma (male)
- (after Pérez Farfante, 1978)

Trachypenaeus similis

- d. thelycum (female)
 - e. thoracic sternites between fourth pereopods and fifth pereopods (male)
- (after Burkenroad, 1934)



Funchalia villosa

- a. lateral view
 - b. thelycum (female)
 - c. petasma (male)
- (after Burukovskii, 1983)

Penaeopsis serrata





- d. anterior region, lateral view (female)
 - e. petasma, ventral view (male)
- (after Pérez Farfante, 1980b)

Trachypeneopsis mobilispinis

- f. rostrum
 - g. telson
 - h. petasma (male)
- (after Rathbun, 1920)

Xiphopenaeus kroyeri

- i. lateral view (female)
 - j. petasma, posterior view (male)
- (after Chace and Hobbs, 1969)



Family Solenoceridae

Key to genera and species
[Adapted from Pérez Farfante, 1977]

1. Upper and lower antennular flagella lamellate; exopod of uropod lacking distolateral spine *Solenocera*
- Upper antennular flagellum subcylindrical, lower subcylindrical or flattened; exopod of uropod armed with distolateral spine 2
2. (1) Lower antennular flagellum conspicuously depressed, orbital spine present *Mesopenaeus tropicalis*
- Lower antennular flagellum subcylindrical, occasionally depressed; if so, orbital spine lacking 3
3. (2) Epigastric and first rostral teeth separated from remaining teeth by long interval; suprahepatic spine absent *Hymenopenaeus*
- Epigastric tooth separated from first rostral tooth by interval not conspicuously greater or smaller than that between first and second rostral teeth 4
4. (3) Rostrum low, with ventral margin straight or concave; submarginal carina present...
..... *Pleoticus robustus*
- Rostrum deep, with ventral margin pronouncedly convex; submarginal carina absent *Hadropenaeus*

Genus *Hadropenaeus* Pérez Farfante, 1977

Key to species

[Adapted from Pérez Farfante, 1977]

Scaphocerite reaching distal end of antennular peduncle or overreaching it by not more than 0.1 of its own length; prosartema extending only to distomesial extremity of first antennular segment; thelycum with median protuberance on sternite between fifth pereopods projecting ventrally, and tooth of median keel of sternite between fourth pereopods directed anteriorly; petasma with distomesial projection of ventromedian lobule directed mesially *H. affinis*

Scaphocerite overreaching antennular peduncle by about 0.25 of its own length; prosartema conspicuously overreaching distomesial margin of first antennular segment; thelycum with median protuberance on sternite between fifth pereopods projecting anteriorly, and tooth of median keel of sternite between fourth pereopods directed ventrally or posteriorly; petasma with distomesial projection of ventromedian lobule directed distally *H. modestus*

Genus *Hymenopenaeus* Smith, 1882

Key to species

[Adapted from Pérez Farfante, 1977]

Eye with cornea hemispherical and disposed such that imaginary line extending from mesial tubercle parallel to basal margin of ocular peduncle intersects lateral border of latter far proximal to proximolateral extremity of cornea ... *H. aphoticus*

Eye with cornea subreniform and disposed such that imaginary line extending from mesial tubercle parallel to basal margin of ocular peduncle intersects posterolateral extremity of cornea *H. debilis*

Genus *Solenocera* Lucas, 1849

Key to species
[Adapted from Williams, 1984]

1. Rostral + epigastric teeth 8-11; postrostral carina high and sharp, extending almost to posterior margin of carapace *S. vioscai*

 Rostral + epigastric teeth 4-8; postrostral carina low or absent posterior to cervical sulcus 2
2. (1) Anterior part of carapace glossy; scaphocerite exceeding distal end of antennular peduncle by 10% of its own length *S. necopina*


 Anterior part of carapace setose; scaphocerite never exceeding distal end of antennular peduncle by 10% of its own length, usually less *S. atlantidis*

Hadropenaeus affinis

- a. lateral view (female)
 - b. thelycum, ventral view (female)
 - c. petasma, dorsal view of right half (male)
- (after Pérez Farfante, 1977)

Hadropenaeus modestus

- d. anterior region, lateral view
 - e. thelycum, ventral view (holotype female)
 - f. petasma, dorsal view of right half (male)
- (after Pérez Farfante, 1977)



Hymenopenaeus aphoticus

female:

a. anterior region, lateral view

b. eye

(after Pérez Farfante, 1977)


Hymenopenaeus debilis

male:

c. lateral view

d. eye

(after Pérez Farfante, 1977)



Solenocera vioscai




- a. carapace and rostrum, lateral view (larger paratype female)
(after Burkenroad, 1934)

Solenocera necopina

- b. anterior region, lateral view (female)
(after Pérez Farfante and Bullis, 1973)

Solenocera atlantidis

- c. anterior region, lateral view (female)
(after Pérez Farfante and Bullis, 1973)



Mesopenaeus tropicalis

a. lateral view (female)

(after Pérez Farfante, 1977)



Pleoticus robustus

male:

b. lateral view

c. petasma, dorsolateral view of left half

(after Pérez Farfante, 1977)



Family Sicyoniidae

Genus *Sicyonia* H. Milne Edwards, 1830

Key to species
[Adapted from Williams, 1984]

1. First pereopod with basis and ischium armed with spine; abdominal somite 2 with dorsal carina notched at junction of transverse sulci 2
 First pereopod with basis and ischium unarmed; abdominal somite 2 with dorsal carina unnotched 3
2. (1) Rostrum (excluding tip) with 2 dorsal teeth anterior to posterior orbital margin; carina of carapace with 3 teeth, first tooth smallest *S. laevigata*
 Rostrum (excluding tip) with 3 dorsal teeth; carina of carapace with 3 evenly spaced, subequal teeth *S. parri*
3. (1) Carapace with 3 large dorsal teeth behind hepatic spine..... *S. brevirostris*
 Carapace with 1 or 2 large dorsal teeth behind hepatic spine..... 4
4. (3) Two teeth on dorsal carina behind hepatic spine.....*S. typica*
 One tooth on dorsal carina behind hepatic spine..... 5
5. (4) Pleura of abdominal somite 4 with both antero- and posteroventral margins spined or angular
 *S. dorsalis*
 Pleura of abdominal somite 4 with posteroventral margin rounded..... 6
6. (5) Antennal spine long, acute, buttressed; pleura of abdominal somites 1-4 with ventral spines laterally recurved *S. burkenroadi*
 Antennal spine short, often minute, not buttressed; pleura of abdominal somite 1 rounded, 2-4 angulate, but without laterally recurved marginal spines
 *S. stimpsoni*

Sicyonia laevigata

- a. carapace and first two abdominal somites,
lateral view

(after Burkenroad, 1934)

Sicyonia parri

- b. carapace and abdomen, lateral view

(after Burkenroad, 1934)

Sicyonia brevirostris


- c. lateral view (male)

(after Cobb et al., 1973)

Sicyonia typica

- d. carapace and first abdominal somite, lateral
view

(after Williams, 1984)



Sicyonia dorsalis

- a. carapace and first abdominal somite, lateral view




(after Williams, 1984)

Sicyonia burkenroadi

- b. lateral view (female)
c. third and fourth abdominal somites (female)
(b, after Cobb, 1971; c, after Huff and Cobb, 1979)

Sicyonia stimpsoni

- d. carapace and part of first abdominal somite, lateral view
e. abdominal somites (male)
(d, after Williams, 1984; e, after Huff and Cobb, 1979)



Family Sergestidae

Key to genera and species

1. Fourth and fifth pereopods absent.....*Acetes americanus carolinae*
Fourth and fifth pereopods present..... 2
2. (1) Specialized luminescent modifications of gastrohepatic gland (organs of Pesta)
present; dermal photophores absent; supraorbital and hepatic spines present or
absent*Sergestes*
Specialized luminescent modifications of gastrohepatic gland (organs of Pesta)
absent; dermal photophores present or absent; if present, with or without cuticular
lenses; supraorbital and hepatic spines absent*Sergia*

Genus *Sergestes* H. Milne Edwards, 1830

Key to species

[based on Crosnier and Forest, 1973]

1. Third maxillipeds at most as long as third pereopods..... 2
 Third maxillipeds much longer than third pereopods..... 4
2. (1) Two distal segments of fifth pereopod setose on only one margin (third segment of antennular peduncle equal to or longer than first; petasma lobes short, stumpy) *S. atlanticus*
 Two distal segments of fifth pereopod setose on both margins..... 3
3. (2) Supraorbital spines always present, acute and easily visible..... *S. henseni*
 Supraorbital spines nearly always absent or, when present, miniscule..... *S. paraseminudus*
4. (1) Two distal segments of fifth pereopod setose on both margins..... 5
 Two distal segments of fifth pereopod setose on only one margin..... 6
5. (4) Dactylus and distal half of propodus of third maxilliped with numerous spines forming comb-like structure; processus ventralis of petasma unarmed *S. pectinatus*
 Dactylus and distal half of propodus of third maxilliped armed with spines but not forming comb-like structure; processus ventralis of petasma armed distally with numerous spines *S. sargassi*
6. (4) Dactylus of third maxilliped subdivided into 6 segments and with 2 terminal spines; external margin of exopod of uropod entirely fringed *S. edwardsii*
 Dactylus of third maxilliped subdivided into 4 segments and with single terminal spine; small proximal portion of external margin of exopod of uropod naked 7
7. (6) About 1/3 or a little more of external margin of exopod of uropod naked; first segment of antennular peduncle much shorter than third *S. armatus*
 About 1/6 or 1/7 of external margin of exopod of uropod naked; first segment of antennular peduncle a little longer than third *S. vigilax*

Genus *Sergia* Stimpson, 1860

Key to species

[Based on Crosnier and Forest, 1973]

Third maxillipeds with propodus and dactylus entire.....*S. splendens*Third maxillipeds with propodus and dactylus subdivided, last into 5 to 7 segments.
.....*S. extenuatus*

Sergestes atlanticus

- a. carapace, lateral view
- b. petasma (male)
- c. fifth pereopod (male)

(a, b, after Kensley, 1971; c, after Hansen, 1922)

Sergestes henseni



- d. rostral region
 - e. petasma (male)
- (after Crosnier and Forest, 1973)

Sergestes paraseminudus

- f. rostral region
 - g. petasma (male)
- (after Crosnier and Forest, 1973)

Sergestes pectinatus

- h. carapace, lateral view
 - i. dactylus and distal end of propodus of third maxilliped
 - j. petasma (male)
- (after Kensley, 1971)



Sergestes sargassi

- a. carapace, lateral view
 - b. dactylus and distal end of propodus of third maxilliped
 - c. petasma (male)
- (after Kensley, 1971)

Sergestes edwardsii




- d. anterior region, lateral view
 - e. petasma (male)
 - f. third maxilliped
- (after Crosnier and Forest, 1973)

Sergestes armatus

- g. anterior region, dorsal view
 - h. uropodal exopod
 - i. third maxilliped
- (after Hansen, 1922)

Sergestes vigilax

- j. anterior region, dorsal view
 - k. uropodal exopod
- (after Hansen, 1922)



Sergia splendens


- a. anterior region, dorsal view
- b. dactylus and propodus of third maxilliped
(after Hansen, 1922, as *Sergestes crassus*)

Sergia extenuatus

- c. anterior region, lateral view
- d. petasma (male)
(after Crosnier and Forest, 1973)

Acetes americanus caroliniae

- e. lateral view (female)
(after Williams, 1965a)



Family Luciferidae**Genus *Lucifer* Thompson, 1829****Key to species****[Adapted from Hansen, 1919]**

Distance between labrum and insertion of eye-stalks somewhat or only a little greater than length of eye-stalks with eyes (basal short joint of stalks included); posterior ventral process on sixth abdominal somite in male with its distal part swollen ***L. typus***



Distance between labrum and insertion of eye-stalks almost or more than twice length of eye-stalks with eyes; posterior ventral process on sixth abdominal somite tapering to narrow, obtuse end ***L. faxoni***

Lucifer typus

- a. anterior end, lateral view
- b. male sixth abdominal somite, lateral view
(after Bowman and McCain, 1967)

Lucifer faxoni

- c. lateral view (male)
(after Williams, 1965a)



Suborder Pleocyemata

Infraorder Caridea

Family Atyidae


Genus *Potimirim* Holthuis, 1954

Pereopods without exopods; orbital margin unarmed.....*P. potimirim*

Potimirim potimirim

a. lateral view

(from Abele's personal drawing)



Family Oplophoridae

Key to genera and species
[Adapted from Chace, 1940a]

1. Exopods of at least third maxillipeds and first pair of pereopods foliaceous and often rigid; outer margin of scaphocerite usually armed with series of spines; telson not truncate at tip, but ending in sharp point; eyes large and well pigmented 2
 None of exopods of pereopods foliaceous or rigid..... 3
2. (1) Abdomen with second somite armed with long, carinate posteromesial spines; fifth somite unarmed *Janicella spinicauda*
 Abdomen with second somite unarmed; fifth somite with posteromesial tooth, sometimes small *Oplophorus*
3. (1) Last four abdominal somites, at least, dorsally carinate (usually no straight ridge or carina running entire length of lateral surface of carapace along median lateral line; hind margin of hepatic furrow not usually cut off abruptly by oblique ridge or carina; incisor process of mandible toothed for entire length of cutting edge) *Acanthephyra purpurea*
 Sixth abdominal somite not dorsally carinate (eyes very large and well pigmented; anterior margin of first abdominal somite armed with distinct lobe or tooth overlapping hind margin of carapace; telson terminating in sharp-pointed end-piece laterally armed with spines) *Systellaspis debilis*

Genus *Oplophorus* H. Milne Edwards, 1837

Key to species



- End of scaphocerite barbed on inner margin; posterolateral angle of carapace with no tooth or spine *O. spinosus*
 No barb on end of scaphocerite; posterolateral angle of carapace with prominent spine *O. gracilirostris*

Oplophorus spinosus

a. lateral view (male)
(after Chace, 1940a)

Oplophorus gracilirostris

b. lateral view
(after Kensley, 1972)



Acanthephyra purpurea



- a. lateral view (male)
(after Chace, 1940a)

Janicella spinicauda

- b. lateral view (young male)
(after Chace, 1940a)

Systellaspis debilis

- c. lateral view (male)
(after Chace, 1940a)



Family Pasiphaeidae**Genus *Leptochela* Stimpson, 1860**

Key to species

[Adapted from Williams, 1984]

1. Sixth abdominal somite bearing movable lappet near anterior end of dorsal surface;
third pereopod with exopod reaching nearly or quite to end of ischium *L. carinata*
.....
Sixth abdominal somite lacking dorsal lappet; third pereopod with exopod not
nearly reaching distal end of ischium 2
2. (1) Suborbital angle dentate; orbital margin serrate dorsolaterally..... *L. serratorbita*
Suborbital angle rounded, unarmed; orbital margin usually entire dorsolaterally.... 3
3. (2) Fifth abdominal somite with 1-3 low prominences on dorsal margin... *L. papulata*
Fifth abdominal somite regularly convex or nearly straight in lateral view.....
..... *L. bermudensis*

Leptochela carinata

- a. left third pereopod (male)
 - b. anterior region, lateral view (ovigerous female)
 - c. abdomen (ovigerous female)
- (after Chace, 1976)

Leptochela serratorbita


- ovigerous female:
- d. right third pereopod
 - e. anterior region, lateral view
 - f. anterior part of carapace and eyes, dorsal view
- (after Chace, 1976)

Leptochela papulata

- g. anterior region, lateral view (holotype ovigerous female)
 - h. abdomen (holotype ovigerous female)
 - i. fifth abdominal somite, lateral view (paratype ovigerous female)
- (after Chace, 1976)

Leptochela bermudensis

- ovigerous female:
- j. anterior region, lateral view
 - k. abdomen
 - l. posterior end of sixth abdominal somite
- (after Chace, 1976)



Family Bresiliidae**Key to genera and species**

[Adapted from Chace and Brown, 1978]

Rostrum armed ventrally with at least 1 small tooth; third maxilliped with terminal segment slender, not flattened; first pereopod no longer than second, with elongate fingers *Pseudocheles chacei*

Rostrum unarmed ventrally; third maxilliped with terminal segment broad, flattened; first pereopod longer than second, fingers short and stout *Discias*

Genus *Discias* Rathbun, 1902**Key to species**

[Adapted from Wilson and Gore, 1979]

Second abdominal somite with posterior dorsal spine..... *D. serratirostris*

Abdominal somites without dorsal spines (rostrum narrow with subparallel margins) *D. atlanticus*

Discias serratirostris

ovigerous female:

- a. anterior region, dorsal view
- b. first three abdominal somites, lateral view

(after Wilson and Gore, 1979)

Discias atlanticus

female:






- c. lateral view
- d. anterior carapace, dorsal view

(after Gore and Wilson, 1978)

Pseudocheles chacei

- e. lateral view

(after Kensley, 1983)



Family Eugonatonotidae**Genus *Eugonatonotus* Schmitt, 1926**

A well-developed toothed rostrum immovable; exopods present on pereopods (first two pairs of pereopods chelate, with dark fingertips; ultimate segment of second maxilliped applied as strip along side of penultimate segment; exopod of first maxilliped with distinct flagellum; first chela more robust than second; carpus of second chela entire) *Eugonatonotus crassus*

Family Rhynchocinetidae**Genus *Rhynchocinetes* H. Milne Edwards, 1837**

Rostrum movable; no exopods on pereopods (first two pairs of pereopods chelate, with dark fingertips; ultimate segment of second maxilliped applied as strip along side of penultimate segment; exopod of first maxilliped with distinct flagellum; first chela more robust than second; carpus of second chela entire) *Rhynchocinetes rigens*

Eugonatonotus crassus

a. lateral view

(after Boone, 1927)

Rhynchocinetes rigens

b. lateral view

(after Gordon, 1936)

Rhynchocinetes rigens




c. rostrum (juvenile 2.5 mm)

d. same (ovigerous female, carapace length, excluding rostrum, 3.9 mm)

e. same (juvenile, carapace length, excluding rostrum, 3.4 mm)

f. same (male, carapace length, excluding rostrum, 6.8 mm)

(after Manning, 1961a)



Family Gnathophyllidae

Key to genera and species
[Adapted from Chace, 1972]

Anterolateral angle of carapace not reaching beyond level of antennal spine; spines on distal margin of telson not very unequal; third maxilliped with exopod considerably overreaching endopod; second pereopod with carpus broader than long; 3 posterior pereopods with dactyli nearly as broad as long, not bifid *Gnathophylloides mineri*

Anterolateral angle of carapace reaching distinctly beyond level of antennal spine; intermediate spines on distal margin of telson nearly twice, or more than twice, as long as median pair; third maxilliped with exopod not overreaching endopod; second pereopod with carpus distinctly longer than broad; 3 posterior pereopods with dactyli distinctly longer than broad and bifid *Gnathophyllum*

Genus *Gnathophyllum* Latreille, 1819

Key to species
[Adapted from Chace, 1972]

1. Posterior tooth of dorsal rostral series situated on rostrum anterior to level of orbital margin; color pattern composed of transverse stripes *G. americanum*
- Posterior tooth of rostral series situated on carapace posterior to level of orbital margin; color pattern composed of spots 2
2. (1) Pereopods slender, propodus of third and fourth pairs 12-15 times as long as wide; color pattern composed of dark rings on slightly lighter background *G. circellum*
- Pereopods not usually slender, propodus of third and fourth pairs 7-8 times as long as wide; color pattern composed of innumerable light dots on dark background (posterior pair of lateral telson spines separated by distinct gap from series of posterior spines; stylocerite falling short of level of articulation between first and second segments of antennular peduncle) *G. modestum*

Gnathophyllum americanum

a. lateral view

b. rostrum

(after Manning, 1963)

Gnathophyllum circellum

c. outline of body, lateral view

d. third pereopod

(after Manning, 1963)

Gnathophyllum modestum

e. anterior portion of carapace, lateral view

f. third pereopod

g. telson and left uropods

(after Manning, 1963)

Gnathophylloides mineri

h. carapace, lateral view





i. telson and left uropods

j. third maxilliped

k. first pereopod

l. major chela

(after Schmitt, 1935a)



Family Palaemonidae

Key to genera
[Based on Chace, 1972]

1. Third maxilliped with well-developed exopod..... 2
 Third maxilliped without exopod..... 12
2. (1) Rostrum armed dorsally with series of prominent teeth..... 3
 Rostrum usually unarmed dorsally, at most with 1 or 2 subapical denticles..... 10
3. (2) Carapace with hepatic spine on lateral surface far posterior to anterior margin..... 4
 Carapace without hepatic spine..... 7
4. (3) Telson bearing 2 pairs of terminal spines and usually 1 or 2 pairs of setae..... 5
 Telson bearing 3 pairs of terminal spines..... 6
5. (4) Three posterior pereopods with biunguiculate dactyli.....
 *Brachycarpus biunguiculatus*
 Three posterior pereopods with dactyli simple, without accessory tooth on inferior margin *Macrobrachium*
6. (4) Rostrum without lateral flange; carapace with antennal spine on anterior margin; 3 posterior pereopods 7-segmented, ischium and merus distinct *Periclimenes*
 Rostrum with lateral flange; carapace without antennal spine on anterior margin; 3 posterior pereopods 6-segmented, ischium and merus indistinguishably fused
 *Tuleariocaris neglecta*
7. (3) Carapace with antennal but without branchiostegal spine on or near anterior margin; telson with 3 pairs of terminal spines; second pereopods massive, unequal
 *Periclimenaeus*
 Carapace with both antennal and branchiostegal spines on or near anterior margin; telson with 2 pairs of terminal spines and 1 or 2 pairs of setae; second pereopods elongate, subequal 8
8. (7) Carapace without branchiostegal groove ventral to antennal spine; endopod of first pleopod of male with accessory appendix *Leander*
 Carapace with branchiostegal groove; endopod of first pleopod of male entire, without accessory appendix 9
9. (8) Mandible with palp..... *Palaemon*
 Mandible without palp..... *Palaemonetes*

10. (2) Scaphocerite rudimentary..... *Typoon*
 Scaphocerite well developed..... 11
11. (10) Telson elongate with no dorsal spines; outer margin of uropodal exopod ending in two spines, inner spine movable *Pontoniopsis paulae*
 Telson rather broad, generally with large dorsal spines; one tooth at distal end of outer margin of uropodal exopod *Pontonia*
12. (1) Rostrum not expanded laterally in basal portion; strongly dentate both dorsally and ventrally *Anchistioides antiguensis*
 Rostrum with eavelike expansions over orbits; unarmed ventrally..... 13
13. (12) Second maxilliped with well-developed exopod..... *Veleroniopsis kimallynae*
 Second maxilliped without exopod..... 14
14. (13) Carapace with hepatic spine on lateral surface far posterior to anterior margin.....
 *Lipkebe holthuisi*
 Carapace without hepatic spine..... 15
15. (14) Basal expansions of rostrum anteriorly acuminate; carapace with longitudinal groove extending almost entire length near lateral margin; abdomen with pleura of at least fourth and fifth somites posterolaterally acuminate
 *Pseudocoutierea antillensis*
 Basal expansions of rostrum evenly convex, not acuminate; carapace without longitudinal groove near lateral margin; abdomen with pleura of all 5 anterior somites rounded *Neopontonides beaufortensis*

Genus *Leander* E. Desmarest, 1849

Key to species

Lateral extension of anterior margin of basal antennular segment convex; stylocerite short, barely reaching to middle of basal antennular segment; scaphocerite slender in both sexes; rostrum shallow in both sexes; fingers of second pereopod armed
..... *L. paulensis*

Lateral extension of anterior margin of basal antennular segment concave or straight; stylocerite may reach to distal third of basal antennular segment; scaphocerite slender in male, but broader and tapering less rapidly to apex in female; rostrum shallow in mature male, but very deep in mature female; fingers of second pereopod unarmed *L. tenuicornis*

Genus *Macrobrachium* Bate, 1888

Key to species

[Based on Williams, 1984]

1. Carpi of second pereopods with maximum length as great or greater than meri..... 2
Carpi of second pereopods distinctly shorter than meri..... 4
2. (1) Palms of chelae on second pair of pereopods greatly swollen; prehensile surfaces of noticeably gaping fingers thickly set with long, stiff setae *M. olfersii*
Palms of chelae on second pair of pereopods cylindrical, not greatly swollen; fingers not noticeably gaping but may be hairy 3
3. (2) Fingers of chelae on second pair of pereopods thickly pubescent throughout length; rostrum with teeth extending to tip *M. acanthurus*
Fingers of chelae on second pair of pereopods with scattered hairs, except thicker on fingers along cutting edges; rostrum with toothless daggerlike tip ... *M. ohione*
4. (1) Adult male with chelae of second pereopods equal in shape..... *M. carcinus*
Adult male with chelae of second pereopods very unequal in shape and size; smaller pereopods with fingers gaping, gap being filled by stiff hairs, implanted on cutting edges *M. crenulatum*

Genus *Palaemon* Weber, 1795

Key to species
[Adapted from Holthuis, 1952]

Rostrum high, ventral margin with 3 or 4 teeth; fingers of second pereopods $\frac{2}{3}$ length of palm or shorter *P. northropi*

Rostrum slender, ventral margin with 5 to 7 teeth; fingers of second pereopods more than $\frac{2}{3}$ length of palm *P. floridanus*

Genus *Palaemonetes* Heller, 1869

Key to species
[Adapted from Holthuis, 1952, and Williams, 1984]

1. Fused part of two rami of upper antennular flagellum distinctly longer than free part (branchiostegal spine situated on anterior margin of carapace, just below branchiostegal groove; posterior pair of dosal spines of telson placed midway between anterior pair and posterior margin of telson) *P. paludosus*

Fused part of two rami of upper antennular flagellum shorter than or as long as free part 2

2. (1) Rostrum with 2 teeth of dorsal series behind posterior margin of orbit, teeth reaching to tip, 3 to 5 ventral teeth; carpus of second pereopod in adult female shorter than palm, in male slightly longer (1.1 times) or shorter; dactylus of second pereopod with 2 teeth, immovable finger with 1 tooth on cutting edge *P. vulgaris*

Rostrum with only 1 tooth of dorsal series behind posterior margin of orbit; carpus of second pereopod in adult female much longer than palm (1.3-1.5 times), in male almost as long as whole chela; dactylus of second pereopod with or without single tooth, fixed finger without tooth on cutting edge 3

3. (2) Rostrum with dorsal teeth reaching to often bifurcate tip, 4 or 5, seldom 3, ventral teeth; dactylus of second pereopod with tiny and sometimes blunt tooth *P. intermedius*

Both margins of rostrum with unarmed stretch before dagger-shaped tip, 2 to 5, generally 3, ventral teeth; fingers of second pereopod without teeth on cutting edge *P. pugio*

Genus *Periclimenaeus* Borradaile 1915

Key to species
[Adapted from Chace, 1972]

1. Telson with anterior pair of dorsal spines arising from anterior fourth of segment.. 2
Telson with anterior pair of dorsal spines arising at end of anterior third of segment or posterior thereto 8
2. (1) Movable finger of major chela of second pereopod extending distinctly beyond tip of immovable finger *P. chacei*
Movable finger of major chela of second pereopod extending very slightly beyond or reaching to tip of immovable finger 3
3. (2) Telson with 3 pairs of distal spines inserted in continuous line..... 4
Telson with lateral pair of distal spines inserted distinctly anterior to intermediate and mesial pairs 6
4. (3) Rostrum with ventral tooth; carapace with small denticle or sharp tubercle posterior to orbit; scaphocerite with anterolateral tooth distinctly overreaching blade *P. caraibicus*
Rostrum unarmed ventrally; carapace without postorbital denticle; scaphocerite with anterolateral tooth not overreaching blade 5
5. (4) Third maxilliped with 2 distal segments broad, penultimate about two and one-half times as long as broad; first pereopod with movable finger tapering to tip, not strongly convex, carpus about one and one-third times as long as chela; minor second pereopod with movable finger elongate, not semicircular ... *P. ascidiarum*
Third maxilliped with 2 distal segments unusually slender, penultimate about five times as long as broad; first pereopod with movable finger strongly convex, carpus about one and one half times as long as chela; minor second pereopod with movable finger short and broad, nearly semicircular *P. pearsei*
6. (3) Major second pereopod with large tooth on opposable margin of immovable finger fitting into cavity in movable finger; minor second pereopod with fingers longer than palm *P. bermudensis*
Major second pereopod with large tooth on opposable margin of movable finger fitting into cavity in immovable finger; minor second pereopod with fingers much shorter than palm 7
7. (6) First pereopod unusually long and slender, carpus nearly twice as long as chela..... *P. perlatus*
First pereopod not abnormally long or slender, carpus less than one and one-half times as long as chela (rostrum with 10-12 dorsal teeth; telson with posterior pair of dorsal spines arising from posterior half of segment) *P. wilsoni*

8. (1) Scaphocerite without anterolateral tooth; third pereopod with dactylus bifid..... *P. schmitti*

Scaphocerite with anterolateral tooth; third pereopod without distinct accessory tooth on inferior margin of dactylus 9

9. (8) Rostrum with 4 dorsal teeth; scaphocerite with large anterolateral tooth reaching about to level of distal margin of blade *P. atlanticus*

Rostrum with 1 or 2 dorsal teeth; scaphocerite with small anterolateral tooth falling far short of level of distal margin of blade *P. maxillulidens*

Genus *Periclimenes* Costa, 1844

Key to species
[Adapted from Chace, 1972]

1. Antennular peduncle with only 1 spine at distolateral angle of basal segment (in addition to stylocerite) 2
 Antennular peduncle with 2 or more spines at distolateral angle of basal segment (in addition to stylocerite) 9
2. (1) Carapace with anterior margin unarmed (antennal spine absent; third pereopod with distinctly biunguiculate dactylus) *P. longicaudatus*
 Carapace armed with antennal spine below suborbital lobe 3
3. (2) Fifth abdominal pleuron with posteroventral angle pointed; telson with anterior pair of dorsal spines arising about one-third of length from base of segment; scaphocerite with distal spine overreaching distal margin of blade . *P. americanus*
 Fifth abdominal pleuron with posteroventral angle rounded; telson with anterior pair of dorsal spines arising at, or posterior to, midlength of segment; scaphocerite with distal spine rarely reaching as far as distal margin of blade, usually falling far short 4
4. (3) Third abdominal somite strongly produced posteromesially into laterally compressed hump 5
 Third abdominal somite sometimes moderately produced posteromesially but never forming laterally compressed hump 6
5. (4) Third pereopod with dactylus simple and considerably more than one-third as long as propodus *P. magnus*
 Third pereopod with dactylus distinctly biunguiculate and not more than one-fourth as long as propodus (carapace with hepatic spine usually arising at, or posterior to, level of posterior tooth of rostral series; carpus of major second pereopod usually more than half as long as chela) *P. pedersoni*
6. (4) Rostrum elongate, more than four times as long as maximum height, one or more of ventral teeth prominent (first pereopod with carpus not noticeably longer than chela; sixth abdominal somite less than twice as long as fifth and shorter than telson; scaphocerite with blade far overreaching distal spine; second pereopod with fingers slightly more than half as long as palm, carpus about one-fifth as long as chela) *P. pandionis*
 Rostrum subtriangular in lateral view, less than four times as long as maximum height, ventral teeth inconspicuous or absent 7

7. (6) Telson with dorsal spines rather large and distinct; major second pereopod with fingers no more than one-fourth as long as palm *P. harringtoni*

Telson with dorsal spines minute and inconspicuous; major second pereopod with fingers more than half as long as palm 8

8. (7) Sixth abdominal somite about twice as long as fifth and longer than telson; telson with anterior pair of dorsal spines arising at about midlength of segment; major second pereopod with movable finger not perceptibly stouter than immovable finger *P. iridescens*

Sixth abdominal somite slightly more than half again as long as fifth and shorter than telson; telson with anterior pair of dorsal spines arising at least two-thirds of length from base of segment; major second pereopod with movable finger usually stout, nearly twice as high as immovable finger *P. rathbunae*

9. (1) Posterior tooth of rostral series far removed from second tooth and from posterior margin of orbit; third pereopod with dactylus deeply biunguiculate *P. yucatanicus*

Posterior tooth of rostral series not widely separated from second tooth, situated slightly posterior or anterior to level of orbital margin; third pereopod with dactylus simple or very obscurely biunguiculate (scaphocerite less than twice as long as broad; major second pereopod with fingers less than one-fourth as long as palm) *P. perryae*

Genus *Pontonia* Latreille, 1829

Key to species
[Adapted from Chace, 1972]

1. Carapace pubescent, cervical groove well marked; major second pereopod with large rounded tooth on movable finger fitting into completely enclosed socket in immovable finger *P. unidens*

Carapace not pubescent, without cervical groove; enlarged tooth on movable finger of major second pereopod, if present, triangular and fitting into shallow, partially open socket in immovable finger 2
2. (1) Telson with dorsal spines minute, inconspicuous..... *P. domestica*

Telson with dorsal spines well developed (three posterior pereopods with dactyli stout, inferior margin convex) *P. margarita*

Genus *Typton* Costa, 1844

Key to species
[Adapted from Chace, 1972]

1. Telson with posterior pair of dorsal spines arising anterior to midpoint of segment; exopod of uropod with outer margin serrate in distal portion *T. prionurus*
 Telson with posterior pair of dorsal spines arising at, or posterior to, midpoint of segment; exopod of uropod with outer margin entire, not serrate distally 2
2. (1) Antennal spine broad, toothlike in lateral view, not spiniform; both second pereopods with movable fingers highly arched, nearly semicircular; major second pereopod with carpus crenulate on proximal portion of angulate margin *T. tortugae*
 Antennal spine strong, spiniform; second pereopods with movable fingers only moderately convex, not nearly semicircular; major second pereopod with carpus not crenulate on angulate margin 3
3. (2) Anterior margin of carapace produced anteriorly to level of tip of antennal spine; exopod of uropod with outer margin rather regularly convex throughout 4
 Anterior margin of carapace less produced, not nearly reaching level of tip of antennal spine; exopod of uropod with outer margin nearly straight in distal half .. 5
4. (3) Rostrum deepest near midlength, ventral margin forming obtuse angle in lateral view; mandible with well-developed incisor process; third pereopod with dactylus bearing small accessory tooth on inferior margin, not clearly symmetrically bifid *T. carneus*
 Rostrum not deepening near midlength, ventral margin straight or convex; mandible without incisor process; third pereopod with dactylus bearing large accessory tooth on inferior margin, nearly symmetrically bifid *T. gnathophylloides*
5. (3) Mandible with incisor process well developed and distally crenulate, molar process tapering distally; major second pereopod with movable finger bluntly hammer-shaped, not noticeably twisted *T. vulcanus*
 Mandible with incisor process reduced to low angulate unarmed lobe, molar process not tapering distally; major second pereopod with movable finger forming pointed hook twisted into plane nearly perpendicular to that of palm *T. distinctus*



Leander paulensis

mature female:

- a. rostrum and anterior portion of carapace
- b. second pereopod
- c. scaphocerite
- d. antennular peduncle
(after Manning, 1961b)

Leander tenuicornis

- e. rostrum and anterior portion of carapace
(adult female)
- f. second pereopod
- g. scaphocerite (adult female)
- h. antennular peduncle (adult female)
(e, g, h, after Manning, 1961b; f, after Holthuis, 1952)



Macrobrachium olfersii

a. lateral view

(after Holthuis, 1952)

Macrobrachium acanthurus

b. lateral view

(after Holthuis, 1952)

Macrobrachium ohione





c. lateral view

(after Holthuis, 1952)

Macrobrachium carcinus


d. lateral view

(after Holthuis, 1952)



Macrobrachium crenulatum

- a. anterior region, lateral view
 - b. major second pereopod (adult male)
 - c. minor second pereopod (adult male)
- (after Holthuis, 1952)





Palaemon northropi

- a. anterior region, lateral view
 - b. second pereopod
- (after Holthuis, 1952)

Palaemon floridanus

- c. lateral view
- (after Holthuis, 1952)



Palaemonetes paludosus

- a. anterior region, lateral view
- b. antennule
- c. telson

(after Holthuis, 1952)

Palaemonetes vulgaris

- d. anterior region, lateral view
- e. second pereopod (female)
- f. second pereopod (male)
- g. fingers of second pereopod (female)

(after Holthuis, 1952)

Palaemonetes intermedius





- h. anterior region, lateral view
- i. fingers of second pereopod (female)

(after Holthuis, 1952)

Palaemonetes pugio

- j. anterior region, lateral view
- k. fingers of second pereopod (female)

(after Holthuis, 1952)



Periclimenaeus chacei

- a. carapace, lateral view (male)
- b. anterior portion of chela (male)
- c. telson and left pair of uropods
- d. second pereopod (male)
(after Abele, 1971)

Periclimenaeus caraibicus

- e. anterior region, lateral view
- f. antenna
- g. major second pereopod
(after Holthuis, 1951b)

Periclimenaeus ascidiarum

- h. anterior region, lateral view
- i. major second pereopod, outer view
- j. minor second pereopod
- k. third maxilliped
(after Holthuis, 1951b)

Periclimenaeus pearsei

- l. carapace, lateral view
- m. minor second pereopod
- n. first pereopod
(after Holthuis, 1951b)