

1966

With best wishes,

LIBRARY
DIVISION OF SCIENCE

Drs. L. J. G.

Reprinted from *Australian Zoologist*, XIII (3), July 6, 1966

259

A REVIEW OF THE AUSTRALIAN MAJID SPIDER CRABS (CRUSTACEA, BRACHYURA)

BY D. J. G. GRIFFIN

(Zoology Department, University of Tasmania, Hobart)
(Plates XV-XVII & Text-figures 1-3)

SUMMARY

An historical account is given of taxonomic studies of spider or masking crabs in Australia and overseas. The basis of the review is a key to all but two of the species known to occur in Australia. The key includes information on synonymies, geographic and bathymetric distribution and references to descriptions and illustrations of the species. Characters which are important in the classification are briefly reviewed. Zoogeographical relationships of the fauna are discussed.

The family Majidae is currently divided into seven subfamilies mainly on the basis of orbital configuration, form of the rostrum, abdomen and first pleopod of the male. All but one of the subfamilies are represented in the Australian fauna which is considered to comprise 95 species in 45 genera. This is about twice the number of species and genera listed by Haswell (1882c) in his "Catalogue." About one-third of the species have been recorded from Australia on only a single occasion. Numerous genera and species are in need of detailed reinvestigation.

The fauna is rather clearly partitioned into a tropical group with widespread Indo-West Pacific relationships and a temperate group related to tropical Australia and/or the Indo-West Pacific rather than to temperate regions outside Australia. There are no clear boundaries between these two faunas but rather quite broad transition areas. Thirty-seven species and five genera, most of which are temperate, are restricted to Australia.

1. INTRODUCTION

One of the most characteristic features of spider crabs of the family Majidae is the presence on the carapace and legs of special curled or "hooked" hairs which aid in the attachment of various kinds of epifauna and flora, especially seaweeds (for example see McNeill 1923). These organisms are placed in position by the crab with the aid of the chelipeds which are able to reach up on to the dorsal surface of the carapace. For this reason majid crabs are sometimes called "masking crabs" or "seaweed crabs." Of special interest in these crabs are the orbits which may be expanded in various ways and surrounded by a seemingly complex array of spines. The legs are often long and slender and the carapace usually triangular or pyriform. Spider crabs range in size from a few millimetres to more than a metre in carapace length and are found in almost all seas and oceans. They form a relatively important part of the benthos and may be locally very abundant on the shelf although some species extend considerable distances down the continental slope to depths as great as 1000 fathoms.

Substantial revisions of the family Majidae have been undertaken on a world-wide scale by Dana (1851), Miers (1879c), Alcock (1895) and Balss (1929). Sakai (1938), Stephensen (1945) and Garth (1958) have adapted Balss's scheme to the faunas of Japan, the Iranian Gulf and America respectively. In the course of these revisions the Majidae have been arranged in several very different fashions (see historical reviews by Miers 1879c, and Garth 1958). Early workers such as Dana and Miers grouped the species in several families and a large number of subfamilies—Dana's arrangement comprised five families and no less than 27 subfamilies. Alcock's was the first of these major revisions to contain the species in a single family. This was divided into four sub-families and seven "alliances." Whereas revisors before Balss had proceeded by rearrangement of previous schemes, Alcock's groups of genera were left almost untouched by Balss, his alliances in some cases being elevated to subfamilial rank much as they were conceived. The number of subfamilies currently recognised stands at seven following the work of Garth.

APR 13 1967

CARDED

Many morphological characters have been used in these different classifications and some have been shown to be unreliable guides to phylogenetic relationships (see Section III). Characters concerned with the form of the orbit still dominate the classification but Garth has found, in the Pacific American fauna, groups of genera showing vast agreements in the form of the male first pleopod. This appendage is now realised to be of great value in the taxonomy of nearly all families of crabs.

The first Australian spider crabs to be described were made known by the European naturalists Herbst, Latreille and H. Milne Edwards in the early nineteenth century. In the second half of that century the work of Dana, Stimpson, Miers and Haswell resulted in substantial additions to knowledge of the Australian spider crab fauna. Dana in 1852 dealt with the Crustacea of the U.S. Exploring Expedition of 1838-1842, Stimpson with the Crustacea of the North Pacific Exploring Expedition led by Ringgold and Rodgers (Stimpson 1857—full report not published until 1907). Haswell published six papers between 1879 and 1882 and in that year (Haswell 1882c) provided the first list of all species known from Australia. But Miers's important reports on the material collected by the "Alert" in 1881-2 (Miers 1884) and by the "Challenger" in 1873-76 (Miers 1886) outdated Haswell's list to some extent. So too did the reports of W. T. Calman (1900) and A. Ortmann (1894) on collections of Brachyura from Torres Strait.

Haswell's work at the Australian Museum was followed by that of Thomas Whitelegge whose report on the Crustacea collected off eastern Australia by the "Thetis" (Whitelegge 1900) may be particularly mentioned. W. H. Baker (1905, 1906) dealt with South Australian Brachyura and S. W. Fulton, F. E. Grant and Allan R. McCulloch all paid some attention to south-eastern Australian species in the first and second decades of the present century. In the 1920's, Herbert M. Hale (1924, 1927, 1929) reported on southern and western species. The numerous reports, from the 1890's to the 1920's, by Mary J. Rathbun of the Smithsonian Institution at the same time added several species to the Australian lists and extended the known ranges of many others; two reports (Rathbun 1914, 1924) dealt with collections from Western Australia. S. K. Montgomery's report on the Brachyura of the Percy Sladen Trust Expedition to the Abrolhos (Montgomery 1931) and Dr. Heinrich Balss's paper (1935) on the material collected by the Hamburg Museum's Expedition to southwestern Australia in 1905 are the latest sources of information on western spider crabs. The contemporary and later work of Frank A. McNeill and Melbourne Ward dealt mainly with north-eastern species.

The present situation with regard to the Australian majid spider crab fauna is thus as follows. No revisionary treatment has ever been attempted and no comprehensive guide has appeared since the time of Haswell. Until recently only McCulloch (1908, 1913) had completed revisions at the generic level. Nearly all reports dealing with majids have merely added species or extended known geographic ranges. The most important recent references to Australian species are those primarily treating overseas faunas (e.g. Sakai 1938, 1965; Buitendijk 1939; Barnard 1950; and Forest & Guinot 1961). The number of known valid genera and species has doubled since the time of Haswell and collections have continued to accumulate at various museums, particularly the Australian Museum, Sydney, and the Western Australian Museum, Perth. Several areas of the Australian coastline remain uninvestigated, particularly that west of Torres Strait south to Broome and most of the Great Australian Bight. The Australian Museum possesses a fine catalogue of all described genera and species. Not the least important of the many problems which therefore remain unanswered is the question of whether the Australian genera can be grouped, on the basis of the male first pleopods (unknown in the vast majority of local species), in an arrangement which parallels that found by Garth.

The present paper attempts to bring together available information on the Australian Brachyura of the family Majidae. This information is presented in the form of a key to genera and species with distributional and other information included. The following discussions centre around this key. Current problems are pointed out but there is no intention here of giving a general revision. Adequate treatments of the morphology of members of this family are

given by Rathbun (1925), Garth (1958) and Griffin (in press, b). Terminology, unless otherwise indicated, follows that used by Rathbun and Garth and by Griffin in previous papers. Although most information used in the key is taken from the literature, a considerable amount of material comprising most of the species represented in Australia, has been examined at the Australian Museum over the past three years.

II. THE FAMILY

The Majidae belong, with the Hymenosomidae (flat-back crabs) and the Parthenopidae (caltrop crabs), to the superfamily Oxyrhyncha. All three families possess an anteriorly narrowed and produced carapace with prominently expanded branchial regions, large epistome, quadrate buccal cavity and longitudinally folded antennules. The characters which chiefly distinguish majid crabs from other families of the Brachyura are: (1) the possession of specially mobile chelipeds; (2) fusion of the well-developed second article of the antenna (usually called the basal antennal article) to the epistome and often also to the front; (3) the comparatively incomplete orbits; (4) articulation of the palp of the external maxillipeds at the anteromedial angle of the merus; and (5) the much greater length of the first pleopods of the male relative to the second pair. The family has usually been considered in the past to share with other Oxyrhyncha nine pairs of gills on each side but Hartnoll (1964) has recently shown that this number is sometimes reduced in the Majidae to seven or eight.

It should be mentioned that the term Maioidea used by Dana and Miers is partially synonymous with Oxyrhyncha (as now understood) and not with the term Majidae (until recently frequently spelt Maiidae). In Dana's and Miers's time the Maioidea included only the Majidae and the Parthenopidae; de Haan's (1839) addition of the Hymenosomidae did not gain general acceptance until after Rathbun (1925).

As to the evolution and relationships of the Oxyrhyncha, Glaessner (1960: 46) states, "The Oxyrhyncha, or spider crabs, are . . . unrepresented before the Tertiary except by unidentifiable fragments. The pointed rostrum . . . and the prominent mesogastral-cardiac ridge, together with the elongate cephalothorax, place this group much closer to the Oystomatidae than to the Brachyrhyncha, but it is more advanced in the organisation of its mouthparts."

III. THE SUBFAMILIES

The subfamilial arrangement adopted here is that proposed by Garth (1958) which is a modification of the schemes of Alcock (1895) and Balss (1929). In the delimitation of subfamilies and in the arrangement of genera within them, particular attention has been paid to the structure of the orbit, i.e., the degree of expansion of the anterolateral part of the carapace above the origin of the eyestalk (the "supraorbital eave"), the presence or absence, and form of, a spine behind the eave (the "postorbital spine") and the degree of expansion of the basal antennal article. These are characters used before Alcock by Miers (1879c) and to some extent by Dana (1851). H. Milne Edwards (1834) considered the length of the ambulatory legs to be important and the classification proposed by de Haan (1839) was based on the shape of the merus of the third maxillipeds. Use of these latter characters was criticised by Miers. In 1861 Claus proposed a classification based on the form of the basal antennal article. Use of this character was criticised by Stimpson in 1871. Calss laid particular stress on a further feature associated with the structure of the orbit: the presence or absence of a spine (the "intercalated spine") between the eave and the postorbital lobe. However, later workers on the group (Sakai 1938 and Garth 1958) have considered that Balss laid too much stress on this feature (see below). The relative size of this spine is sometimes variable with either growth or geographical locality (Garth 1958; Griffin, unpublished). Balss also introduced three other characters: the number of free segments of the abdomen (basically seven but sometimes reduced by

coalescence or fusion involving segments four to six inclusive), the degree of development of the interantennular spine (the true rostrum of the zoea) and the degree of fusion of the (pseudo) rostral spines.

In the key presented here genera within each subfamily are arranged in a series which begins with forms with a double rostrum and seven free abdominal segments in both sexes and ends with those in which the rostral spines are partly or wholly fused and the number of free segments in the abdomen is reduced. In this respect the key resembles those provided by Garth (1958). Implicit in such a presentation is the concept that parallel evolution has proceeded independently within each subfamily. The genera at the beginning of the series are considered to be primitive and those at the end advanced. This is in accordance with the ideas of Balss who considered that a well developed interantennular spine and the presence of an intercalated spine were also primitive features.


Some aspects of orbital structure require further clarification. The term "commencing orbit," used in respect of some of the Ophthalmiinae, Acanthonychinae and Pisinae, is intended to contrast with the unformed orbits of the Inachinae and some members of the Acanthonychinae on the one hand and with the complete or almost complete orbits of the Majinae and Mithracinae on the other. The commencing orbit is thus intermediate between two extremes. It involves partial enlargement of the supraorbital eave, most commonly by the development of a prominent spine either anteriorly or, less often, posteriorly; this spine scarcely conceals the eyestalk from dorsal view.

In some inachines there is above the origin of the eyestalk a variously developed spine, considered by Balss to represent the intercalated spine. However, in the Inachinae, this spine is very seldom separated by distinct fissures from the surrounding parts of the orbit as it is in the Majinae and some Mithracinae. Further, in some inachine genera which appear to be natural groups, some species possess a spine above the eyestalk and some do not. It would seem better to call this spine merely a supraorbital spine and to disregard, at least for the present, any possible homology between it in the Inachinae and the intercalated spine of other majids.


A similar difficulty exists in many of the Pisinae. In some the postorbital lobe is provided on the upper anterior edge close to the base with a prominent accessory lobe and in a few species this lobe is quite distinct from the postorbital lobe. In other pisines where the eave is virtually unexpanded a small denticle is present on the supraorbital margin. There this lobe or denticle appears to be the intercalated spine and is generally so treated here.

To sum up, emphasis on the intercalated spine led Balss to divide the Inachinae and the Pisinae each into two further subfamilies. In at least the Inachinae it is difficult to work out the homologies of a spine above the orbit in those species in which it is present and in the Pisinae Garth has found that such a division, in the Pacific American forms, is not supported by the male first pleopods. The term supraorbital spine is used, especially in the Inachinae, to denote a spine above the orbit where homologies are not clear. In other groups the term preorbital spine is restricted to mean an anterior outgrowth of the supraorbital eave and the term antorbital spine is used to denote a posterior outgrowth (see figure in Griffin in press, a).


Figure 1. Generalised morphology of the subfamilies Inachinae, Ophthalmiinae and Acanthonychinae. Front of the carapace from above at left and from below at right; whole carapace from above in the middle. Abbreviations: a_1 , antennule; a_2 , antenna; a.o., antorbital spine; b.a., basal antennal article; e, eye; i, intercalated spine; m, mouthfield; p.o., postorbital lobe; pr, preorbital spine; r, rostrum; s.o., supraorbital eave.



INACHINAE



OPHTHALMIINAE



ACANTHONYCHINAE

The characters of the six subfamilies represented in Australia are now given (the number of the couplet in the key at which genera of each subfamily begin is indicated in brackets after the name):

INACHINAE (4): Orbit undeveloped, or eave weakly expanded only; a supraorbital spine sometimes present. A postorbital spine sometimes present but affording no concealment to the cornea of the retracted eyestalk. Basal antennal article extremely slender and usually long. Eyestalk very long and visible almost to its base in both dorsal and ventral view. Carapace subtriangular or sometimes subpyriform or occasionally circular. Rostrum sometimes of a single spine, and frequently short. Ambulatory legs frequently extremely long and slender.

OPHTHALMIINAE (26): Orbit consisting of a well developed and laterally expanded eave or of a greatly elongated spine. Postorbital spine short. An intercalated spine never present. Basal antennal article moderately expanded only. Eyestalk very long, often concealed in dorsal view but always largely visible in ventral view. Carapace elongate and often truncate in front and provided with a medial expansion or spine posteriorly. Rostral spines distinct and usually short. Ambulatory legs seldom very long.




ACANTHONYCHINAE (29): Orbit undeveloped, eave rounded or forwardly produced as a preorbital spine. Postorbital spine, if present, simple and affording no concealment to eyestalk. Basal antennal article not very wide and characteristically truncate triangular. Eyestalks extremely short and often sunk in sides of rostrum. Carapace basically pyriform but characteristically twice expanded laterally at hepatic and branchial margins. Rostrum consisting either of two distinct spines or sometimes of a huge "beak." Ambulatory legs short or at most of moderate length.

PISINAE (36): Orbit with a weakly expanded eave either produced anteriorly as a preorbital spine or acute or sometimes posteriorly acute, or else weakly expanded midway along and almost confluent with the postorbital lobe. Intercalated spine present or absent, sometimes a small denticle or lobe close to the postorbital lobe. Postorbital lobe always well developed and cupped but not completely concealing the eyestalk from dorsal view. Basal antennal article slightly to broadly expanded and usually produced into a spine anterolaterally. Eyestalk typically short and bulbous with a large cornea. Carapace always pyriform although sometimes very wide. Rostrum frequently bifid for distal half only. Ambulatory legs often very long and slender.



MAJINAE (62): Orbit well developed, comprising above a laterally expanded eave which is always acute posteriorly and sometimes produced into a spine, an intercalated spine and a postorbital spine which is sometimes simple and sometimes cupped and occasionally armed with an accessory spine on the upper anterior edge near the base. Basal antennal article moderately broad and rectangular and frequently armed with spines at both anteromedial and anterolateral angles. Eyestalks of moderate length and generally slender; only the distal half is visible in dorsal view. Carapace pyriform and sometimes broad. Rostrum always of two distinct spines. Ambulatory legs generally of moderate length.

MITHRACINAE (81): Orbit extremely well developed with the eave and basal antennal article laterally expanded and usually tubular, completed behind by the prominent postorbital lobe which is often cupped. An intercalated spine is sometimes present. Eave sometimes forwardly produced into a preorbital spine or lobe, and sometimes prolonged posteriorly into an antorbital spine. Eyestalk moderately long but only the distal portion generally visible in either dorsal or ventral view. Carapace basically pyriform but anteriorly broadened due to expansion of orbits. Rostrum of two distinct spines which are often contiguous and sometimes fused into a broad lamella. Ambulatory legs seldom of great length.



Figure 2. Generalised morphology of the subfamilies Pisinae, Majinæ and Mithracinæ. Arrangement and abbreviations as in text-fig. 1.



PISINAE



MAJINAE



MITHRACINAE

The seventh subfamily, the OREGONIINAE contains three genera which are boreal in distribution and not represented in Australia. They resemble the Inachinae in most features but are characterised by the terminally broadened male abdomen, the seventh segment being quadrate and deeply inserted into the sixth segment, and the longitudinally grooved male first pleopod, the groove being margined by rows of filamentous setae. A detailed treatment is given by Garth (1958).

IV. THE GENERA

The characters which are of value at the generic level are essentially the same as those used at the subfamilial level and already discussed. It need only be emphasised that the application of these characters at the generic level is invariably more rigorous than at the higher levels. Thus congeneric species are usually considered to share very similar orbital structure, form of the rostrum and of the male first pleopods and to agree in the number of free segments in the abdomen. They also generally resemble each other closely in carapace shape and ornamentation, form of the third maxillipeds, shape of the male abdomen (a character introduced by Shen in 1932) and often shape of the male chela but seldom in actual size and in relative length of the ambulatory legs.

The remaining part of this section and most of the succeeding one will be devoted to a discussion of the several lower taxa of the Australian Majidae which are considered to pose important problems at the present time.

Of the 24 majid genera listed by Haswell (1882c), ten now either have quite different meanings or are known under different names. These are as follows:

1. *Stenorhynchus* Lamarck. Of the three species included here by Haswell, two have been shifted to *Achaeus* and one, *S. curvirostris*, has not been collected since its discovery (see next section). Lamarck's genus is currently recognised for two species confined to the Americas and north-west Africa (Garth 1958, Garth & Holthuis 1963); since Haswell mentions the rostrum being formed of two spines, not one, it is reasonable to assume that he intended to refer to *Stenorhynchus* Latreille (now considered a synonym of *Macropodia* Leach, 1814).
2. *Halimus* Latreille. The species included here by Haswell are now known as species of *Naxia* Latreille, *Halimus* being a junior synonym of that genus. The composition of the abdomen (of both sexes) and several other characters in these species require investigation to determine their relationships to the New Zealand representative of the genus (Griffin, in press, b). The comparatively well developed orbits and distally expanded ambulatory propodi set this genus apart from other inachines. Haswell's subgenus *Microhalimus*, with its single species, *M. deflexifrons*, was included as a subgenus of *Naxia* by McCulloch (1908). For the sake of convenience, *Microhalimus* is here not given subgeneric status though it is true that *M. deflexifrons* appears to differ in some important features of the orbit from other species of *Naxia*, particularly in the number and arrangement of the spines above and behind the orbit. Such disagreement is especially obvious when it is compared with those species with which it agrees in the slight expansion of the ambulatory propodi.
3. *Naxia* H. Milne Edwards. Haswell listed one species under this genus, *Pisa serpulifer* Guérin. This was transferred to *Naxioides* A. Milne Edwards (of which *Naxia* Miers is a synonym) by Rathbun (1914) and later (Rathbun 1924) set apart in a genus of its own, *Paranaxia*; H. Milne Edwards's genus is a synonym of the latter.
- 4-6. *Gonatorhynchus* Haswell, *Paramithrax* H. Milne Edwards and *Chlorinoides* Haswell. *Gonatorhynchus* has recently (Griffin 1963b) been sunk in *Paramithrax* which has been recognised as monotypic. There appear to be some important similarities in orbital structure between *P. barbicornis* and *Anacinetops stimpsoni* although the species are at present placed in different subfamilies. The meaning attributed *Paramithrax* in Haswell's time has also been changed through the removal of several species to *Chlorinoides* (see Griffin in press, a). These latter species are the ones often placed in *Acanthophrys* A. Milne Edwards, a genus synonymous with *Hyastenus* White.

7-9. *Egeria* Latreille, *Cyclomaia* Stimpson and *Parathoe* Miers. These have been replaced respectively by *Phalangipus* Latreille, *Cyclax* H. Milne Edwards and *Perinia* Dana without change in meaning. In addition *Eucinetops* Stimpson no longer appears in the Australian lists, *Anacinetops* Miers having been accepted for *E. stimpsoni* Miers (see Balss 1935). One genus listed by Haswell among the Oxyrhyncha, the monotypic *Pleurophricus* A. Milne Edwards, has recently been included among the Cymopodidae (now known as the Palicidae—see Holthuis 1962: 244, 249) by Balss (1957: 1662) and is not treated here.

10. *Chorilabinia* Lockington. This genus was listed by Haswell under the name *Chlorolabinia*; it will be discussed in more detail at the end of this section.

A nomenclatural point which should be mentioned here concerns the genus *Camposcia*. Balss (1957: 1620) lists this as *Camposcia* Desmarest, 1823. Yaldwyn (pers. comm.) has kindly informed me that Desmarest's reference (*Dict. Sci. nat.* 28: 259, 262) is to an MS name of Leach, *Camposia*. Other authors before Latreille (1829) (in Cuvier's *Reg. Anim.* IV. (2): 60) also used this spelling. However, it appears that Latreille who used the name *Camposcia*, was the first to include a species in the genus. *Camposcia* of Desmarest and others would, therefore, appear to be "nomen nudum." The species included in *Camposcia* Latreille, 1829 was *C. retusa* which was listed without comment by H. Milne Edwards (1834:283) as *Camposcia retusa* Latreille, 1829. All subsequent workers used these spellings which are therefore followed here.

Among other genera which need re-examination in regard to their distinctness and relationships are *Xenocarcinus*, *Zewa*, *Phalangipus*, *Doclea*, *Antilibinia*, *Hyastenus* and *Micippa*.

1. *Xenocarcinus* White. Sakai (1938: 324) has placed this genus in the subfamily currently known as the Mithracinae because he considers that at least two of the Japanese species show traces of an intercalated spine. Although this structure is unknown in the Acanthonychinae, *Xenocarcinus* is here placed in that subfamily because of its numerous resemblances to other acanthonychinines.

2. *Zewa* McCulloch. This genus was introduced in 1913 for a (new) species from north-eastern Australia. At the same time the earlier described *Pseudomicippe varians* Miers was included because, as McCulloch noted, previous workers had expressed some doubt about the wisdom of considering this latter species to be congeneric with *P. tenuipes* A. Milne Edwards. McCulloch considered the presence of a large lobe above the orbit and of a spine on the posterior border of the carapace important characters in this genus. Balss (1957), however, lists *Zewa* as a synonym of *Pseudomicippe* Heller. Although comparisons of descriptions and figures of the three species (see Buitendijk 1939: 255, pl. VIII figs. 3 and 4; and Sankarankutty 1962: 160, figs. 17-23 for *P. tenuipes*) certainly inclines me towards Balss's view, I retain McCulloch's genus pending a fuller investigation of the Australian species.

3. *Phalangipus* Latreille and *Doclea* Leach. These two genera are among those pisines in which positive identification of an intercalated spine is difficult. In both genera the eave is poorly expanded and separated by a more or less wide hiatus from the postorbital lobe. *Phalangipus* is usually regarded as possessing an intercalated spine and *Doclea* to be lacking it. Thus *P. australiensis*, which possesses a small denticle in the hiatus, is widely separated in the key from *D. profunda*, in which there is no denticle. But among species of *Doclea*, the South African *D. muricata* (Herbst) (see Barnard 1950: 49, fig. 11a) does possess such a denticle just forward of the postorbital lobe. Some re-arrangement of species may be necessary. Rather striking differences in carapace shape exist between the species of *Doclea* (see Sakai 1938: pl. XXXVII) just as in *Naxioides*.

4. *Antilibinia* Macleay. Three species are at present included in this genus, one each from South Africa, the Philippines and Australia. The last two were both described by Rathbun from a few specimens and have not been since discussed on the basis of additional material. Barnard (1950:37) doubted that the Australian *A. lappacea* belongs in the same genus as the South African *A. smithii* because of differences in carapace shape and length of the rostrum; he also considered that the latter species might well be shifted to the American *Taliepus* A. Milne Edwards, again because of similarities in carapace shape. Indeed both

Antilibinia and *Taliepus* have in the past been considered subgenera of *Epiactus* H. Milne Edwards (see Garth 1958:207). A new genus may be required for the Australian and Philippine species.

5. *Hyastenus* White. As the basis for Balss's (1929) *Hyasteniinae*, this extremely large genus (Balss in 1935 listed 38 species) is usually considered to be characterised especially by an absence of an intercalated spine from the orbit and by the distinctness of all seven abdominal segments. However, Laurie (1906) has noted that in his *H. irami* there is a distinct lobe between the eave and postorbital lobe. *H. convexus* Miers also appears to possess such a structure and many species of *Hyastenus* have on the upper anterior edge of the postorbital lobe a well developed accessory lobe. Sakai (1938:280) has pointed out that the abdomen of the female of *H. diacanthus* (de Haan) consists of only five segments. Such features are usually considered sufficient to warrant generic separation.

6. *Micippa* Leach and *Paramicippa* H. Milne Edwards. When *Paramicippa* was first set up in 1834, two species, *Micippa platipes* Rüppell and *P. tuberculosa* Milne Edwards were included. The first was designated type species of *Paramicippa* by Miers (1879c: 662) and a definition of that genus given at the same time. The second species was alone included in *Paramicippa* in Miers's later revision of the two genera (Miers 1885). Such a procedure was quite invalid. Since *M. platipes* is now generally accepted as belonging to *Micippa* (Sakai 1938, Buitendijk 1939), *Paramicippa* should fall to that genus. If a separate genus is to be established for *P. tuberculosa* because of the pronounced bifid nature of its rostrum or its comparatively long eyestalk or for any other reason, a new name must be used. The situation awaits further investigation and *P. tuberculosa* is here included amongst species of *Micippa*.

Finally, among the genera introduced to the Australian lists in this paper, two require further mention; in one way they may be regarded as merely replacing names already in use.

1. *Scyramathia* A. Milne Edwards. In 1918 Rathbun placed *Hyastenus fultoni* Grant, 1905 in this genus which has recently been recognised as a synonym of *Rochinia* A. Milne Edwards (Garth 1958: 282). Examination of material of Grant's species (11 specimens including relatively large males and females from the Australian Museum's collections: reg. no. P.4515, 20 miles east of Babel I., Tasmania, 65-70 fms., "Endeavour" Expedition) reveals the male first pleopod as being of the "pisoidiform" type (Garth 1958:249; see also his pl. Q) with a truncate but poorly expanded apex and similar to that of *Rochinia occidentalis* (Faxon) (see Garth 1958: pl. Q fig. 7). *H. fultoni* also resembles this species in shape and type of ornamentation of the carapace, relative length of the rostrum, form of the orbit and in several other features (see Rathbun 1925: pls. 228, 229 fig. 5). It seems to fit very satisfactorily into *Rochinia* which genus is now added to the Australian fauna. In the Indo-West-Pacific, species of *Rochinia* are known also from Japan, India and South Africa.

2. *Chorilbinia* Lockington. Garth (1958: 282) has recently orphaned the Australian and New Guinean *C. gracilipes* Miers, 1879 and the Indian *C. andamanica* Alcock, 1895 by the transference, to the mithracine *Stenocionops* Desmarest, 1823, of *Chorilbinia angusta* Lockington, 1877, type species (by monotypy) of *Chorilbinia* Lockington, 1877. The Indian and Australian species are completely different from *C. angusta* and resemble species of *Libinia* Leach and *Libidoclea* H. Milne Edwards & Lucas among the American Pisinae. However, examination of *C. gracilipes* (numerous specimens including relatively large males and females from the Australian Museum's collections: reg. no. P. 14931, Albany Passage, N. Queensland, Melbourne Ward, August 1928) shows that it differs from such American species in two important characters. First, the female abdomen comprises only five free segments (fourth to sixth inclusive fused) in contrast to the seven of *Libinia* and *Libidoclea*. Secondly, the male first pleopod is of the pisoidiform type with a tapered, acute apex and subterminal aperture in contrast to the scyriform type possessed by the American species. *C. andamanica* and *C. gracilipes* resemble each other in structure of the orbit, form of the rostrum, shape of the carapace and comparative length of the ambulatories; *C. andamanica* also has a five-segmented abdomen in the female (Alcock 1895). The form of the male first pleopod

is unfortunately unknown for the Indian species. A new genus thus seems to be required for these two Indo-West-Pacific species. The name *Chlorolibinia* published by Haswell (1882:17) is surely a mistake for Lockington's genus as the latter's name follows the name of the genus. Being an incorrect subsequent spelling, *Chlorolibinia* is unavailable for use under article 33 (b) of the International Code of Zoological Nomenclature. The name *Austrolibinia* is therefore proposed and the genus is diagnosed below.

AUSTROLIBINIA n. gen.

Chorilbinia Lockington; Miers 1879b: 7; (part: *C. gracilipes* Miers, 1879).

Alcock 1895: 221 (part: *C. andamanica* Alcock, 1895).

Chlorolibinia Haswell, 1882c: 17; incorrect subsequent spelling of *Chorilbinia* Lockington, 1877.

Carapace pyriform, armed with a few slender spines and bearing posteriorly a broad, medially acute lobe. Rostrum united in basal half, consisting distally of two acute, divergent spines. Supraorbital eave well expanded, anterolaterally and posterolaterally acute, separated from the large, cupped postorbital lobe by a very narrow fissure; intercalated spine absent. Basal antennal article of moderate width, provided with a prominent lobe laterally at its base. Ambulatory legs long and slender, the first about twice carapace length. Chelipeds shorter than the first ambulatory leg in both sexes, chelae moderately inflated in male. Abdomen of seven segments in the male, of five in the female, fourth to sixth fused. First pleopod of male slender, tapering, apically acute, aperture subterminal (based on *C. gracilipes* only).

Type species: *Chorilbinia gracilipes* Miers, 1879.

V. THE SPECIES

The characters which are of value at the specific level are for the most part different in each genus. For instance, in the genus *Notomithrax*, the form of the crests on the carpus of the cheliped is of importance; in nearly all majines the number and arrangement of the spines on the carapace are a reliable guide but appear to be of little use in many pisines. Several characters which are widely used at the specific level unfortunately differ with age and/or sex. These include the shape of the chela and abdomen which change with growth, often in a single moult, and also differ according to sex; relative proportions of the carapace and length of the spines which change with growth, the carapace becoming wider (particularly in pisines and majines) and the spines shorter and blunter; and the number of free segments in the abdomen which is often different in males and females of the one species. In *Huenia* species the shape of the carapace is strikingly different in males and females. Cases in which sexual dimorphism has resulted in the original description of two species, one based on the male and one on the female, are, as in other groups of animals, not infrequent. The shape of the merus of the third maxilliped and of the basal antennal article are often used diagnostically but in some species the degree of spinulation or tuberculation of these two structures may change during growth.

Among the 45 species listed by Haswell (1882c), 16 have since suffered specific name changes. Some of these have been mentioned in the preceding section. Some species, particularly among the Acanthonychinae, are now recognised as highly polymorphic so that numerous names are reduced to a single valid one. This is true, for example, in the genera *Oncinopus* de Haan, *Menaethius* Latreille and *Huenia* de Haan, although the number of Australian species currently recognised is the same as that listed by Haswell. Specific name changes have taken place, since Haswell's time, particularly in the genera *Naxia* Latreille, *Chlorinoides* Haswell, *Micippa* Leach and *Tiarinia* Dana (see references in key). Special mention should be made of the relatively recent recognition that *Platymaia wyvillethomsoni* Miers is a western Pacific species distinct from the Indian Ocean *P. alcocki* Rathbun (*P. wyvillethomsoni* of Alcock and later authors) (Rathbun 1918); Calman's (1900) Torres Strait material of *Xenocarcinus tuberculatus* is correctly referable to *X. depressus* Miers (Gordon 1934, Sakai 1965), whilst *Xenocarcinus tuberculatus* is a western Pacific species distinct from the Indian Ocean *X. alcocki* Laurie (Sakai 1965); *Paramicippa hispida* Baker (*Eruma hispida* of McCulloch) is synonymous with *Anacinetops*

stimpsoni Miers (Balss 1935); *Schizophyrs dama* (Herbst) is a western species in Australia distinct from the widespread *S. aspera* (H. Milne Edwards) (Balss 1935, Yaldwyn 1964); *Cyclax spinicinctus* Heller (? = *C. perryi* Dana) is distinct from *C. suborbicularis* (Stimpson) (Forest & Guinot 1961); *Micippa platipes* Rüppell and *M. philyra* (Herbst), often considered synonymous, are in fact distinct (Buitendijk 1939); *Notomithrax ursus* (Herbst) includes *Paramithrax latreillei* Miers (Bennett 1964, Griffin 1963a) and definitely occurs in Australia (McNeill 1953); the Australian material of *Paramithrax peronii* mentioned by Haswell (1882c) is correctly referable to *Notomithrax minor* (Filhol) (Bennett 1964); *Leptomithrax australiensis* Miers and *L. spinulosus* Haswell are both synonyms of *L. gaimardi* (H. Milne Edwards) (Griffin 1963b); *Achaeus fissifrons* (Haswell) includes *A. tenuicollis* Miers (Griffin and Yaldwyn 1965); and that *Sargassocarcinus foliatus* Ward also occurs in Japan where it has been known under the older but generically inaccurate name of *Mimulus cristatus* Balss (Sakai 1965). In this paper advantage is taken of the remarks of previous carcinologists to consider *Cancer aragnoides* Rumphius (specific name misspelt *arachnoides* by later workers), *Cancer longipes* Linnaeus, *Egeria indica* Leach and *E. herbstii* H. Milne Edwards a single species, *Phalangipus longipes* (Miers 1884: 182, Alcock 1895: 224 and Grant & McCulloch 1906: 27). Similarly, *Naxia cerastes* Ortmann is considered synonymous with *Naxia taurus* Pocock and included as *Naxioides taurus* (Alcock 1895: 220, Calman 1900: 37). Both these species need investigation.

It is brought out later in this paper that a large number of species considered to be restricted to Australia have been recorded only once from little material. Nothing is to be gained by listing these—material of all is to be eagerly sought. Only two are mentioned here as they are not included in the key.

1. *Stenorhynchus curvirostris* A. Milne Edwards, 1873. This was recorded from Bass Strait and the original description was repeated by Haswell (1882c: 2). I am unable to confidently place it in a genus but it should key out somewhere near *Achaeus* or *Achaeopsis*; Miers (1886: 6, 18) tentatively placed it in *Achaeopsis*. As already indicated (see preceding section) it may belong in *Macropodia* which so far is unknown from Australia.

2. *Inachus australis* Gray, 1831. This was not listed by Haswell. Gray's description was very short and I am unable to comment on the relationships of the species; Gray mentions that it is "somewhat allied" to *Inachus arabicus* Rüppell (a synonym of *Menaethius monoceros* Latreille) whilst Miers (1886: 19) considered that it might belong in *Chlorinoides*.

Special difficulties in satisfactory delineating species appear to have been concentrated in a few genera, most of which still await study. Three require mention here:

1. *Hyastenus*. Many species appear to show rather wide geographic or age variation in ornamentation of the carapace and relative length of the rostral spines. Yet both these characters were heavily relied upon by Alcock (1895) whose arrangement is largely followed here because it is these features which have most often been mentioned in descriptions in the literature. Buitendijk (1939) has illustrated the male first pleopods of several Australian species.

2. *Micippa*. Species of this genus also appear to show the kind of wide variation exhibited in *Hyastenus*. Characters such as the presence of knobs on the spines should be used with caution, as in *Chlorinoides*.

3. *Tiarinia*. Species of this genus are in at least as much confusion as those of any other. Only *T. angusta*, which stands out because of its spinulous rostrum, is easily recognised. The status of all other species is in doubt although descriptions and illustrations provided by Sakai (1938, 1965) and by Buitendijk (1939) are of assistance. The main requirement is definition of the species on the basis of the male first pleopod (as in many other species groups), followed by an analysis of the tuberculation and spinulation of the dorsal surface and margins of the carapace and their changes during growth. Measurements given by Buitendijk of three dimensions of the carapace in *T. cornigera* (Latreille) and *T. gracilis* Dana (of which the Australian and western Pacific *T. depressa* Stimpson may be a synonym) reveal no relative growth differences.

Only four other species need be mentioned.

1. *Entomonyx spinosus* Miers. This quite widespread Indo-West Pacific species, of which *Macrocoeloma nummifer* Alcock is a synonym, has been recorded only once from Australia (as *Acanthophrys spinosus* by Bäss 1929). *Entomonyx* is resurrected as a monotypic genus to accommodate it (Griffin, in press, a).
2. *Criocarcinus superciliatus* (Herbst). This widespread Indo-West Pacific ophthalmine is included in the key on the basis of material from the Great Barrier Reef identified by F. A. McNeill (in prep.).
3. *Achaeus* sp. A single specimen in the Australian Museum's collections appears to constitute an unnamed species of *Achaeus*. Additional characters of this species have been given elsewhere (Griffin & Yaldwyn 1965).
4. *Chlorinoides goldsboroughi* Rathbun. This species, previously known only from Hawaii, is included in the key on the basis of two specimens in the Australian Museum's collections taken off New South Wales. Additional characters for separating this species from the closely related Japanese *C. brevispinosa* Yokoya have recently been given by Sakai (1965:88).

VI. THE KEY

The following key to genera and species includes all species recorded in the literature from Australian localities, with the exception of *Stenorhynchus curvirostris* and *Inachus australis* (see above).

Apart from the characters given for each taxon, the following information is also presented. In the case of genera, the name is followed by the world geographic range; references to important accounts of the genus as a whole, or merely of non-Australian species; and the approximate number of known valid species. In the case of species the name is followed by specific synonyms (as recorded in the Australian literature); size; distribution within Australia; distribution outside Australia; bathymetric range (for details of last four see below); further distinguishing characters or other information which may be of assistance in more positively identifying the species; and finally (in brackets), one or two references to descriptions and illustrations of the species. (Note: Six species are figured in this paper—pls. xv-xvii—and a reference to these is inserted in the appropriate part of the key. In the case of four species, viz., *Huenia bifurcata*, *Hyastenus auctus*, *Hyastenus sebae* and *Tiarinia elegans*, these are the first illustrations provided; that of *Paranaxia serpulifera* is the first illustration of an adult specimen). Where a genus is monotypic or represented in Australia by only a single species, the above information follows on immediately from the remarks concerning the genus.

Size: Relative size (total carapace length including rostrum) is indicated as follows: very small: less than 10 mm; small: 10-20 mm; medium: 20-40 mm; large: 40-80 mm; very large: more than 80 mm.

Australian distribution: The Australian coastline is divided into six regions for the purposes of this review. SE (south east): Tasmania, Victoria and New South Wales, or, more correctly, Kangaroo Island to Brisbane; NE (north east): remaining part of the eastern seabord; N (north): Torres Strait; NW (north west): remaining part of northern coastline to Shark Bay; SW (south west): west coast from Shark Bay and along south coast to Eucla; and S (south): rest of the Great Australian Bight to Spencer's Gulf. The divisions south and north are used here to emphasise the fact that some species are recorded from South Australia but not Victoria, or from Torres Strait but not the adjacent Barrier Reef or Gulf of Carpentaria; they are thus convenient divisions only and may have no real zoogeographic basis (see Section VII). It should be noted that this review treats only species recorded from the Australian mainland, Tasmania, the inshore islands off the western and eastern coastlines, New Guinea and Lord Howe Island.

Overseas Distribution: The grouping and designation of overseas localities in the main follows Ekman (1953). Thus the tropical to warm temperate area of the Indian and western Pacific Oceans are considered to comprise the following faunal subregions (outside Australia): Indian Ocean; Indo-Malaya (including the Philippines); tropical and subtropical Japan (designated simply as Japan); central Pacific (excluding Hawaii); and Hawaii. Taxa, the distribution of which extends through the Indian Ocean and Japan and any other subregion, are designated in the key as Indo-West Pacific.

Bathymetric Distribution: The depth at which species occur is divided into five categories: littoral; intertidal; sublittoral: down to 10 fms; shallow offshore: 10-50 fms; lower shelf: 50-100 fms; and slope: over 100 fms. In the case of species which extend down the slope, the deepest recorded occurrence is given.

Finally, an attempt has been made in the key, at least at the generic level and above, to use as far as possible characters which are considered important from the phylogenetic point of view; in most cases ready separation is achieved. The key is set out with contrasting couplets adjacent so as to permit easier comparison of the divisions.

**KEY TO SUBFAMILIES OF MAJID BRACHYURA AND AUSTRALIAN
GENERA AND SPECIES**

- | | | |
|------|---|-----------------------|
| 1 | Eyes either without orbits, or with incomplete or commencing orbits. Basal antennal article rather slender | 2 |
| — | Eyes with nearly complete, or complete, orbits. Basal antennal article very broad | 61 |
| 2(1) | Male abdomen terminally broadened, seventh segment sub-quadratae and inserted deeply into sixth segment. Male first pleopod longitudinally grooved, with rows of filamentous setae on either side of groove | Subfamily OREGONIINAE |
| — | Male abdomen not terminally broadened, seventh segment sub-triangular and not inserted deeply into sixth segment. Male first pleopod exceedingly varied but not as in Oregoniinae | 3 |
| 3(2) | Eyes without orbits; eyestalks generally long, either non-retractile, or retractile against an acute postorbital spine affording no concealment. Basal antennal article extremely slender and usually long | Subfamily INACHINAE |
| — | Eyes with incomplete or commencing orbits. Basal antennal article not extremely slender | 24 |
| 4(3) | Seven free abdominal segments in male and usually in female | 5 |
| — | Abdomen of six or fewer segments in both sexes | 15 |
| 5(4) | Carapace circular or subcircular. Rostrum appearing trispinose, interantennular spine forwardly projecting and visible from above. Postorbital lobe a prominent spine. Basal antennal article cylindrical. Ambulatory legs very long, the longest more than 3 times carapace length | 6 |
| — | Carapace pyriform or triangular, never circular. Interantennular spine not visible from above. Postorbital lobe spinous or absent. Basal antennal article flattened. Ambulatory legs variously elongate, never more than 2½ times carapace length | 7 |
| 6(5) | Interantennular spine projecting well beyond rostrum. Ambulatory propodi flattened, oar-like. Chelipeds not very long, less than one-third length of ambulatories in adult male, palm in male inflated, widest about midway along. No long protogastric spines <i>Platymata</i> Miers, 1886
Indo-West Pacific, 6: <i>P. wylieithomsoni</i> Miers, 1886; medium; in Australia known only from four localities near Eucla, Great Australian Bight (SW); Indo-Malaya, C. Pacific, Japan; lower shelf and slope to 400 fms; further distinguished by spine on upper anterior margin of postorbital lobe (Rathbun, 1918: 7, pls. III, IV, XIV). | |
| — | Interantennular spine scarcely, if at all, exceeding rostrum. Ambulatory propodi cylindrical. Chelipeds very long, as long as ambulatories in adult male, palm in male widest distally. Protogastric regions each with a very long, forwardly directed spine <i>Cyrtomaia</i> Miers, 1886
Indo-West Pacific, 13: <i>C. maccullochi</i> Rathbun, 1918; medium to large; known only from near Eucla, Great Australian Bight (SW) in 190-450 fms; further distinguished by cylindrical basal segments of antennal flagellum and long rostral spines (Rathbun 1918: 4, figs. 1, 2, pls. 1, 2). | |
| 7(5) | Rostrum very short, less than 1/8 postrostral carapace length, of two rounded or blunt lobes. Carapace smooth | 8 |
| — | Rostrum long, at least 1/5 postrostral carapace length, of two acute spines. Carapace with scattered tubercles | 10 |

- 8(7) Eyestalks small, retractile beneath edge of carapace. Supraorbital eave unexpanded, without lobes or spines. Ambulatory legs slender and moderately long, propodi 1 & 2 dilated and compressed, dactyli of all ambulatories subchelate *Oncinopus* de Haan, 1839.
Widespread Indo-West Pacific; monotypic: *O. aranea* de Haan, 1839 (= *O. subpellucidus* Stimpson, 1857; *O. angulatus* Haswell, 1880); small; S, SE, NE, N; littoral to shallow offshore; carapace weakly calcified (Hale 1927: 125, fig. 122; Sakai 1965: 66, pl. 27, fig. 1). 9
- Eyestalks very long. Supraorbital eave variously expanded. Ambulatory legs variously elongate, cylindrical
- 9(8) Ambulatory legs long, the first twice carapace length, slender. Supraorbital eave unexpanded, lacking lobes or spines. Basal antennal article very slender. Third maxillipeds with merus narrower than ischium *Camposcia* Latreille, 1829
Widespread Indo-West Pacific, monotypic: *C. retusa* Latreille, 1829; medium; NE, N, NW; littoral and sublittoral; often masked by seaweed and sponges (Barnard 1950: 12, fig. 1; Sakai 1965: 69, pl. 30, fig. 1).
- Ambulatory legs short, the first less than $\frac{1}{4}$ carapace length, quite stout. Supraorbital eave moderately expanded with a small antorbital lobe; a prominent conical spine between eave and postorbital lobe separated from both by wide fissures. Basal antennal article moderately broad. Third maxillipeds with merus as wide as ischium *Anacinetops* Miers, 1879
Australia, Timor; monotypic: *A. stimpsoni* (Miers, 1879) (= *Paramicippa hispida* Baker, 1905); medium; S, NE, N; littoral (McCulloch 1913: 336, figs. 47, 48). 10
- 10(7) Abdomen showing no coalescence of segments. Orbita with eave unexpanded and unarmed except for a small, widely removed postorbital spine. Ambulatory legs long, cylindrical, first pair more than twice carapace length, stout . . . *Ephippias* Rathbun, 1918
Restricted to Australia, monotypic: *E. endeavouri* Rathbun, 1918; very large; SE, S, SW, NW; shallow offshore (Rathbun 1918: 9, pl. 15). 11
- Abdomen sometimes, especially in female, showing coalescence of segments. Orbita with moderately expanded eave followed by one or two spines. Ambulatory legs of only moderate length, less than $1\frac{1}{2}$ carapace length, generally slender, propodi distally expanded, compressed, dactyli subchelate . . . *Naxia* Latreille, 1825
Australasia; see reviews by Baker (1905), McCulloch (1913), Balss (1935), Griffin (in press, b); 6; all species (except *N. deflexifrons*) figured by Hale (1927); all except *N. spinosa* restricted to Australia. 12
- 11(10) Ambulatory propodi weakly expanded distally. Supraorbital eave usually acute anterolaterally 12
- Ambulatory propodi strongly expanded distally. Supraorbital eave rounded anterolaterally, armed with a small spine posterolaterally 14
- 12(11) Supraorbital eave rounded anterolaterally and with a small spine posterolaterally. Carpus of cheliped smooth
N. deflexifrons (Haswell, 1880)
Medium; SE; shallow offshore (McCulloch 1913: 330, pl. X figs. 1-4). 13
- Supraorbital eave acute both anterolaterally and posterolaterally. Carpus of cheliped ridged 13

- 13(12) Rostral spines stout, almost straight. Carpus of cheliped with lateral ridge rounded . . . *N. aurita* (Latreille, 1825)
 (= *Halimus laevis* Haswell, 1880); large; SW, S, SE; littoral
 (Hale 1927: 129, fig. 127).
 — Rostral spines slender, strongly curved outwards distally. Carpus of cheliped with lateral ridge proximally acute . . .
N. aires (Guérin, 1825)
 (= *Halimus gracilis* Baker, 1905); medium; S, SE; littoral
 (Hale 1927: 129, fig. 128).
- 14(11) Anterolateral spine of basal antennal article laterally denticulate. Merus of cheliped weakly tuberculate dorsally . . .
N. tumida (Dana, 1851)
 (= *Halimus tumidus* var. *gracilis* Baker, 1905); medium; S, SE, NE; littoral (Hale 1927: 128, fig. 126).
 — Anterolateral spine of basal antennal article unarmed. Merus of cheliped smooth . . . *N. spinosa* (Hess, 1865)
 (= *Halimus truncatipes* Miers, 1879); large; SW, S, SE; Kermadecs; littoral (Hale 1927: 127, fig. 125).
- 15(4) Abdomen of six segments in both sexes. Carapace triangular 16
 — Abdomen of five segments in female, of five or six in male.
 Carapace subpentagonal to subpyriform . . .
Paratymolus Miers, 1879 21
 Indo-West Pacific; See also Sakai (1938); 5.
- 16(15) A prominent postorbital spine. Rostrum generally of moderate length, about 1/3 postrostral carapace length, of two acute spines. Carapace with a few prominent spines . . .
Achaeopsis Stimpson, 1857 17
 Indo-West Pacific, Atlantic, Mediterranean; see Sakai (1938); 6; key to species in Griffin (in press, b).
 — No postorbital spine. Rostrum very short, less than 1/10 postrostral carapace length, of two generally blunt lobes. Carapace with a few tubercles but not spines . . . *Achaeus* Leach, 1817
 Indo-West Pacific, E. Atlantic, Mediterranean; review of Australasian species in Griffin & Yaldwyn (1965); 20. 18
- 17(16) Rostral spines unarmed. Branchial margins with two prominent spines and a few spinules . . .
A. thomsoni (Wyville-Thomson, 1873)
 Small, SE, SW; South Africa, E. & W. Atlantic, Mediterranean; lower shelf and slope down to 1000 fms; further distinguished by spinous merus and carpus of cheliped (Hale 1927: 124, fig. 120).
 — Rostral spines with 2 or 3 spinules laterally. Branchial margins with four prominent spines . . . *A. ramusculus* (Baker, 1906)
 Small; in Australia known only from off Neptune I. (S), in 104 fms; New Zealand (Hale 1927: 124, fig. 121; Griffin in press, b, fig.).
- 18(16) Rostrum of two short, acute spines. Supraorbital eave bearing laterally a prominent spine and several minute spinules . . .
A. fissifrons (Haswell, 1879)
 (= *A. tenuicollis* Miers, 1886); small; SE; New Zealand, Indo-West Pacific; shallow offshore and lower shelf (Griffin & Yaldwyn 1965: 38, figs. 1-8).
 — Rostrum of two generally rounded lobes. Supraorbital eave sometimes with minute spinules but never with a prominent spine 19

- 19(18) Carapace smooth, lacking prominent tubercles. Dactyli of ambulatories 3 and 4 almost semicircular, ventrally spinulated for distal 2/3 *A. lacertosus* Stimpson, 1857 (= *A. breviceps* Haswell, 1880); small; SE, NE, NW; Indo-West Pacific; sublittoral (Stimpson 1907: 20, pl. III fig. 7).
— Carapace with at least two prominent tubercles on cardiac region. Dactyli of ambulatories 3 and 4 strongly falcate but not semicircular, ventrally spinulated for entire length 20
- 20(19) Rostrum of two strong lobes separated by a narrow V or U sinus, often weakly convergent distally. Intestinal region with a small tubercle *A. brevirostris* (Haswell, 1879) (= *A. affinis* Miers, 1884); small; SE, NE, N, NW; Indian O., Indo-Malaya; sublittoral to shallow offshore (Haswell 1880a: 432, pl. XXVII fig. 5).
— Rostrum of two pointed lobes, each with a small, sharp, distal projection. Intestinal region without a tubercle *Achaeus* sp. Known only from a single small female taken off Cairns (NE) in 28 fms. (Griffin & Yaldwyn 1965:38).
- 21(15) Carapace with protogastric regions each bearing a strong tubercle; branchial margin also with a tubercle or spine 22
— Carapace smooth dorsally and without spines posterolaterally on branchial margin 23
- 22(21) Palm of chela dorsally prolonged into a terminal spine. Merus of cheliped with four ventral spines *P. bituberculatus* Haswell, 1880 (= *P. bituberculatus* var. *gracilis* Miers, 1884); very small; restricted to Australia, NE; sublittoral (Haswell 1880c: 303, pl. XVI figs 1, 2).
— Palm of chela not prolonged into a terminal spine. Merus of cheliped unarmed *P. pubescens* Miers, 1879 Very small; N; Indo-West Pacific; sublittoral to shallow offshore (Miers 1879a: 45, pl. 2 figs 6, 6a, b; Sakai 1965: 66, pl. 26 figs. 3, 4).
- 23(21) Palm of chela smooth. Merus of cheliped smooth dorsally and with two tubercles ventrally *P. sexspinosa* Miers, 1884 Very small; NE, N; Indian O., Japan; sublittoral (Miers 1884: 261, pl. XXVII figs., B, b, b').
— Palm of chela dorsally spinulous, outer surface with granulations and spinules. Merus of cheliped with numerous spinules dorsally and a few ventrally *P. latipes* Haswell, 1880 (= *P. latipes* var. *quadridentata* Baker, 1906); small; restricted to Australia, SE, S, SW; sublittoral (Hale 1927: 123, fig. 119).
- 24(3) Eyes without true orbits, lacking a postorbital cup. Eyestalks variously elongate 25
— Eyes with commencing orbits having, in addition to the suprabranchial eave which is sometimes produced into a preorbital spine, a large cupped postorbital process into which the eyestalk retracts. Eyestalks short Subfamily PISINAE 36
- 25(24) Eyestalks long, orbit partially protected by a hornlike supraorbital spine or by a jagged postorbital tooth or by both Subfamily OPHTHALMIINAE 26
— Eyestalks short, little moveable and either concealed by a preorbital spine or sunk in sides of rostrum Subfamily ACANTHONYCHINAE 29

- 26(25) Supraorbital eave hardly expanded and bearing only a small antorbital spine. Rostrum of two weakly curved spines less than 1/3 postrostral carapace length. Branchial margin with small tubercles or unarmed . . . *Zewa* McCulloch, 1913 . . .
Australia, Japan; see McCulloch (1913), Sakai (1938); 4; Australian species, both of which are not known overseas, lack prominent marginal branchial tubercles. 27
- Supraorbital eave prominently expanded with a strongly developed antorbital spine and/or preorbital spine. Rostrum either of two straight spines 2/3 postrostral carapace length or of two exceedingly short and strongly curved spines. Branchial margins with 2-3 long spines 28
- 27(26) Postorbital lobe with a strong tubercle. Posterior intestinal margin rounded *Z. varians* (Miers, 1879)
Medium; SE, NE, N, NW; sublittoral (Calman 1900: 39, pl. 2 figs. 25, 26).
- Postorbital lobe without an anterior tubercle. Posterior intestinal margin with a strong medial tubercle
Z. banfieldi McCulloch, 1913
Medium; known only from near Cairns (NE); littoral (McCulloch 1913: 332, pl. X figs. 5, 6).
- 28(26) Supraorbital eave rounded anteriorly, antorbital spine greatly elongated. Rostral spines widely separated, straight, 2/3 postrostral carapace length. Branchial margins with 3 spines
Picroceros A. Milne Edwards 1865
N. Caledonia, Lord Howe, N. Hebrides, Japan; monotypic:
P. armatus A. Milne Edwards, 1865; very large; NE; sublittoral; further distinguished by single gastric spine and single posterior intestinal spine; carapace spines sometimes knobbed (Sakai 1938: 247, fig. 24; pl. XXV fig. 2).
- Supraorbital eave with strongly developed, subequal preorbital and antorbital spines. Rostral spines arising close together, strongly curved outwards, less than 1/6 postrostral carapace length. Branchial margins with two spines
Criocarcinus H. Milne Edwards, 1834
Indo-West Pacific; monotypic: *C. superciliosus* (Herbst, 1803); large; known in Australia only from Low Isles, Great Barrier Reef (NE) (to be reported elsewhere by F. A. McNeill); littoral (A. Milne Edwards, 1872: 242, pl. 12 fig. 3; Sakai 1938: 251, text-fig. 26).
- 29(25) Rostrum of two spines separate from base. A forwardly produced preorbital spine and a postorbital spine both present. Abdomen in female of 6 or 7 free segments
Rostrum single or only distally bifid. Preorbital spine present or absent, postorbital spine absent. Abdomen in female of only 5 free segments 30
- 30(29) Rostrum short, less than 1/4 postrostral carapace length. Branchial margin with a prominent spine or lobe *Pugettia* Dana, 1851
Indo-West Pacific and E. Pacific to California, 18: *P. mosaica* Whitelegge, 1900; small; restricted to Australia, SE; shallow offshore to lower shelf; further distinguished by strongly granulated carapace and dorsally narrow postorbital lobe (Whitelegge 1900: 141, pl. XXXV figs. 5-7). 31
- Rostrum moderately long, more than 1/3 postrostral carapace length. Branchial margin lacking a lobe or spine
Antilibinia Macleay, 1838
South Africa, Philippines and Australia, 3: *A. lappacea* Rathbun, 1918; small; known only from Great Australian Bight (SW) near Eucla in 200-300 fms; further distinguished by prominent hepatic spine and length of rostral spines and preorbital spine (Rathbun 1918: 12, fig. 3; pl. vii fig. 3).

- 31(29) Supraorbital eave with a strong, forwardly directed preorbital spine. Branchial regions, at least, with prominent lateral expansions. Rostrum slender, abdomen in male of 7 free segments 32
- Orbit with a weakly expanded eave lacking spines. Margins of carapace weakly expanded. Rostrum a stout "beak." Abdomen in male of only 5 free segments
Xenocarcinus White, 1847 35
 Indo-West Pacific, 5; see Gordon (1934), Sakai (1938, 1965); upper orbital margin may be weakly notched once or twice.
- 32(31) Rostrum distally acute or bifid, either downwardly deflexed and depressed, or horizontal, compressed and vertically deep. Carapace smooth or with only a few (1-5) tubercles dorsally. Hepatic margin, at least in female, prominently expanded 33
- Rostrum distally acute, horizontal and depressed, vertically shallow. Carapace with numerous scattered tubercles dorsally. Hepatic margin not prominently expanded in either sex
Menaethius H. Milne Edwards, 1834
 Widespread Indo-West Pacific; monotypic: *M. monoceros* (Latreille, 1825); small; NE, N, NW; littoral (Barnard 1950: 43, figs. 9 g, h; Sakai 1965: 74, pl. 33 fig. 4).
- 33(32) Hepatic expansions well developed in both sexes and separated from branchial expansions by a shallow concavity or closed fissure. Rostrum downwardly deflexed
Sargassocarcinus Ward, 1936
 Australia, Philippines and Japan; see Sakai (1965); 2: *S. cristatus* (Balss, 1924) (= *S. foliatus* Ward, 1936); small; in Australia known only from Lindeman I., Whitsunday Passage (NE); Japan; littoral; further distinguished by sharply carinate and lobate ambulatory legs (Ward 1936: 9, pl. III figs. 4-6; Sakai 1965: 77, text-figs. 11a, b, pl. 34 fig. 3). 34
- Hepatic expansions well developed in female and clearly separated by a wide fissure from branchial expansions, poorly developed in male. Rostrum horizontal
Huenia de Haan, 1839
 Widespread Indo-West Pacific, 3; species show extreme variation in carapace shape, particularly between sexes.
- 34(33) Rostrum distally simple and acute. Branchial expansion rounded, subacute, or sometimes bilobed
H. proteus de Haan, 1839
 Medium; S, NW, N; widespread Indo-West Pacific; littoral, (Hale 1927: 133, fig. 132; Barnard 1950: 41, figs. 9 a-f; Sakai 1965: 75, pl. 34 figs. 1, 2). 34
- Rostrum distally bifid. Branchial expansion bi- or trilobed.
H. bifurcata Streets, 1870
 Medium; restricted to Australia, SE; littoral; distally bifid rostrum appears most reliable character (Haswell 1882c: 8). Figured here pl. XV, figs. d, e.
- 35(31) Carapace uneven, bearing several very prominent tubercles. Rostrum distally notched. Ambulatory legs smooth
X. tuberculatus White, 1847
 Medium; NE; Hong Kong, Japan; littoral and shallow offshore (Sakai 1965: 91, text-fig. 13, pl. 42 fig. 5). 37
- Carapace depressed, bearing a few indistinct granules. Rostrum deeply bifid. Ambulatory legs tuberculate
X. depressus Miers, 1874
 Small; SE, N; Indo-Malaya, N. Caledonia, Japan; littoral (Gordon 1934: 70, fig. 36 a-d). 43
- 36(24) Intercalated spine present 37
 — Intercalated spine absent 43

- 37(36) Abdomen of seven segments in both sexes. Carapace pyriform. Supraorbital eave laterally expanded. Ambulatory legs generally short, seldom more than twice carapace length, usually less 38
- Abdomen of seven segments in male, of five in female. Carapace subtriangular. Supraorbital eave almost unexpanded laterally. Ambulatory legs extremely long and slender, the first three times carapace length . . . *Phalangipus* Latreille, 1825 42
Indo-West Pacific; Rathbun (1916); 6.
- 38(37) Rostrum without an accessory spinule. Carapace variously smooth or granular, tuberculate or spinous. Ambulatory legs little longer than carapace 39
- Rostrum with an accessory spinule not far from tip. Carapace with well defined granules, spinules and spines. First ambulatory legs at least $1\frac{1}{2}$ times carapace length . . .
Naxioides A. Milne Edwards, 1865 41
Indo-West Pacific, 9.
- 39(38) Carapace, broad, smooth or granular. Basal antennal article broad, bearing two strong spines distally, one medial and one lateral
Herbstia H. Milne Edwards, 1834
E. & W. central America, S.E. Atlantic, Mediterranean, Australia; see Garth (1958), Rathbun (1918), 10: *H. crassipes* (A. Milne Edwards, 1873); known only from Bass Strait (SE) in unknown depth; further distinguished by absence of spines on lateral margin of carapace (Haswell 1882: 12, no fig.).
- Carapace narrow, dorsally bearing numerous rounded tubercles. Basal antennal article slender, truncate distally or bearing anterolaterally a lobe or spine 40
- 40(39) Supraorbital eave rounded anterolaterally. Tubercles of carapace arranged in distinct groups. Rostral spines widely separated from base . . . *Eurynome* Leach 1817
Indo-West Pacific, E. Atlantic, Mediterranean; genus reviewed in Griffin (1964), 8-9: *E. granulosa* Baker 1906; very small; known only from "South Australian coast" in 100-104 fms. (Griffin 1965:30, figs. 1-5).
- Supraorbital eave produced into a strong, forwardly directed spine. Tubercles of carapace scattered. Rostral spines fused basally
Tylocarcinus Miers, 1879
Widespread Indo-West Pacific; 2: *T. styx* (Herbst, 1803); small; N, NE; Indo-West Pacific; littoral; further distinguished by spinous ambulatory meri and denticulate dactyli (Sakai 1938: 271, pl. XXXVI fig. 5).
- 41(38) Rostral spines about $\frac{1}{3}$ postrostral length of carapace, divergent from base. Dorsal surface of carapace with numerous tubercles sometimes enlarged into straight spines in mid-dorsal regions . . .
N. taurus (Pocock, 1890)
(? = *N. cerastes* Ortmann, 1894); small; N; Indian Ocean, shallow offshore (Alcock 1895: 219-20; Alcock & Anderson 1897: pl. XXXIII figs. 2-2a, 5-5a).
- Rostral spines as long as postrostral length of carapace, subparallel basally, weakly divergent distally only. Dorsal surface of carapace with numerous large, curved spines . . . *N. robillardii* Miers, 1882
Very large; SE; Mauritius, China S.; shallow offshore (Miers 1882: 339, pl. 20).

- 42(37) Median suborbital tooth much deflexed and bounded by a broad U sinus on both sides. Lobe on first abdominal segment in both sexes arcuate and occupying nearly whole width of segment . . .
P. australiensis Rathbun, 1918
 Small; restricted to Australia, known only from Platypus B. (NE) in 7-9 fms; further distinguished by absence of accessory spinule on rostral spines (Rathbun 1918: 15, pl. VI).
- Median suborbital tooth bounded by a V-sinus on both sides. Lobe on first abdominal segment in both sexes small, almost pointed, occupying much less than whole width of segment . . .
P. longipes (Linnaeus, 1767)
 (= "Egeria Aragnoides" Rumphius, 1705; *E. herbstii* H. Milne Edwards, 1834); small; NE, N; Indian Ocean; shallow offshore; carapace generally prominently spinous (Haswell 1882:12—no recent fig.).
- 43(36) Abdomen of seven segments in both sexes. Rostral spines generally distinct from base. Carapace with slender spines or tubercles. Ambulatory legs generally long, if short then carapace weakly tuberculate 44
- Abdomen in male of seven segments, of five segments in female. Rostral spines basally coalescent. Carapace with stout spines, ambulatory legs short or carapace with slender spines, ambulatory legs long 60
- 44(43) Rostral spines slender, in length at least 1/6 postrostral portion of carapace and usually longer. Supraorbital eave not in close contact with postorbital lobe 45
- Rostral spines stout, in length less than 1/6 postrostral portion of carapace. Supraorbital eave separated from postorbital lobe by an extremely narrow fissure or completely coalesced with it 59
- 45(44) Supraorbital eave rounded anterolaterally and posterolaterally. Carapace completely smooth. *Micippoides* A. Milne Edwards, 1873 Widespread Indo-West Pacific; monotypic: *M. angustifrons* A. Milne Edwards, 1873 (= *Hyastenus andrewsi* Calman, 1909); small; E. New Guinea; littoral; further distinguished by weakly denticulate ambulatory dactyls, body and legs covered by very long hair (Calman 1909: 711, pl. LXII figs. 6, 7).
 — Supraorbital eave at least anterolaterally acute, sometimes produced into a preorbital spine. Carapace with several prominent spines and/or tubercles or else granular 46
- 46(45) Supraorbital eave separated from postorbital lobe by a broad U sinus, posterolaterally unarmed, anterolaterally armed with a strong preorbital spine. Rostral spines fused for a short distance basally. Basal antennal article distally truncate
Rochinia A. Milne Edwards, 1875
 Indo-West Pacific and E. Pacific, Atlantic and Mediterranean; see Garth (1958); 18: *R. fultoni* (Grant, 1905) n. comb.; medium; restricted to SE Australia; lower shelf and slope to 300 fms.; further distinguished by 1 medial gastric, 1 medial cardiac, 1 medial intestinal and 1 lateral branchial spine (Rathbun 1918: 14, pl. V).
 — Supraorbital eave posterolaterally acute, separated from postorbital lobe by a wide V or sometimes U sinus, anterolaterally acute but seldom produced. Rostral spines separated from base. Basal antennal article with an anterolateral spine or lobe
Hyastenus White, 1847 47
- Tropical Indo-Pacific; list of species in Balss (1935): 38.
- 47(46) Denuded carapace with dorsal surface bearing numerous tubercles or spines and erosions 48
- Denuded carapace with a few spines or tubercles, often smoothly polished 49

- 48(47) Rostral spines as long as postrostral portion of carapace. Surface of carapace eroded but indistinctly tuberculate
H. sebae White, 1847
 Small; N, NW; Indian O., Indo-Malaya; sublittoral (Alcock 1895: 213). Figured here pl. XV, figs. b, c.
- Rostral spines about $\frac{1}{2}$ postrostral carapace length. Surface of carapace distinctly tuberculate
H. oryx A. Milne Edwards, 1872
 Medium; NE, N, NW, SW; Indian O., Indo-Malaya; sublittoral to shallow offshore (A. Milne Edwards 1872: 250, pl. XIV fig. 1; Haswell 1882:20).
- 49(47) Rostral spines less than 1/5 postrostral portion of carapace
 — Rostral spines at least 1/3 postrostral carapace length and generally longer 50
- 50(49) Gastric regions with three medial tubercles; a prominent medial tubercle on cardiac region and on posterior intestinal margin; dorsal surface otherwise with several tubercles laterally
H. verrucosipes (Adams & White, 1848)
 Small; in Australia known only from Torres Strait in 15-20 fms; Indo-Malaya (Calman 1900: 36, pl. 2 figs. 23, 24).
 — Gastric region with a single tubercle medially, dorsal surface otherwise with some tubercles laterally but none mid-dorsally 51
- 51(50) Rostral spines outwardly curved distally
H. minimus Rathbun, 1924
 Very small; restricted to Australia, known only from C. Jaubert (NW) in 5-7 fms. (Rathbun 1924: 4, fig. 1).
 — Rostral spines subparallel, weakly curved inwards distally.
H. planasius (Adams & White, 1848)
 Medium; in Australia known only from Pt. Denison (NE); New Guinea, Indian O., Indo-Malaya; shallow offshore (Adams & White, 1848: 9, pl. ii figs. 4, 5; Alcock 1895: 212).
- 52(49) Branchial regions with a spine close to lateral margin, at widest part of carapace, markedly longer than any other spine or tubercle except rostral spines
 — Branchial regions with a small tubercle or spine close to lateral margin, at widest part of carapace, little longer than other tubercles of carapace 53
- 53(52) Gastric regions with two medial tubercles or spines, cardiac region with a central tubercle. Rostral spines sinuous
H. spinosus A. Milne Edwards, 1872
 Large; in Australia known only from Torres Strait in 5-7 fms; South Africa, Fiji (Barnard 1950: 53, fig. 11f).
 — Gastric regions with a single medial tubercle; cardiac region without a tubercle. Rostral spines straight 54
- 54(53) Intestinal region without a tubercle. Lateral branchial spine without a subdorsal tubercle in front of it
H. diacanthus (de Haan, 1839)
 Large; SE, NE, N, NW, SW; widespread Indo-West Pacific sublittoral to shallow offshore; rostrum in female often shorter than in male, lateral branchial spine often greatly reduced, abdomen of female of 5 segments (Alcock 1895: 210; Sakai 1965: 81, pl. 36 fig. 1).
 — Intestinal region with a strong tubercle. Lateral branchial spine with a tubercle above and in front of it
H. auctus Rathbun, 1916
 Small; in Australia known only from near C. Jaubert (NW) in 12-14 fms; Philippines (Rathbun 1916: 543). Figured here, pl. xv, fig. a.

- 55(52) Carapace strongly tuberculate. Rostral spines almost as long as postrostral carapace length *H. brockii* de Man, 1888
Medium; in Australia known only from Torres Strait (N); Indian O., Indo-Malaya; sublittoral (de Man 1888: 221, pl. 7 fig. 1).
— Carapace weakly tuberculate or granular or sometimes smooth. Rostral spines usually $\frac{1}{2}$ postrostral carapace length, sometimes longer 56
- 56(55) Rostral spines $\frac{2}{3}$ postrostral carapace length. A single medial gastric tubercle and several granules laterally *H. borradalei* (Rathbun, 1907)
Medium; in Australia known only from C. Jaubert (NW); Indo-West Pacific; sublittoral to shallow offshore; further distinguished by 5 gastric granules in a transverse row, body covered by a thick tomentum (Rathbun 1911: 251, pl. 20 fig. 5).
— Rostral spines $\frac{1}{2}$ postrostral carapace length. Carapace with gastric and intestinal tubercles medially or else smooth 57
- 57(56) Carapace completely smooth except for lateral branchial tubercle. Supraorbital eave produced as a blunt forwardly directed spine *H. irami* (Laurie, 1906)
Very small; in Australia known only from C. Jaubert (NW); Ceylon; sublittoral (Laurie 1906: 379, pl. 1 figs. 4, 4a).
— Carapace with at least a trace of 1 or 2 medial tubercles in gastric, cardiac and intestinal regions. Supraorbital eave acute anteriorly but not forwardly produced 58
- 58(57) Rostral spines curved slightly outwards distally. Branchial regions with 2-3 small tubercles laterally including that at widest part of carapace *H. convexus* Miers, 1884
Small; NE, N, NW; Indian O., Indo-Malaya; sublittoral to shallow offshore (Miers 1884: 196, pl. XVIII figs. B, b).
— Rostral spines curved slightly inwards distally. Branchial regions completely smooth laterally except for small spine at widest part of carapace *H. espinosus* Borradaile, 1903
Small; in Australia known only from C. Jaubert (NW); Maldives Archipelago; sublittoral (Borradaile 1903: 688, pl. XLVII figs. 4a-d).
- 59(44) Carapace armed with several prominent tubercles and generally a few spines. Supraorbital eave and postorbital lobe separated by a narrow fissure. Ambulatory legs smooth . . *Doclea* Leach, 1814
Tropical Indo-Pacific, 13: *D. profunda* Rathbun, 1918; small; known only from a single ovigerous female taken in Great Australian Bight, south of Eucla (SW), in 250-450 fms; further distinguished by single cardiac and intestinal spine and single hepatic and branchial marginal spines; narrowly pyriform carapace unusual for this genus (Rathbun 1918: 16 pl. VII figs. 1, 2).
— Carapace unevenly and indistinctly tuberculate, lacking spines. Supraorbital eave completely fused with postorbital lobe. Ambulatory legs armed with a few spines and tubercles
Perinia Dana, 1851
Widespread Indo-West Pacific, monotypic: *P. tumida* Dana, 1851 (= *Parathoe rotundata* Miers, 1879); very small; NE; littoral; further distinguished by short, broad, apically inwardly curved rostral spines and denticulate dactyli (Miers 1879b: 16, pl. V fig. 2, 2a; Sakai 1938: 294, fig. 40).

- 60(43) Supraorbital eave well expanded posterolaterally, separated from postorbital lobe by an extremely narrow fissure. Rostrum bifid for distal half only. Carapace armed with a few slender spines Ambulatory legs long, the first at least twice carapace length, slender, smooth *Austrolibinia*, n. gen. India, New Guinea, Australia, 2: *A. gracilipes* (Miers, 1879) n. comb.; small; NE, N, NW; New Guinea; shallow offshore; further distinguished by 2 medial gastric and 1 medial cardiac spine, a broad, acute intestinal lobe and 2 dorsal branchial spines (Miers 1879b: 7, pl. IV figs. 4, 4a).
- Supraorbital eave weakly expanded posterolaterally, separated from postorbital lobe by a wide U sinus. Rostrum less than 1/6 postrostral carapace length, bifid for at least distal 2/3. Carapace armed with numerous, short, very coarse spines. Ambulatory legs little longer than carapace, stout, spinous *Hoplophrys* Henderson, 1893 Tropical Indo-West Pacific; see Alcock (1895), Sakai (1932), Buitendijk (1939); 2: *H. ogilbyi* McCulloch, 1908; small; in Australia known only from Moreton B. (NE); Indo-Malaya (Moluccas, Ceram), ?Japan; sublittoral; further distinguished by simple spine at lateral branchial angle (McCulloch 1908: 51, pl. xii fig. 2, 2a).
- 61(1) Basal antennal article not specially expanded to form a floor to the orbit, which is formed by a supraorbital eave, a postorbital lobe and an intercalated spine between the two Subfamily MAJINAE 62
- Basal antennal article expanded to form a floor to the orbit which is formed above by eave and postorbital lobe; intercalated spine present or absent Subfamily MITHRACINAE 81
- 62(61) Postorbital lobe a simple acute spine more or less isolated from orbit and affording no concealment to cornea of eyestalk 63
- Postorbital lobe cupped, close to orbit and affording some concealment to cornea of retracted eyestalk 66
- 63(62) Rostral spines shorter than width at base, fused for basal 1/3. Postorbital spine no longer than intercalated spine. Basal antennal article slightly narrowed anteriorly, lateral margin notched distally. Carapace very weakly tuberculate *Paramithrax* H. Milne Edwards, 1834 Restricted to Australia, monotypic: *P. barbicornis* (Latreille, 1825) (= *Gonatorhynchus tumidus* Haswell, 1880); medium; SW, S, SE; littoral to shallow offshore; further distinguished by two small marginal branchial tubercles (Griffin 1963b: 137, figs. 7-14).
- Rostral spines longer than their width at base, distinct from base. Postorbital spine distinctly longer than intercalated spine. Basal antennal article of even width throughout, or produced into a lobe anterolaterally, not notched laterally or narrowed distally. Carapace spinous or densely tuberculate 64
- 64(63) Chelipeds in adult male robust, merus tuberculate or spinous, carpus with a dorsal and lateral longitudinal ridge. Eyestalks not especially slender and hardly reaching postorbital spine *Notomithrax* Griffin, 1963 65
- South Pacific to Juan Fernandez; review of species in Griffin (1963a); 5.

- Chelipeds in both sexes slender and unarmed, carpus sub-cylindrical, lacking ridges. Eyestalks long and slender, reaching postorbital spine *Maja* Lamarck, 1801
Tropical Indo-West Pacific, E. Atlantic & Mediterranean, 15: *M. miersii* Walker, 1890; large; in Australia known only from C. Jaubert (NW); Singapore, Japan; sublittoral to shallow offshore; further distinguished by 2 medial and 5 marginal spines (Sakai 1938: 298, pl. XXXVIII fig. 2).
- 65(64) Carapace with both spines and tubercles dorsally. Hepatic margin with two spines, branchial margin with seven spines
N. minor (Filhol, 1885)
(= *Paramithrax peronii*; Haswell, 1882); large; SE, NE; New Zealand; sublittoral to lower shelf; further distinguished by spines of branchial margin being alternately large and small; usually thickly covered by algae, hydroids and sponges (Rathbun 1918: 18, pl. VIII).
- Carapace with numerous low tubercles dorsally. Hepatic and branchial margins each with three spines
N. ursus (Herbst, 1788)
(= *Paramithrax latreillei* Miers, 1876); large; SE; New Zealand; littoral; hepatic and branchial marginal spines decrease in size posteriorly, body thickly hirsute (Griffin in press, b, fig.).
- 66(62) Postorbital lobe sometimes with spines or tubercles on posterior edge but not on anterior upper edge 67
- Postorbital lobe with a prominent accessory spine on upper anterior edge near base 78
- 67(66) Supraorbital eave seldom with a preorbital spine. Rostral spines straight but divergent, seldom more than 1/6 postrostral portion of carapace in length. Carapace densely tuberculate, margins spinous. Chelipeds with merus and carpus variously tuberculate
Leptomithrax Miers, 1876 68
Australasia, Indo-Malaya, Japan; review of Australian and New Zealand species in Griffin (in press, b); 16; all Australian species except *L. tuberculatus* restricted to Australia.
- Supraorbital eave with a prominent preorbital lobe directed outwards or upwards. Rostral spines straight and divergent, or, more usually, outwardly curved distally, in length at least 1/2 postrostral portion of carapace. Carapace with a few scattered long or short spines or lamellae, otherwise smooth. Chelipeds with merus and carpus smooth or with well-developed longitudinal crests *Chlorinoides* Haswell, 1880 73
Indo-West Pacific; reviewed in detail in Griffin in press, a); 12.
- 68(67) Intercalated spine not nearly reaching tip of antorbital spine, almost excluded from outer rim of supraorbital margin by distal approximation of antorbital spine and postorbital lobe. Ambulatory legs hardly longer than carapace 69
- Intercalated spine reaching to tip of antorbital spine, not excluded from rim of supraorbital margin. Ambulatory legs at least 1½ times carapace length 71
- 69(68) Sternum and abdomen smooth in adults and juveniles of both sexes. Carapace weakly tuberculate. Two marginal branchial spines *L. parvispinosus* (Ward, 1933)
Medium; SE, NE; littoral; orbit becomes slightly more open in larger specimens (Ward 1933: 392, pl. XXIII fig. 4).

- Sternum or abdomen, or both, excavated as rimmed pits in adult males and juveniles. Carapace strongly tuberculate. Three or four marginal branchial spines 70
- 70(69) Postorbital lobe acuminate. Four marginal branchial spines *L. tuberculatus* (Whitelegge, 1900)
Medium to large; SE; New Zealand, Kermadecs; shallow offshore to lower shelf; Australian forms with short marginal and long dorsal spines (Whitelegge 1900: 146, pl. XXXIV figs. 1, 2).
— Postorbital lobe truncate distally. Three marginal branchial spines *L. sternocostulatus* (H. Milne Edwards, 1851)
Medium to large; S, SE, NE, NW; shallow offshore; further distinguished by sternal excavations being wholly segmental (Hale 1927: 137, fig. 137).
- 71(68) Carapace narrowly pyriform, width no more than $\frac{1}{3}$ postrostral length. Postorbital lobe distally slender with a single small spinule close to tip and numerous small tubercles around base. First ambulatory leg twice carapace length *L. globifer* Rathbun, 1918
Large to very large; SW; lower shelf and slope to 120 fms. (Rathbun 1918: 23, pls. X, XI).
— Carapace broadly pyriform, at least in adult, width at least 2/3 postrostral length. Postorbital lobe subtriangular, distal portion not especially slender, a prominent spine or tubercle close to tip and another about halfway along posterior edge. First ambulatory leg no more than $\frac{1}{4}$ times carapace length 72
- 72(71) Mid-dorsal regions of carapace thickly covered by both spines and tubercles. Ambulatory meri with a blunt terminal dorsal lobe *L. gaimardi* (H. Milne Edwards, 1834)
(*= L. australiensis* Miers, 1876; *L. spinulosus* Haswell, 1880); very large; SW, S, SE; sublittoral to slope down to 450 fms; postorbital lobe with sharp spinules in juvenile and blunt tubercles in adult (Hale 1927: 135, fig. 135: Griffin 1963b: 133, figs. 1-6, pls. 6, 7).
— Mid-dorsal regions of carapace smooth except for a few short, prominent spines. Ambulatory meri with a prominent sharp terminal dorsal spine *L. waitei* (Whitelegge, 1900). Very large; SE; shallow offshore to lower shelf; postorbital lobe with sharp accessory spines in juveniles and adults (Whitelegge 1900: 143, pl. XXXIII).
- 73(67) Preorbital lobe vertically directed upwards from base. Carapace with a few long spines but no lamellae. Rostral spines with a small spinule on dorsal surface near tip *C. tenuirostris* Haswell, 1880
Medium; restricted to Australia; NE, N; shallow offshore (Griffin in press, a, fig.).
— Preorbital lobe outwardly directed, at least basally. Carapace with a few spines and some lamellae, particularly around margins and above orbit. Rostral spines without a spinule on dorsal surface 74
- 74(73) Rostral spines straight, divergent, distal width less than twice basal width. Branchial margins with three spines and, anteriorly, a small lamella. Ambulatory legs with numerous prominent tubercles dorsally arranged in rows *C. goldsboroughi* Rathbun, 1906
Medium; in Australia known from 2 specimens taken off N.S.W. coast in 70-120 fms; Hawaii (Rathbun 1906: 881, pl. XIV fig. 7).
— Rostral spines weakly or strongly outwardly curved, very divergent, distal width at least $2\frac{1}{2}$ times basal width. Branchial margins with two spines posteriorly. Ambulatory legs smooth or minutely spinous 75

- 75(74) Antorbital lobe a narrow, flattened, distally rounded lamella, preorbital lobe much wider and somewhat longer. Rostral spines weakly curved 76
- Antorbital and preorbital lobes subequal, acuminate. Rostral spines very strongly curved outwards distally 77
- 76(75) Preorbital lobe simple, acuminate. Rostral spines unarmed. Posterior intestinal margin with a short, cylindrical spine
C. albanyensis (Ward, 1933)
 Medium; restricted to Australia, NE; sublittoral to shallow offshore (Ward 1933: 391, pl. XXIII fig. 3).
- Preorbital lobe wide distally, truncate or bifid. Rostral spines armed with several strong spines medially. Posterior intestinal margin with a wide, flattened lamella
C. spatulifer (Haswell, 1882)
 (= *Acanthophrys aculeatus* A. Milne Edwards, 1865); medium; restricted to Australia, SW, S, SE; sublittoral to slope down to 250 fms (Hale 1927: 137, fig. 138).
- 77(75) Preorbital spine simple. A single cardiac spine. Intestinal region with two medial spines . . . *C. aculeatus* (H. Milne Edwards, 1834)
 (= *Acanthophrys aculeatus* var. *armatus* Miers, 1884); large; NE, N, NW; Indian O., Japan; sublittoral to lower shelf (Miers 1884: 193, pl. XVIII, fig. A).
- Preorbital lobe divided into two distinct spines. Cardiac region with a pair of widely divergent, outwardly curved spines. Intestinal region with a single spine
C. longispinus (de Haan, 1839)
 (= *Paramithrax coppereri* Haswell, 1880); large; SE, NE, NW; widespread Indo-West Pacific; shallow offshore (Sakai 1965: 87, pl. 40 fig. 1).
- 78(66) Carapace suborbicular. Rostrum exceedingly short, 1/8-1/20 postrostral carapace length, unarmed . . . *Cyclax* Dana, 1851 79
 Widespread Indo-Pacific; detailed account in Forest & Guinot (1961); 2.
- Carapace pyriform. Rostrum of moderate length, more than 1/5 postrostral carapace length, bearing one or two spines or tubercles laterally near base . . . *Schizophrys* White, 1847 80
 Widespread Indo-West Pacific; 2.
- 79(78) Intercalated spine distally with three subequal spinules. Basal antennal article with a strong accessory spine between antero-lateral and anteromedial spines. Marginal spines of carapace granular almost to tip . . . *C. suborbicularis* (Stimpson, 1858)
 (= *Cyclomaia margaritata* A. Milne Edwards, 1872); SW, NW; Seychelles to Tahiti; littoral (Forest & Guinot 1961: 15, figs. 5, 6, 10; pl. VI figs. 1, 2).
- Intercalated spine triangular, granulate basally only. Basal antennal article lacking an accessory spine between main anterior spines. Marginal spines of carapace granular only at their bases
C. spinicinctus Heller, 1861
 Medium; SW, N, NE; E. Africa to Samoa; littoral (Forest & Guinot 1961: 15, figs. 7, 8, 11; pl. VI, fig. 3).
- 80(78) Rostrum with a single lateral spine. Surface of carapace bearing a few groups of prominent tubercles, especially posteriorly and several spines . . . *S. aspera* (H. Milne Edwards, 1834)
 Large; S, NE, N, NW, SW; widespread Indo-West Pacific; littoral (Hale 1927: 134, fig. 139). Figured here pl. xvi, figs. a, b.
- Rostrum with two lateral spines or tubercles. Surface of carapace densely covered by low tubercles dorsally
S. dama (Herbst, 1804)
 Large; SW, NW; Indo-Malaya; littoral (Yaldwyn, 1964, fig.).

81(61)	Intercalated spine present	82
—	Intercalated spine absent (possibly present in <i>Paranaxia</i>)	88
82(81)	Rostrum weakly deflexed, of two slender spines distinct from base Indian O., Japan; monotypic: <i>E. spinosus</i> Miers, 1884 (= <i>Acanthophrys spinosus</i> ; Balss, 1929); small; in Australia known only from Dampier I. (NW); shallow offshore and lower shelf; further distinguished by two marginal branchial spines and densely tuberculate chelipeds (Miers 1884:526, pl. XLVII fig. B; Sakai 1965: 88, pl. 40 fig. 2).	
—	Rostrum generally steeply deflexed, broad, lamellar, of two spines fused for at least basal third <i>Micippa</i> Leach, 1817 Tropical Indo-Pacific; see Miers (1885), Sakai (1938), Buitendijk (1939); 8.	83
83(82)	Eyestalks projecting laterally well beyond postorbital lobe. Rostrum of two distally distinct, truncate lobes <i>M. tuberculosa</i> (H. Milne Edwards, 1834) (= <i>Micippa parvirostris</i> Miers, 1879); small; restricted to Australia, S, SE; littoral; further distinguished by 4 large marginal branchial spines and broad antennal flagellum (Hale 1927: 140, fig. 142).	
—	Eyestalks reaching only to postorbital lobe. Rostrum of two acute spines, or distally notched	84
84(83)	Rostrum terminating in two strong submedial lobes flanked by a short, broad, recurved spine. Carapace strongly tuberculate and lacking spines dorsally	85
—	Rostrum distally bifid or notched, without lateral spines. Carapace smooth or granular with a few tubercles or strongly tuberculate and spinous	86
85(84)	Orbit open below, a wide hiatus between smooth basal antennal article and postorbital lobe. Anterolateral borders of carapace with 8-10 spines, anterior spines broad, posterior spines acuminate <i>M. platipes</i> Rüppell, 1830 (= <i>M. spatulifrons</i> A. Milne Edwards, 1872); small to medium; NE; widespread Indo-West Pacific; littoral and sublittoral (Sakai 1938: 316, fig. 46; pl. XXXII fig. 2, pl. xxxviii fig. 4; Buitendijk 1939: 254, text-fig. 22, pl. X figs. 2, 4).	
—	Orbit closed below, strongly tuberculate basal antennal article in broad contact with postorbital lobe. Anterolateral borders with 3-6 acuminate spines or spinules <i>M. philyra</i> (Herbst, 1803) (= <i>M. superciliosa</i> Haswell, 1880; <i>Paramicippa asperimanus</i> Miers, 1884; <i>M. mascarenica</i> var. <i>nodulifera</i> Baker, 1905); medium; SW, NW, N, S; widespread Indo-West Pacific; littoral and sublittoral (Haswell 1880a: 446, pl. 26 figs. 2, 2a; Buitendijk 1939: 253, text-fig. 21, pl. X figs. 1, 3; Sakai 1965: 90, pl. 42 fig. 1).	
86(84)	Rostrum of two, outwardly curved, sharply pointed spines, distinct for distal half <i>M. thalia</i> (Herbst, 1803) (= <i>M. inermis</i> Haswell, 1880); SW, N, NE; widespread Indo-West Pacific; medium; sublittoral to shallow offshore; further distinguished by 2 medial gastric, 2 submedial cardiac, 1 dorsal branchial and 9 marginal branchial spines (Haswell 1880a: 445 pl. 26 figs. 3, 3a; Sakai 1965: 90, pl. 42 fig. 3).	
—	Rostral spines fused throughout their length	87

- 87(86) Carapace minutely granular dorsally with a few tubercles, anterolateral margins with about 3 small spinules. Merus of cheliped dorsally carinate; ambulatory legs tuberculate
M. curtispina Haswell, 1880
 Small; NE, N; Singapore; sublittoral; further distinguished by basally vertically deflexed and apically inflexed rostrum (Haswell 1880a: 446, pl. 25 figs. 1, 1a).
- Carapace strongly tuberculate and spinous, anterolateral margins with about 9 prominent spines of various sizes. Merus of chelipeds smooth; ambulatory meri smooth except for a terminal dorsal spine
M. spinosa (Stimpson, 1857)
 (= *Paramicippa affinis* Miers, 1879); medium; restricted to Australia, S, SE; sublittoral and shallow offshore (Hale 1927: 140, fig. 143).
- 88(81) Supraorbital eave separated from postorbital lobe by a wide fissure, not expanded anteriorly into a preorbital spine or lobe. Rostral spines short, broad
Tumulosternum McCulloch, 1913
 Restricted to Australia; monotypic: *T. longimanus* (Haswell, 1880); medium; SE; littoral (Haswell 1880a: 444, pl. XXVI; McCulloch 1913: 334, fig. 45).
- Supraorbital eave separated from postorbital lobe by a very narrow fissure or completely unexpanded posterolaterally, armed with a prominent preorbital spine or lobe. Rostral spines moderately to very long
- 89(88) Rostrum of two distinct, subparallel spines, each apically bifid. Basal antennal article narrowed anteriorly. Carapace weakly tuberculate. Ambulatory legs smooth
Paranaxia Rathbun, 1924
 Restricted to Australia; monotypic: *P. serpulifera* (Guérin, 1829); very large; SW, NW, N; sublittoral to shallow offshore Rathbun 1924: 7, Montgomery 1931:417). Figured here, pl. xvii, fig. a.
- Rostrum of two spines, contiguous throughout their length or at most only apically divergent. Basal antennal article with anterolateral angle forwardly produced. Carapace with groups of distinct or confluent tubercles dorsally. Ambulatory legs tuberculate
Tiarinia Dana, 1851
 Tropical Indo-West Pacific; see Stimpson (1907), Sakai (1938), Buitendijk (1939); 6.
- 90(89) Rostrum with two or three lateral spines close to base. Carapace bearing numerous coarse tubercles dorsally
T. angusta Dana, 1851
 (= *T. spinosirostris* Haswell, 1882); medium; NE, N; Indo-Malaya, Japan; littoral; further distinguished by 3 small, curved, marginal branchial spines (Sakai 1936: 160, fig. 4).
- Rostrum unarmed. Carapace smooth, uneven, or with distant tubercles
- 91(90) Rostrum moderately long, about 1/3 postrostral portion of carapace
T. elegans Haswell, 1882
 Medium; known only from off Broughton I., Pt. Stephens (SE) in 25 fms; further distinguished by 2-3 conical submarginal branchial tubercles (Haswell 1882c: 29). Figured here, pl. xvii, fig. b.
- Rostrum short, less than 1/5 postrostral portion of carapace

89

90

91

92

- 92(91) Carapace with distinct erect tubercles and granules dorsally, branchial region with six obtuse tubercles laterally . . .
T. cornigera (Latrelle, 1825)
 (= *T. mammillata* Haswell, 1880); medium to large; NW; Indian O., Indo-Malaya, Japan; littoral; carapace very wide—width equal to postrostral length (Sakai 1965: 91, pl. 42 fig. 2; Buitendijk 1939: pl. XI fig. 1).
- Carapace with confluent depressed tubercles or smooth dorsally, branchial regions with five, or fewer, acute tubercles laterally . . . 93
- 93(92) Branchial regions with five large tubercles laterally. Preorbital spine quite stout. Seventh abdominal segment in male wider than long
T. gracilis Dana, 1852
 (= ?*T. depressa* Stimpson, 1857); medium; SE, NE; Indo-Malaya, Japan; littoral; further distinguished by 3 large, subequal, blunt tubercles on posterior margin (Stimpson 1907: 12, pl. III fig. 2; Sakai 1938: 321, fig. 49; Buitendijk 1939: 259, text-fig. 26, pl. XI fig. 2).
- Branchial regions with a single sharp spine at widest part of carapace laterally. Preorbital spine very slender. Seventh abdominal segment in male much longer than wide
T. tiarata (Adams & White, 1848)
 Small; N. Guinea; West Pacific; sublittoral; further distinguished by completely straight rostral spines (Sakai 1938: 322, pl. XXXVIII fig. 7).


VII. ZOOGEOGRAPHY

Two features of the Australian majid fauna stand out—(1) the relatively large proportion of species (31%) which have been recorded only once from a single locality in Australia; and (2) the very clear partitioning of the fauna into tropical and temperate components.

Of the poorly known species, 12 are not known outside Australia whilst 16 are very widely distributed species known from several parts of the Indo-West Pacific. Ten of the species were recorded by Rathbun, either from the "Endeavour" collections or from near Cape Jaubert (Rathbun 1918, 1924), two were recorded by Whitelegge (1900) and two by Baker (1905, 1906). Fewer species have been recorded from Western Australia than from eastern coasts.

The geographical boundaries between the tropical and temperate faunas are in the form of very broad transition areas (containing a mixture of tropical and temperate species) extending from Shark Bay to Fremantle on the west coast and from about Mast Head Island to Cape Howe on the east. Slightly narrower transition areas between south-eastern and south-western provinces on the one hand and between north-eastern and north-western provinces on the other, appear to exist just west of Kangaroo Island in the south and around Torres Strait in the north. The four faunal provinces which emerge here are in general agreement with the findings of workers on other groups of marine animals (see Bennett & Pope 1953). Two points should be mentioned here—(1) the Spencer Gulf area of South Australia shows a very close affinity with the rest of south-eastern Australia but possesses five species which are tropical and not otherwise known from temperate latitudes (*Oncinopus aranea*, *Anacinetops stimpsoni*, *Huenia proteus*, *Schizophrys aspera* and *Micippa philyra*); (2) the Torres Strait fauna is a mixture of species otherwise known from north-eastern and to a lesser extent north-western, Australia with very few species confined to the region.

The tropical fauna is clearly part of that of the Indo-West Pacific region and possesses few species restricted to Australia. The first feature is borne



out most strikingly by the fact that a larger proportion of it is shared with other Indo-West Pacific areas than with temperate Australian provinces. Thus, of the north-eastern species only 18% extend southward compared with 50% which are shared with the Indian Ocean, 47% with Japan and 36% with Indo-Malaya. Similarly, for the north-western fauna, 28% extend southward whereas 72% are shared with the Indian Ocean, 48% with Japan and 60% with Indo-Malaya. However, the proportion of species which are distributed throughout the Australian tropical area is not very high. For example 42% of the species found in north-eastern Australia are shared with north-western Australia and the proportion falls to 29% if Torres Strait is excluded. Indian Ocean species are represented to approximately equal extents (about 60%) in both the north east and north west but the Japanese species are definitely best represented in the north east (also about 60%). Widespread Indo-West Pacific species which are also widely distributed in the Australian tropics include *Oncinopus aranea*, *Menaethius monoceros*, several species of *Hyastenus*, *Schizophrys aspera*, two species of *Chlorinoides*, three of *Micippa* and two of *Tiarinia*.

The low degree of restriction in the tropical fauna is evidenced by fewer than 30% of the species in either of the tropical provinces which are not found outside Australia and about 30% which are restricted to any one province. Restricted Australian species found in the tropics include species of *Zewa*, *Hyastenus minimus*, *Phalangipus australiensis* and two species of *Chlorinoides*.

The temperate provinces contain fewer species and overall there is a much greater restriction of these species both to Australia and to particular provinces. Thus, only nine species are found in both south-western and south-eastern Australia out of a total of 34 south-eastern and 20 south-western species; 40% of the south-eastern and 50% of the south-western species are not known outside Australia. Eight species appear to be widespread temperate forms (two species of *Naxia*, *Ephippias endeavouri*, *Paratymolus latipes*, two species of *Leptomithrax* and *Chlorinoides spatulifer*). The relationships of the temperate species are either with tropical Australia (e.g., species of *Paratymolus*, *Zewa*, *Huenia*, *Xenocarcinus*, *Hyastenus*, *Chlorinoides* and *Micippa*) or with the Indo-West Pacific (e.g. species of *Achaeopsis*, *Platymaia*, *Cyrtomaia*, *Pugettia*, *Doclea*, *Eurynome* and *Leptomithrax*). The relationships with temperate regions outside Australia are slight. Only one species which does not have a tropical distribution, *Achaeopsis thomsoni*, is shared with South Africa. Five species, all of which are found in south-eastern Australia, are shared with New Zealand. Of the nine genera shared with New Zealand, seven are more or less widespread in the Indo-West Pacific. The strong restricted element in the Australian temperate fauna is exemplified by species of *Naxia*, *Ephippias endeavouri*, *Paramithrax barbicornis* and *Tumulosternum longimanus*. Of the 37 species restricted to Australia, 69% are temperate.

If only species which penetrate the transition zones between tropical and temperate regions are considered, there is indeed only a very slight partitioning of the fauna into eastern and western elements. Such eastern species probably number no more than six (e.g. *Naxia tumida* and *Notomithrax minor*) and western ones no more than five (e.g., *Paranaxia serpulifera* and *Schizophrys dama*). If more stenothermal species are considered a division between eastern and western regions is quite clear.

Distant relationships of the fauna are shown, at the specific level, by nine species shared with South Africa, 10 with the Red Sea and eight with Hawaii; one species, *Achaeopsis thomsoni*, is also found in the Atlantic. The vast majority of these widespread species are represented in the tropical fauna of Australia.

Figure 3. Known Australian distribution of eight species of majid spider crab. (Each circle represents a single recorded locality).

The zoogeographical features shown by the species are strongly emphasised at the generic level. Thus 58% of the 43 genera have widespread tropical Indo-West Pacific representation. A further 16%, also represented in the Indo-West Pacific, have wider relationships, four genera (*Achaeus*, *Achaeopsis*, *Eury nome* and *Maja*) being found in the Atlantic and three (*Pugettia*, *Herbstia* and *Rochinia*) in the eastern Pacific. An additional 7% (three genera—*Picroceros*, *Sargassocarcinus* and *Leptomithrax*) are western Pacific only. Two genera, *Naxia* and *Notomithrax*, both temperate, are mainly Australasian. Five genera (12%) (*Anacinetops*, *Epippia*, *Paramithrax*, *Tumulosternum* and *Paranaxia*) are restricted to Australia; all are monotypic and part of the temperate fauna. Evidence of bipolarity or antitropicality in the Australian majid fauna is, I think, difficult to discern, although the genera *Cyrtomaia*, *Zewa*, *Xenocarcinus*, *Pugettia*, *Sargassocarcinus*, *Eury nome*, *Rochinia*, *Hoplophrys* and *Leptomithrax* may provide some evidence of such distribution patterns.

Of the other Australian crabs the two best known families are the Portunidae (Stephenson 1962) and the Grapsidae (Tesch 1918, Banerjee 1960, Campbell and Griffin, 1966). In both these families the existence of a few more or less widespread temperate species, such as *Leptograpsus variegatus* (Fabricius), *Plagusia capensis* de Haan, *Macropipus corrugatus* (Pennant), *Ovalipes punctatus* (de Haan) (last two also bipolar) and *Nectocarcinus integrifrons* (Latreille), should not be allowed to obscure the fact that the temperate representatives as a whole relate themselves most closely to the tropical faunas. These two families are proportionally less represented in temperate regions of Australia than are the Majidae.

In conclusion it can be stated that the features shown by the Australian Majidae are those shown by other Australian marine animals (see Ekman 1953). This is particularly true of the partitioning into a tropical fauna with tropical Indo-West Pacific relationships and a temperate fauna related to the tropical fauna of Australia rather than to temperate faunas outside Australia. There is also agreement in the apparent position of transition areas and the geographical extent of the faunal provinces; only the Peronian province (central eastern Australia) (see Bennett and Pope 1953) may be of doubtful validity for the majids.

VIII. ACKNOWLEDGMENTS

I wish to thank first the Trustees and Director (Dr. J. W. Evans) of the Australian Museum, Sydney, for providing a grant which allowed me to examine much of the Museum's collections of spider crabs in January 1963. I wish similarly to acknowledge the financing of those studies by a Research Grant from the University of Tasmania. I am grateful also to Drs. Danièle Guinot and J. Forest (Muséum National d'Histoire Naturelle) and T. Sakai (Yokohama National University) for sending literature, Drs. Isabella Gordon (British Museum (Natural History)) and R. B. Manning (U.S. National Museum) for providing photographs of type material in their care, Messrs. Frank A. McNeill, Melbourne Ward (Australian Museum) and Bruce Campbell (Queensland Museum) for discussion and aid with literature, and Messrs. Athol Beswick and Anthony Healy for photography. The manuscript was read by Drs. John S. Garth (Allan Hancock Foundation), R. W. George (Western Australian Museum) and J. C. Yaldwyn (Australian Museum), all of whom provided most helpful comments for which I am very grateful; this review has also been considerably aided by Dr. Yaldwyn's kindness in forwarding material and literature.

IX. REFERENCES

- Adams, A. and White, A. (1848). Crustacea. In A. Adams (ed.), *Zoology of the Voyage of H.M.S. "Samarang" under the command of Captain Sir Edward Belcher, during the years 1843-46*. London. Pp. viii, 66, 13 pls.
- Alcock, A. W. (1895). Materials for a Carcinological Fauna of India. 1. The Brachyura Oxyrhyncha. *J. Asiatic Soc. Beng.* 64: 157-291 pls. 3-5.
- Alcock, A. W. and Anderson, A. R. (1897). Illustrations of the Zoology of H.M. Indian Marine Survey Steamer "Investigator." Crustacea Decapoda. Pls. 33-35. Calcutta.
- Baker, H. W. (1905). Notes on South Australian Decapod Crustacea. Part 2. *Trans. roy. Soc. S. Aust.* 29: 116-131, pls. XXI-XXIV.
- (1906). Notes on South Australian Decapod Crustacea. Part 4. *Trans. roy. Soc. S. Aust.* 30: 104-117, pls. I-III.
- Balss, H. (1929). Decapoden des Roten Meeres. IV. Oxyrhyncha und Schlussbetrachtungen. *Denkschr. Akad. Wiss. Wein. Math-nat. Kl.* 102: 1-30, text-figs. 1-9, pl. 1.
- (1935). Brachyura of the Hamburg Museum Expedition to South-western Australia, 1905. *J. roy. Soc. W. Aust.* 21: 113-151, figs. 1-5, pl. 13.
- (1957). Abteilung Brachyura, Borradaile, 1907. In H. G. Bronns, *Klassen und Ordnungen des Tierreichs* 5 (1) 7, Decapoda 12: 1505-1672, figs. 1131-1199.
- Banerjee, S. K. (1960). Biological Results of the Snellius Expedition. XVIII. The genera *Grapsus*, *Geograpsus* and *Metopograpsus* (Crustacea, Brachyura). *Temminckia* 10: 132-199, text-figs. 1-6.
- Barnard, K. H. (1950). Descriptive Catalogue of South African Decapod Crustacea (Crabs and Shrimps). *Ann. S. Afr. Mus.* 38: 1-837, figs. 1-154.
- Bennett, I. and Pope E. C. (1953). Intertidal Zonation of the Exposed Rocky Shores of Victoria, together with a rearrangement of the Biogeographical Provinces of Temperate Australian Shores. *Aust. J. Mar. Freshw. Res.* 4(1): 105-159, figs. 1-5, pls. 1-6.
- Borradaile, L. A. (1903). The Spider Crabs (Oxyrhyncha). In J. S. Gardiner, *The Fauna and Geography of the Maldives and Laccadive Archipelagos* 2 (10): 681-690, figs. 122-124, pl. XLVII.
- Buitendijk, A. M. (1939). Biological Results of the Snellius Expedition. V. The Dromiacea, Oystomatidae and Oxyrhyncha of the Snellius Expedition. *Temminckia* 4: 223-276, text-figs. 1-27, pls. VII-XI.
- Calman, W. T. (1900). On a Collection of Brachyura from Torres Straits. *Trans. Linn. Soc. Lond. (Zool.)* (ser. 2) 8: 1-50, pls. 1-3.
- (1909). On Decapod Crustacea from Christmas Island, collected by Dr. C. W. Andrews. *Proc. zool. Soc. Lond.* 1909: 703-713, pl. 72.
- (1913). Note on the Brachyuran Genera *Micippoides* and *Hyastenus*. *Ann. Mag. nat. Hist.* (ser. 8) 11: 312-314.
- Campbell, B. M. and Griffin, D. J. G. (1966). The Australian Sesarmae (Crustacea: Brachyura): Genera *Helice*, *Helograpsus* nov., *Cyclograpsus* and *Paragrapsus*. *Mem. Queensl. Mus.* 14 (5): 127-174, 10 figs., pls. xx-xxiii.
- Chhapgar, B. F. (1957). On the marine crabs (Decapoda: Brachyura) of Bombay State. *J. Bombay nat. Hist. Soc.* 54: 399-439, 503-549, 3 figs., 16 pls.
- Dana, J. D. (1851). On the Classification of the Maioid Crustacea or Oxyrhyncha. *Amer. J. Sci.* (ser. 2) 11: 425-434.
- (1852). Crustacea. In "U.S. Exploring Expedition during the years 1838-1842 under the command of Charles Wilkes, U.S.N." 13. Philadelphia: C. Sherman. Pp. viii, 1618. *Atlas* (1855): 27 pp., 96 pls.

- Ekman, S. (1953). Zoogeography of the Sea. London: Sidgwick & Jackson. 417 pp., 121 figs.
- Forest, J. and Guinot D. (1961). Crustacés Décapodes Brachyures de Tahiti et des Tuamotu. In Expédition française sur les récifs coralliens de la Nouvelle-Calédonie. Vol. prélim: i-xi, 1-195, 178 figs. 7 maps, 18 pls.
- Fulton, S. W. and Grant, F. E. (1906a). Some little known Victorian Decapod Crustacea, with Descriptions of New Species. No. III Proc. roy. Soc. Vict. 19 (1): 5-15, pls. III-V.
- (1906b). Census of the Victorian Decapod Crustacea. Part 1. Brachyura. Proc. roy. Soc. Vict. 19 (1): 16-20.
- Garth, J. S. (1958). Brachyura of the Pacific Coast of America: Oxyrhyncha. Allan Hancock Pacif. Exped. 21: 1-854, 106 tables, 9 text-figs., 82 pls.
- Garth, J. S. and Holthuis, L. B. (1963). *Stenorhynchus* Lamarck, 1818 (Crustacea, Decapoda): Proposed Validation under the Plenary Powers with Designation of *Cancer seticornis* Herbst, 1788, as Type Species. Z.N. (S.) 751. Bull. zool. Nomencl. 20(6): 424-428.
- Glaessner, M. F. (1960). The Fossil Decapod Crustacea of New Zealand and the Evolution of the Order Decapoda. N.Z. geol. Surv. paleont. Bull. 31: 1-63, figs. 1-25, pls. 1-7.
- Gordon, I. (1934). Crustacea Brachyura. Résultats Scientifiques du Voyage aux Indes Orientales Néerlandaises de LL. AA. RR. le Prince et la Princesse Léopold de Belgique. Mem. Mus. roy. Hist. nat. Belg. Hors. ser., III (15) (15): 1-78, figs. 1-37.
- Grant, F. E. (1905). Crustacea dredged off Port Jackson in Deep Water. Proc. Linn. Soc. N.S.W. 30 (2): 313-324, pl. x, xi.
- Grant, F. E. and McCulloch, A. R. (1906). On a Collection of Crustacea from the Port Curtis District, Queensland. Proc. Linn. Soc. N.S.W. 31 (1): 1-53, 4 pls.
- Gray, J. E. (1831). Zoological Miscellany, no. 1 (February 1831): 40. London.
- Griffin, D. J. G. (1963a). *Notomithrax* gen. nov. and the Status of the Genus *Paramithrax* H. Milne Edwards (Crustacea, Brachyura, Majidae). Trans. roy. Soc. N.Z. Zool. 3 (22): 229-237.
- (1963b). Redescriptions of the Australian majid spider crabs *Leptomithrax gaimardi* (H. Milne Edwards) and *Paramithrax barbicornis* (Latreille). Rec. Aust. Mus. 26 (4): 131-143, figs. 1-14, pls. 6, 7.
- (1964). A Review of the Genus *Eurynome* Leach (Decapoda, Majidae) and a New Species from New Zealand. Crustaceana 6 (3): 195-206, figs. 1-11.
- (1965). A Redescription of the Australian Spider Crab *Eurynome granulosa* Baker (Crustacea, Brachyura, Majidae). Rec. S. Aust. Mus. 15 (1): 29-37, 5 figs.
- (in press, a). The Genus *Chlorinoides* (Crustacea, Brachyura, Majidae). 1. A Redescription of *C. tenuirostris* Haswell and the Status of the Genus *Acanthophrys* A. Milne Edwards. Rec. Aust. Mus.
- (in press, b). The Marine Fauna of New Zealand: Spider Crabs of the Family Majidae (Crustacea, Brachyura). N.Z. Oceanogr. Inst. Mem. 35
- Griffin, D. J. G. and Yaldwyn, J. C. (1965). A Record of the Majid Brachyuran Genus *Achaeus* from New Zealand, with Notes on the Australian Species. Trans. roy. Soc. N.Z. Zool. 6 (4): 33-51, figs. 1-8.
- Haan, W. de (1833-1850). Crustacea. In P. F. von Siebold, "Fauna Japonica." Amsterdam: Muller. Pp. xvii, xxxi, 244, 55 pls.
- Hale, H. M. (1924). The Flora and Fauna of the Nuyts Archipelago and the Investigator Group. No. 16. The Crustacea. Trans. roy. Soc. S. Aust. 48: 67-73, figs. 1, 2, pl. 1.
- (1927). The Crustaceans of South Australia. Pt. 1: 1-201, figs. 1-202. Adelaide: Govt. Printer.

- Hale, H. M. (1929). Notes on the Fauna of Dirk Hartog Island, Western Australia. No. 4. Crustacea. *Trans. roy. Soc. S. Aust.* 53: 67-70, pl. 5.
- Hartnoll, R. G. (1964). Reduction of the gill number in Spider Crabs. *Crustaceana*. 7 (2): 145-148.
- Haswell, W. A. (1879). On two new species of the genus *Stenorhynchus*. *Proc. Linn. Soc. N.S.W.* 3: 408-409.
- _____. (1880a). On the Australian Brachyura Oxyrhyncha. *Proc. Linn. Soc. N.S.W.* 4: 431-458, pls. 25-27.
- _____. (1880b). Notes on the Australian Maioid Brachyura. *Ann. Mag. nat. Hist.* (ser. 5) 5: 145-147.
- _____. (1880c). On two new species of the genus *Paratymolus* Miers from Australia. *Ann. Mag. nat. Hist.* (ser. 5) 5: 302-304, pl. XVI.
- _____. (1882a). On some new Australian Brachyura. *Proc. Linn. Soc. N.S.W.* 6: 540-551.
- _____. (1882b). Descriptions of some New Species of Australian Decapoda. *Proc. Linn. Soc. N.S.W.* 6: 750-763.
- _____. (1882c). Catalogue of the Australian Stalk- and Sessile-eyed Crustacea. Sydney: Australian Museum. Pp. xxiv, 326, 4 pls.
- Holthuis, L. B. (1962). Forty-seven genera of Decapoda (Crustacea); Proposed Addition to the Official List. Z.N. (S.) 1499. *Bull. zool. Nomencl.* 19 (4): 232-253.
- Laurie, R. D. (1906). Report on the Brachyura collected by Professor Herdman at Ceylon in 1902. *Rep. Pearl. Oyster Fish., Ceylon* 5: 349-432, figs. 1-12, pls. I, II.
- McCulloch, A. R. (1908). Studies in Australian Crustacea. No. 1. *Rec. Aust. Mus.* 7: 51-59, pl. xii.
- _____. (1913). Studies in Australian Crustacea. No. 3. *Rec. Aust. Mus.* 9: 321-353, pls. x, xi.
- Macleay, W. S. (1838). Brachyurous Decapod Crustacea. In A. Smith (ed.), Illustrations of the Zoology of South Africa. 5. Invertebrata 3: 53-71, pls. 2, 3. London: Smith, Elder and Co.
- McNeill, F. A. (1923). Crustacean Camoufleurs. *Aust. Mus. Mag.* 1 (8): 243-6, illus.
- _____. (1926). The Biology of North-west Islet, Capricorn Group. (J.) Crustacea. *Aust. Zool.* 4 (5): 299-318, text-figs. 1, 2, pl. XLI.
- _____. (1953). Carcinological Notes. No. 2. *Rec. Aust. Mus.* 23 (3): 89-96, pl. VII.
- _____. (In prep.) Crustacea Decapoda and Stomatopoda. *Sci. Rep. Great Barrier Reef Exped. 1928-29.*
- Man, J. G. de (1888). Bericht über die von Herrn Dr. J. Brock im indischen Archipel gesammelten Decapoden und Stomatopoden. *Arch. Naturgesch.* 53 (1): 215-600, pls. 7-22.
- Miers, E. J. (1879a). On a collection of Crustacea made by Capt. H. C. St. John, R.N. in the Corean and Japanese Seas. I. Podophthalmia. *Proc. zool. Soc. Lond.* 1879: 18-61, pls. 1-3.
- _____. (1879b). Descriptions of New or little known species of Maioid Crustacea (Oxyrhyncha) in the Collections of the British Museum. *Ann. Mag. nat. Hist.* (ser. 5) 4: 1-28, pls. IV, V.
- _____. (1879c). On the Classification of the Maioid Crustacea or Oxyrhyncha, with a synopsis of the Families, Subfamilies and Genera. *J. Linn. Soc. Lond. (Zool.)* 14: 634-673, pls. 12, 13.
- _____. (1882). On some Crustaceans collected at Mauritius. *Proc. zool. Soc. Lond.* 1882: 339-343, pls. XX.
- _____. (1884). Crustacea. In "Report on the Zoological Collections made in the Indo-Pacific Ocean during the Voyage of H.M.S. "Alert", 1881-2": 178-322, 513-575, pls. XVIII-XXXIV, XLV-LII. London: British Museum (Natural History).
- _____. (1885). On the species of *Micippa* Leach and *Paramicippa* M. Edw. *Ann. Mag. nat. Hist.* (ser. 5) 15: 1-13, pl. 1.

- Miers, E. J. (1886). Report on the Brachyura collected by H.M.S. "Challenger" during the years 1873-1876. *Rep. Voy. Challenger, Zool.* 17(2): 1-362, pls. 1-29. London, Edinburgh and Dublin.
- Milne Edwards, A. (1865). Descriptions de quelques Crustacés nouveaux, appartenant à la tribu des Maiens. *Ann. Soc. entom. Fr.* (ser. 4) 5: 133-147, pls. 3-5.
- (1872). Recherches sur la faune carcinologique de la Nouvelle-Calédonie. *Nouv. Arch. Mus. Hist. nat., Paris.* 8: 229-267, pls. X-XIV.
- (1873). Descriptions de quelques Crustacés nouveaux ou peu connus provenant du Musée de M. C. Godeffroy. *J. Mus. Godeffroy* 1(4): 253-264, pls. 12, 13.
- Milne Edwards, H. (1834). Histoire Naturelle des Crustacés, comprenant l'anatomie, la physiologie et la classification de ces animaux. I: i-xv, 1-468. Atlas. Paris: Lib. Roret.
- Montgomery, S. K. (1931). Report on the Crustacea Brachyura of the Percy Sladen Trust Expedition to the Abrolhos Islands under the leadership of Prof. W. J. Dakin in 1913, along with other Crabs from Western Australia. *J. Linn. Soc. Lond. (Zool.)* 37: 405-465, 1 fig., pls. 24-30.
- Ortmann, A. (1894). Crustaceen. In R. Semon, *Zoologische Forschungsreisen Australien und dem Malayischen Archipel.* 5 (1): 30-80, 3 pls. (*Denkschr. med. naturw. Ges Jena* 8: 1-80.).
- Rathbun, M. J. (1892). Catalogue of the Crabs of the family Periceridae in the U.S. National Museum. *Proc. U.S. nat. Mus.* 15: 231-277, pls. 28-40.
- (1893). Catalogue of the Crabs of the family Maiidae in the U.S. National Museum. *Proc. U.S. nat. Mus.* 16: 63-103, pls. 3-8.
- (1894). Catalogue of the Crabs of the family Inachidae in the U.S. National Museum. *Proc. U.S. nat. Mus.* 17: 43-75, pl. 1.
- (1897). A revision of the nomenclature of the Brachyura. *Proc. biol. Soc. Wash.* 11: 153-167.
- (1906). The Brachyura and Macrura of the Hawaiian Islands. *Bull. U.S. Fish. Comm.* 23 (3): 827-930, text figs. 1-79, pls. 3-25.
- (1911). The Percy Sladen Trust Expedition to the Indian Ocean in 1905: Marine Brachyura. *Trans. Linn. Soc. Lond. (Zool.)* (ser. 2) 14 (2): 191-261, pls. 15-20.
- (1914). Stalk-eyed Crustaceans collected at the Monte Bello Islands. *Proc. zool. Soc. Lond.* 1914: 653-664, pls. I, II.
- (1916). Scientific Results of the Philippine Cruise of the Fisheries Steamer "Albatross" 1907-10. No. 34. New Species of Crabs of the Families Inachidae and Parthenopidae. *Proc. U.S. nat. Mus.* 50: 527-559.
- (1918). Report of the Spider Crabs obtained by the F.I.S. "Endeavour" on the Coasts of Queensland, New South Wales, Victoria, South Australia and Tasmania. *Biol. Res. Endeavour 1909-14* V (1): 1-29, 15 pls.
- (1924). Results of Dr. E. Mjoberg's Swedish Scientific Expeditions to Australia 1910-1913. 37. Brachyura, Albuneidae and Porcellanidae. *Ark Zool.* 16 (23): 1-33, 1 pl.
- (1925). The Spider Crabs of America. *Bull. U.S. nat. Mus.* 129: 1-613, 153 figs., 283 pls.
- Sakai, T. (1932). Notes on some rare materials of Japanese Oxyrhyncha. *Sci. Rep. Tokyo Bunrika Daig.* (B) 1 (4): 41-59, pls. II, III.
- (1934). Brachyura from the coast of Kyushu, Japan. *Sci. Rep. Tokyo Bunrika Daig.* (B) 1 (25): 281-330, text-figs. 1-26, pls. XVII, XVIII.
- (1935). New or rare species of Brachyura, collected by the "Misago" during the zoological survey around the Izu-peninsula. *Sci. Rep. Tokyo Bunrika Daig.* (B) 2 (32): 63-88, text-figs. 1-17, pls. VI-VIII.


- Sakai, T. (1936). Report on the Brachyura collected by Mr. F. Hiro at Palao Islands. *Sci. Rep. Tokyo Bunrika Daig.* (B) 2 (37): 155-177, text-figs. 1-7, pls. XII-XIV.
- (1938). Studies on the Crabs of Japan. III. Brachynathidae, Oxyrhyncha: 193-364, text-figs. 1-55, pls. I-XLI. Tokyo: Yokendo Co.
- (1965). The Crabs of Sagami Bay collected by His Majesty the Emperor of Japan. Tokyo: Maruzen. Pp. xvi, 1-206, 1-26 (English), 1-92, 27-32 (Japanese), 26 text-figs., 100 pls. (coloured), 1 map.
- Sankarankutty, C. (1961). On some crabs (Decapoda-Brachyura) from the Laccadive Archipelago. *J. Mar. biol. Ass. India* 3 (1 & 2): 120-136, figs. 1, 2.
- (1962). On Decapoda Brachyura from the Andaman and Nicobar Islands. 3. Families: Calappidae, Leucosiidae, Parthenopidae, Maiidae, and Gecarcinidae. *J. Mar. biol. Ass. India* 4 (1): 151-164, figs. 1-23.
- Stephensen, K. (1945). The Brachyura of the Iranian Gulf, with an Appendix; The Male Pleopoda of the Brachyura. *Danish sci. Invest. Iran* IV: 57-237, 60 figs.
- Stephenson, W. (1962). Evolution and Ecology of Portunid Crabs, with especial reference to Australian Species. In G. W. Leeper (ed), "The Evolution of Living Organisms": 311-327, 3 tables. Melbourne: University Press.
- Stimpson, W. (1857). Prodromus descriptionis animalium evertebratorum, quae in Expeditione ad Oceanum Pacificum Setentrionalem, e Republica Federate missa, C. Ringgold et J. Rodgers Ducibus, observavit et descripsit. III. Crustacea Maioidea. *Proc. Acad. nat. Sci. Philad.* 9 (25): 216-221.
- (1907). Report on the Crustacea (Brachyura and Anomura) collected by the North Pacific Exploring Expedition. *Smithson. misc. Coll.* 49 (1717): 1-240, pls. i-xxvii.
- Tesch, J. J. (1918). The Decapoda Brachyura of The Siboga Expedition. I. Hymenosomatidae, Retroplumidae, Ocypodidae, Grapsidae and Gecarcinidae. *Siboga Exped. Monogr.* 39c (82): 1-148, pls. 1-6.
- Ward, M. (1933). New genera and species of marine Decapoda Brachyura. *Aust. Zool.* 7: 337-394, pls. 21-23.
- (1936). Crustacea Brachyura from the coasts of Queensland. *Mem. Queensl. Mus.* II: 1-13, pls. 1-3.
- White, A. (1847). Descriptions of new Crustacea from the Eastern seas. *Ann. Mag. nat. Hist.* 20: 61-63.
- (1848). Descriptions of New or little-known Crustacea in the Collections of the British Museum. *Ann. Mag. nat. Hist.* (ser. 2) 1: 221-226.
- Whitelegge, T. (1900). Scientific Results of the Trawling Expedition of H.M.C.S. "Thetis", off the coast of New South Wales, in February and March, 1898. Crustacea. Part 1. *Mem. Aust. Mus.* IV: 135-199, pls. 32-35.
- Yaldwyn, J. C. (1964). The Ornamental Sea-toad. *Aust. nat. Hist.* XIV (11): 362. (see also: Anon (1965). *Nature* 205 (4977): 1163).

EXPLANATION OF PLATES.


- Plate xv* (a) *Hyastenus auctus* Rathbun. HOLOTYPE, male, carapace length from photo (including rostral spines in all cases) about 30 mm. Sulu Sea, Philippine Islands, *Albatross Exped.*, U.S. National Museum no. 48214 (Photo: U.S. Nat. Mus.).
(b) *Hyastenus sebae* White. LECTOTYPE (selected on the advice of Dr. I. Gordon), dorsal view, female, carapace length from photo about 11.5 mm. Corregidor, Philippine Islands, British Museum (N.H.) no. 43.6 (Photo: British Museum).
(c) *Hyastenus sebae* White. LECTOTYPE, ventral view (Photo: British Museum).
(d) *Huenia bifurcata* Streets. Male, carapace length 22 mm., setae cleaned from shaft of rostrum only. New South Wales, Australian Museum no. P. 14961 (Photo: Anthony Healy).
(e) *Huenia bifurcata* Streets. Female, carapace length 21 mm, uncleaned. Port Jackson, N.S.W., Aust. Mus. no. G. 5102 (Photo: Anthony Healy).

- Plate xvi* (a) *Schizophrys aspera* (H. Milne Edwards). Dorsal view, male, carapace length 60.5 mm. Lord Howe Island, W.R.B. Oliver collection, Dominion Museum, Wellington (Photo: Athol Beswick).
(b) *Schizophrys aspera* (H. Milne Edwards). Ventral view of same specimen (Photo: Athol Beswick).

- Plate xvii* (a) *Paranaxia serpulifera* (Guérin). Male, carapace length 102 mm. Darnley Island, Torres Strait, Aust. Mus. no. G. 2469 (Photo: Anthony Healy).
(b) *Tiarinia elegans* Haswell. HOLOTYPE, male, carapace length 14.5 mm. Off Broughton Island, near Port Stephens, N.S.W., 25 fms. Aust. Mus. no. G. 5140 (Photo: Anthony Healy).
-



SPIDER CRABS (for explanation see page 298).



SPIDER CRABS (for explanation see page 298).