

Boston Society of Natural History.

FROM

Mr. Walter Faxon.

Received Nov. 15, 1895.

DESCRIPTIONS OF A NEW GENUS AND FOUR NEW SPECIES
OF CRABS FROM THE ANTILLEAN REGION.

By MARY J. RATHBUN,
Aid. Department of Marine Invertebrates.

THE CRABS described below were, with one exception, obtained in the extended cruise of the United States Fish Commission steamer *Albatross* to the Gulf of Mexico and Caribbean Sea in 1885. The species of *Actaea* formed a part of the large collection of invertebrates brought from Florida by Dr. Edward Palmer in 1884.

Family PARTHENOPIDAE.

THYROLAMBRUS, new genus.

Carapace broader than long, deeply eroded. Frontal and antero-lateral region strongly deflexed. Entire surface covered with stellar granules, which unite to form ridges outlining irregular pits. Maxillipeds broad, fitting closely together and filling the buccal cavity; ischium subrectangular posteriorly, slightly oblique anteriorly; merus broader than long, with a slight notch at the antero-internal angle, in which the first joint of the palpus is fitted in a transverse direction; the remainder of the palpus is concealed beneath the merus. Chelipeds of moderate length; manus much more slender than the merus and armed on the inner or anterior side with 2 rows of long, sharp, curved spines, which are continued on the fingers.

THYROLAMBRUS ASTROIDES, new species.

Carapace about two-thirds as long as wide, thick, slightly wider at the postero-lateral than at the lateral angles; frontal and antero-lateral regions almost perpendicular. Posterior margin directed slightly forward and outward. Besides the small pits everywhere present on the surface there are other larger depressions. A deep hollow between the orbits is continued backward by a shallow sulcus to the post-medial region. Two deep depressions occur at the inner branchial angles. The cardiac area is well defined and is bounded posteriorly by a trans-

verse linear sulcus. The hepatic region is outlined by a series of depressions. The ridges of the surface are elevated at intervals into rough acute tubercles. Rostrum very broad, arcuate as seen from above, produced downwards at the middle in a small, triangular, denticulate tooth which extends backward to the antennular cavities. Orbits small, circular; eye-peduncles covered with stellar granules, and with a row of 3 or 4 spinules next the cornea on the upper side. Hepatic region with a triangular marginal tooth. Lateral margin of the branchial region with about 7 small granulate teeth. Teeth of posterior margin very shallow. Antero-internal angle of the basal antennal joint barely touching the front. Exognath of maxilliped slender; endognath with a longitudinal row of 3 spinules. The pterygostomial groove is continued on the subbranchial regions. The sternum in the male has 3 prominent ridges on either side of the abdomen, and is deeply hollowed at the anterior end.

Merus of cheliped thick, with short spines on the anterior and upper surfaces. Carpus with 3 spinules on inner margin. The spines of the propodus and dactylus number 5 or 6 in the lower series and 6 or 7 in the upper. They are curved inward and directed toward the extremity of the fingers. The fingers are slender, curved inward, their tips prolonged in sharp spines; the stellar granules are arranged longitudinally; prehensile edges armed with fine sharp irregular teeth or spines. The ambulatory legs are very rough. The meral joints have 1 crest above and 2 below. Dactyli short and slender, armed with sharp spines, and terminating in an acuminate horny tip.

Measurements.—Length of female, 16 mm.; width, 23.5; thickness at epistome, 6; length of cheliped, about 32; length of merus, below, 11; length of propodus, 14. Length of male, 14; width, 20; length of cheliped, about 34; of merus, 12; of propodus, 15.

Locality.—Off Havana, Cuba, in lat. $23^{\circ} 10' 42''$ N., long. $82^{\circ} 18' 24''$ W., 67 fathoms, white coral, 2 females (No. 9507, U. S. N. M.); and in lat. $23^{\circ} 10' 40''$ N., long. $82^{\circ} 20' 15''$ W., 189 fathoms, coral, 1 male (No. 9515, U. S. N. M.).

SOLENOLAMBRUS DECEMSPINOSUS, new species.

Closely allied to *S. typicus*, Stimpson. Antero-lateral margin convex, area between the gastric ridges narrower than in *S. typicus*, gastric and cardiac prominences slender spines. There are 8 additional dorsal spines: 2 on each branchial ridge, of which the marginal is the longer, 1 at each posterior angle, and 1 on the postero-lateral margin midway between the last and the branchial spine. The punctures of the carapace are very fine and scattered; in *S. typicus* they are coarse and anteriorly crowded. The sternum (in the male) is smooth in front of the abdomen. The terminal segment of the abdomen is much longer and narrower distally than in *S. typicus*, its sides deeply concave. The merus of the maxillipeds is narrower and more produced at the antero-external angle than in *S. typicus*.

The chelipeds are similar in ornamentation to those of *S. typicus*; the upper margin of the outer surface of the manus is furnished with 10 granulated teeth, the lower margin with about 12; as in *S. typicus*, the surfaces of the palm have bunches of granules arranged in 2 longitudinal rows. The immovable finger is shorter and more deflexed than in *S. typicus*, and in consequence the dactylus is also more deflexed, being, when closed, nearly at a right angle with the outer or upper surface of the palm. Color of fingers in alcohol, red.

Measurements.—Length of carapace of male, 6 mm.; width, 7.

Locality.—Gulf of Mexico, in lat. 28° 44' N., long. 85° 16' W., 60 fathoms, gray sand, station 2404, one male (No. 18157, U. S. N. M.).

Family CANCRIDÆ.

ACTEA PALMERI, new species.

Carapace covered with 36 large, nodose prominences separated by deep sinuses filled with long silky hair, which also conceals the posterior portion of the carapace as well as the entire lower surface of the crab. The nodules of the surface are very convex and are each composed of a number of smooth, shining, bead-like granules crowded close together. The frontal lobes or nodules are thick, with convex, entire margins, and are separated by a deep sulcus. There are 6 orbital nodules, one very small inner-orbital, followed by 1 large and 2 small, and 2 suborbital nodules. The basal antennal joint is also a thick, shining, compound nodule. The carapace has 4 nodules on the lateral margin which project upward and not outward, the margin itself being entire. A small median lobule is visible near the posterior margin.

Chelipeds with merus smooth and hairy, carpus with 6 nodules, and manus with 5, the remainder of the surface silky-hairy. Immobile finger and distal half of dactylus smooth and shining, horn-colored, with white tips. The fingers are broad, compressed and sharp-pointed. The ambulatory legs have 2 small nodules on the carpal joints, 1 or 2 on the propodal joints, and 2 on the meral joints of the last pair.

Measurements.—Length of male, 16 mm.; width, 21. Length of female, 14; width, 19.

Locality.—Rodriguez Creek, Florida, Dr. Edward Palmer; 1 male, 1 female carrying a large quantity of minute eggs, and 2 young specimens, male and female (No. 13927, U. S. N. M.).

PILUMNUS DIOMEDEÆ, new species.

Carapace of moderate width, beset with long yellow hairs arising from low spinules. Front with 2 produced lobes, each bearing 4 slender spines; a longer incurved spine is placed near the antenna. Orbital spines 9, 2 on the upper margin, 1 at the outer angle, and 6 below. Of the suborbital spines, the 2 outermost are separated by a deep fissure. There are 4 strong antero-lateral spines, including the orbital; between the first and second there is a small spine, and the second

spine has 1 or 2 accessory spinules. The subhepatic and pterygostomial regions are spinulous. There are 2 small spines forming a longitudinal line with the inner suborbital spine.

Chelipeds unequal, spinous and long-hairy. Merus with surface minutely spinulous, margins spinous, the upper margin furnished distally with 2 spines longer than the others. Carpus with outer surface spinous and spinulous, a strong spine at the inner angle. Manus with 4 slender spines on upper margin, lower margin spinulous, spines of outer surface arranged in longitudinal rows, inner surface minutely granulous. Fingers spinulous and hairy proximally, horn-colored, with teeth and tips almost white. Ambulatory legs very long, slender and hairy; margins of meral joints and upper margin of carpal and propodal joints spinous.

The type specimen has the posterior portion of the carapace and sternum broken off and is without the last 3 pairs of legs. The only other individual is a soft-shell female, very imperfect.

Measurements.—Width of carapace, 16 mm.; width of front, 6; length of longest hairs of carapace, about 6.

Localities.—Off Havana, Cuba, in lat. $23^{\circ} 10' 40''$ N., long. $82^{\circ} 20' 15''$ W., 184 fathoms, fine gray and white coral, station 2345, type (No. 9526, U. S. N. M.). Off Yucatan, lat. $20^{\circ} 59' 30''$ N., long. $86^{\circ} 23' 45''$ W., 130 fathoms, coral, station 2354 (No. 18158, U. S. N. M.).

This species in its long ambulatory legs resembles *P. gracilipes*, A. Milne-Edwards, which differs, according to that writer, in its unarmed superior orbital margin and short hair.

DIGEST OF THE
LIBRARY REGULATIONS.

No book shall be taken from the Library without the record of the Librarian.

No person shall be allowed to retain more than five volumes at any one time, unless by special vote of the Council.

Books may be kept out one calendar month; no longer without renewal, and renewal may not be granted more than twice.

A fine of five cents per day incurred for every volume not returned within the time specified by the rules.

The Librarian may demand the return of a book after the expiration of ten days from the date of borrowing.

Certain books, so designated, cannot be taken from the Library without special permission.

All books must be returned at least two weeks previous to the Annual Meeting.

Persons are responsible for all injury or loss of books charged to their name.