

## Epibenthic communities on the Chatham Rise, New Zealand

D. G. McKNIGHT

National Institute of Water & Atmospheric  
Research Ltd  
P. O. Box 14 901, Kilbirnie  
Wellington, New Zealand

P. K. PROBERT

Department of Marine Science  
University of Otago  
P. O. Box 56  
Dunedin, New Zealand

**Abstract** Samples of epibenthic macrofauna were collected using a small sledge trawl from 40 stations on the Chatham Rise (42°34'–44°47'S; 175°34'E–179°03'W) at water depths of 237–2039 m. 218 taxa were collected, predominantly echinoderms, crustaceans, and molluscs. Multivariate analysis based on 85 species indicated three epibenthic communities, the shallowest characterised mainly by crustaceans and two deeper water communities mainly characterised by echinoderms: (1) a community on predominantly sandy sediments on the crest and shallower flanks of the Rise at 237–602 m that included as characteristic species *Munida gracilis*, *Phylladorhynchus pusillus*, *Campylonotus rathbunae*, *Pontophilus acutirostris*, and *Acutiserolis bromleyana* (Crustacea), *Amphiura lanceolata* (Ophiuroidea), *Cuspidaria fairchildi*, and *Euciroa galatheae* (Bivalvia); (2) a community associated with muddy sediments at 462–1693 m that included *Ypsilothuria bitentaculata* and *Pentadactyla longidentis* (Holothuroidea), *Brissopsis oldhami* (Echinoidea), and *Amphiophiura ornata* (Ophiuroidea); and (3) a community on muddy sediments at 799–2039 m that included *Ophiomusium lymani* (Ophiuroidea), *Porcellanaster ceruleus* (Asteroidea), *Gracilechinus multidentatus* (Echinoidea), and

*Aeneator recens* (Gastropoda). Bathymetric range of the communities was not symmetrical on either side of the Rise; there was a marked submergence on the north side similar to the vertical displacement of the core of the Antarctic Intermediate Water.

**Keywords** New Zealand; Chatham Rise; bathyal benthos; epifauna; communities; Echinodermata; Crustacea; Mollusca

### INTRODUCTION

New Zealand has an extensive bathyal zone that includes a number of major plateaux and ridges (Carter 1980). A prominent feature of the New Zealand bathyal is the Chatham Rise, a broad ridge lying to the east of central New Zealand and extending for c. 1400 km to c. 168° W, east of the Chatham Islands. To the west of the Chatham Islands the Rise is generally flat-topped at depths of 200–400 m, but to the east the depth gradually increases to c. 3000 m (Thompson 1991). At the western end of the Rise is the Mernoo Gap, a narrow depression, deeper than 500 m, that separates the Rise from the New Zealand continental shelf. On the Rise lie four banks with depths of less than 250 m: Mernoo, Veryan, and Reserve Banks toward the western margin, and the smaller Matheson Bank nearer the Chatham Islands. On large-scale bathymetric charts the Rise appears smooth-topped, but more detailed mapping reveals considerable local relief (Mitchell & Cullen 1984). Surface sediments are in general relatively fine-grained, with terrigenous sediments giving way to pelagic sediments at c. 179° E (McDougall 1982; Mitchell et al. 1989). Located over the Rise is the Subtropical Convergence, a circumglobal front where subtropical and subantarctic surface waters meet (Heath 1985). The Rise is of economic importance, with major fisheries (Francis & Fisher 1979; Fenaughty & Uozumi 1989) and large deposits of potentially mineable phosphorite (Cullen 1987).

Earlier benthic studies of the Rise have dealt mainly with discrete elements of the fauna and have




Fig. 1 Locality diagram showing transects.

been largely systematic in approach (e.g., Fell 1960; Knox 1960) although Hurley (1961) provides some observations on benthic structure. Dawson (1984) and Probert et al. (1997) examined effects of mining and trawling on benthos of the Rise. Probert & McKnight (1993) and Probert et al. (1996) noted relatively higher macrobenthic biomass and faunal densities on the southern side of the Rise, which they suggest may be the result of the enhanced organic flux associated with the Subtropical Convergence. As part of a multidisciplinary study of the Chatham Rise region the following contribution examines epibenthic communities on the Rise, mainly between 177°E and 179°W.

## MATERIALS AND METHODS

Twenty-four epibenthic samples were taken with a small sledge-trawl on the Chatham Rise during September 1989 (V-prefixed stations in Table 1). Stations were sited along three north-south transects: a western transect at 177°E, a central transect at 179°E, and an eastern transect at 179°W (Fig. 1). Positions of the V stations are those in Probert & McKnight (1993), except that weather conditions prevented sampling at the southern end of the eastern transect (>568 m) and three off-transect stations were added: V361 (30 km west of the central transect), and stations V393 and V394 (65 and 117 km west of the western transect, respectively). (These three stations are considered

below as belonging to the central and western transects.) A further 16 samples were taken during September 1993 (W stations in Table 1) on the central transect. Transect positions are shown in Fig. 1 and sample sites detailed in Table 1. In total, 24 of the stations were on the central transect, 10 on the western and six on the eastern. The trawl was similar to that described by McDougall (1973), with a width of 1.2 m and a height of 0.45 m. The outer bag, c. 2 m long, had a mesh of 15 mm knot to knot, with a finer mesh of 7 mm knot to knot lining the distal half of the net. At each site the trawl was towed for 20 min at c. 0.5 m s<sup>-1</sup>. Most samples retained some fine sediment and all were washed on a 1-mm mesh then preserved. In the laboratory the animals were sorted to major taxa and identified, if possible to species. Polychaetes were relatively uncommon in the samples and most were assigned only to family. A total of 85 taxa present in two or more samples was used in the subsequent analyses. The smaller and infaunal species were largely excluded from the subsequent analyses as they were likely to have been sampled inconsistently by the trawl. The presence/absence data were subject to numerical classification and ordination analyses. Classification was performed using the Czekanowski coefficient and by group-average sorting (Lance & Williams 1967). Ordination was carried out using multidimensional scaling (MDS) (Field et al. 1982). Community scores for species were calculated as in Grange (1979), but modified

to exclude the bioindex value. Thus the community score for a species in a site group is the percentage occurrence within the group multiplied by the proportion of total sites where the species is present. Sediment samples taken from a small dredge also used at some of the sampling sites (Table 1) were analysed for grain-size in the size categories: gravel (>2.00 mm), sand (2.00–0.062 mm), silt (4–62  $\mu\text{m}$ ), and clay (<4  $\mu\text{m}$ ), and expressed as percentage weight.

## RESULTS

A total of 218 taxa was recognised from the entire collection, including a considerable number of as yet undescribed species. The best represented major taxa were Echinodermata (52 spp.), Crustacea (44 spp.), and Mollusca (36 spp.). Table 2 shows the species used in multivariate analysis.

The main feature of the dendrogram (Fig. 2) is a central cluster of 20 stations, here called Group

**Table 1** Station positions, water depth, and community grouping.

Station no.	Latitude (S)	Longitude (E or W)	Depth (m)	Group designation
V361†	43° 30.6'	178° 38.11'E	345	A
V363†	44° 15.22'	178° 58.53'W	560–568	A
V365†	43° 44.92'	179° 00.35'W	399	A
V366†	43° 29.69'	178° 59.55'W	499	A
V367†	43° 14.31'	179° 02.18'W	471	A
V368†	42° 49.87'	179° 00.45'E	1040	B
V369†	43° 05.67'	178° 59.94'E	399	A
V370†	42° 41.70'	179° 03.42'W	1024	B
V371†	42° 59.39'	179° 03.30'W	533	A
V372†	43° 20.22'	178° 58.88'E	415–409	A
V373†	43° 35.48'	178° 59.50'E	385	A
V374†	43° 51.19'	178° 58.81'E	480	A
V375	44° 05.97'	179° 01.08'E	815	C
V376	44° 20.29'	179° 00.34'E	1239	C
V385	44° 19.81'	176° 59.90'E	1086	C
V386	44° 05.25'	177° 00.05'E	665	B
V387	43° 49.62'	176° 59.82'E	498–497	A
V388	43° 34.83'	176° 59.91'E	331–328	A
V389	43° 20.02'	176° 59.83'E	237	A
V390	43° 05.00'	177° 00.00'E	330	A
V391	42° 49.97'	176° 59.91'E	476	A
V392	42° 34.32'	176° 59.70'E	1390	B
V393	42° 54.89'	176° 12.53'E	538	A
V394	43° 01.49'	175° 34.25'E	396	A
W247	44° 47.4'	178° 59.8'E	1932–1963	C
W248	44° 36.0'	178° 55.6'E	1442–1463	C
W249	44° 18.9'	179° 00.0'E	1200–1230	C
W250	44° 05.4'	179° 00.0'E	799–815	C
W251	43° 50.0'	178° 59.5'E	469–462	B
W252	43° 37.7'	178° 59.8'E	400–428	A
W255	44° 40.9'	179° 01.2'E	1600–1706	C
W257	43° 22.5'	179° 00.0'E	400–390	A
W258	42° 57.9'	178° 59.8'E	602–499	A
W259	42° 54.1'	179° 00.2'E	812–715	B
W260	42° 57.0'	179° 00.0'E	665–625	B
W261	42° 47.2'	178° 59.6'E	1212–1126	B
W262	42° 44.0'	179° 00.2'E	1410–1281	B
W263	42° 41.0'	179° 00.3'E	1660–1693	B
W265	42° 38.3'	178° 59.9'E	2039–1687	C
W273	42° 39.0'	178° 57.2'E	1915–1491	C

†Sediment particle size analyses available.

**Table 2** Occurrence of species used for multivariate analysis with community scores (> 20). (E, endemic; +, present; - absent; G, ovigerous or gonads appear ripe.) (Also recorded as G were the crustaceans *Acanthophyra pelagica*, *Prionocrangon curvicaulis*, and *Nectocarcinus antarcticus*.)

Taxa	Community group			
	A	B	C	
<b>Cnidaria</b>				
<i>Flabellum knoxi</i>	+	+	+	
<i>Edwardsia</i> sp.	+	-	+	
<b>Polychaeta</b>				
<i>Hyalinoecia tubicola</i>	29	+	+	
<i>Aphrodita talpa</i>	43	+	+	
<b>Crustacea</b>				
<i>Acutiserolis bromleyana</i>	51	+	+	G
<i>Pontophilus acutirostris</i>	55 E	-	-	G
<i>Notopandulus magnoculus</i>	41	+	-	G
<i>Campylonotus rathbunae</i>	70 E	-	-	G
<i>Prionocrangon curvirostris</i>	+	+	+	
<i>Metacrangon knoxi</i>	+	-	-	
<i>Pasiphaea notosivado</i>	+	+	-	G
Euphausiid sp.	+	-	+	
<i>Nephrops challengerii</i>	20 E	-	-	
Polychelid sp.	+	-	-	
<i>Munida gracilis</i>	81	+	+	G
<i>Gastroptychus novaezealandiae</i>	+	-	-	G
<i>Phylladorhynchus pusillus</i>	55	+	-	G
<i>Carcinoplax victorienesis</i>	31	+	-	G
<i>Trichopeltarion fantasticum</i>	+	+	-	
<b>Mollusca</b>				
<i>Dentalium zealandicum</i>	+	-	-	
<i>Dentalium</i> sp.	-	-	20 E	
<i>Poroleda lanceolata</i>	+	-	-	
<i>Neilo wairoana delli</i>	+	-	-	
? <i>Anomia</i> sp.	+	-	-	
<i>Cuspidaria fairchildi</i>	55	-	+	
<i>Cuspidaria morelandi</i>	+	-	-	
<i>Euciroa galathea</i>	50	-	+	
<i>Falsilunatia powelli</i>	+	24	+	
<i>Waihaoa knoxi</i>	+	+	+	
<i>Baryspira novaezealandica benthicola</i>	+	+	-	
<i>Waipaoa marwicki</i>	-	20 E	-	
<i>Surculina expeditionis</i>	20 E	-	-	
<i>Fusitriton cancellatus</i>	-	+	+	
<i>Otukaia blacki</i>	-	-	20 E	
<i>Aeneator recens</i>	-	-	30 E	
<i>Retusa pachys</i>	+	+	+	
<b>Echinodermata</b>				
<i>Psilaster acuminatus</i>	+	+	+	
<i>Plutonaster knoxi</i>	+	+	+	
<i>Plutonaster fragilis</i>	-	-	+	
<i>Porcellanaster ceruleus</i>	-	-	42 E	
<i>Benthopecten pikei</i>	-	+	27	
<i>Pectinaster mimicus</i>	-	+	23	
<i>Pseudarchaster garricki</i>	+	+	+	
<i>Ceramaster lennoxkingi</i>	-	-	+	
<i>Pillsburiaster aotearanus</i>	+	-	+	

(continued)

Table 2 (continued)

Taxa	Community group			
	A	B	C	
<i>Hippasteria trojana</i>	–	–	20 E	
<i>Crossaster japonicus</i>	21	–	+	
<i>Zoroaster spinulosus</i>	+	–	20 E	
Ophiomyxid sp.	+	–	20 E	
<i>Ophiacantha vilis</i>	37	–	+	
<i>Ophiacantha yaldwyni</i>	+	–	–	
<i>Ophiophthalmus relictus</i>	–	+	32	
<i>Amphiura lanceolata</i>	63	+	–	G
<i>Amphiura</i> sp.	–	–	+	
<i>Amphioplus pegasus</i>	+	–	–	
<i>Ophiactis hirta</i>	+	–	–	
<i>Ophiactis abyssicola</i>	–	–	23 E	
<i>Ophiochiton lentus</i>	–	20 E	–	
<i>Ophiura irrorata</i>	42	+	–	
<i>Ophiura ooplax</i>	+	–	–	
<i>Amphiophiura ornata</i>	–	30 E	–	
<i>Ophiomisidium irene</i>	+	+	–	
<i>Ophiozonella stellamaris</i>	+	+	+	
<i>Ophiozonella stellata</i>	–	20 E	–	
<i>Ophiomusium lymani</i>	–	+	49	
<i>Ophiernus</i> sp.	–	–	20 E	
<i>Goniocidaris parasol</i>	23	+	–	
<i>Phormosoma bursarium</i>	+	+	+	
<i>Caenopedina otagoensis</i>	–	–	20 E	
<i>Gracilechinus multidentatus</i>	–	–	30 E	G
<i>Urechinus antipodeanus</i>	–	–	20 E	
<i>Hemiaster expergitus gibbosus</i>	+	+	+	
<i>Brissopsis oldhami</i>	32	34	+	
<i>Echinocardium lymani</i>	31 E	–	–	
<i>Paramaretia peloria</i>	21	–	+	
<i>Spatangus multispinus</i>	+	+	–	
<i>Spatangus lutkeni</i>	+	+	–	
<i>Heterothyone alba</i>	+	–	+	
<i>Pentadactyla longidentis</i>	+	41	+	
<i>Ypsilothuria bitentaculata</i>	+	49	+	
<i>Echinocucumis hispida</i>	35 E	–	–	
<i>Paracaudina chilensis</i>	43	+	–	
<i>Bathyploetes natans</i>	+	+	+	
Holothurian sp. E	+	+	+	
Holothurian sp. F	+	–	–	

A, occupying a water depth range of 237–602 m and two groupings each of 10 stations: Group B at depths of 462–1693 m and Group C at 799–2039 m. The two smaller clusters are less discrete and subdivide at a similarity level lower than that required to recognise the integrity of the large grouping (<0.25). The three groups were less well defined in the MDS plot and, in the absence of clear alternatives, the groups referred to above were adopted for further examination.

Community A comprised 20 stations from the crest and shallower flanks of the Rise, in water

depths of 237–602 m, and was present on all three transects (Table 1). Sediment particle size analyses, available for 10 stations, indicated predominantly muddy sands; mean percent weights were: clay 8% (SD 4.1), silt 20% (6.2), sand 68% (7.1), and gravel 4% (8.0). Sixty-three species were recorded, of which 19 were “endemic”, 13 shared with Community B, 11 with Community C, and 20 with both. Community scores for the species ranged from 81.0 to 1.7. Eight species had a score of 50.0 or more, and a further 10 a score between 50.0 and 25.0. Twenty-three species had a score of 20.0 or




Fig. 2 Dendrogram of stations.

more (Table 2). These comprised 8 crustaceans, 4 echinoids, 3 ophiuroids, 2 bivalves, 2 polychaetes, 2 holothurians, 1 asteroid, and 1 gastropod. Mobile crustaceans were noticeable among the high-scoring characteristic species.

Community B was recorded from 10 stations on both flanks of the Rise (Table 1). To the north it was present at eight sites on all three transects, at 625–1693 m, and to the south at two sites on the western and central transects at 462–665 m.

Sediment particle size analyses, available for only two stations, indicated sandy mud; mean percent weights were: clay 29% (SD 6.0), silt 35% (7.2), and sand 36% (13.1). Forty-two species were recorded of which four were endemic, 13 shared with Community A, five with Community C, and 20 with both. Community scores for the species ranged from 49.0 to 0.5, and eight species had a score of >20.0 (Table 2). These comprised 3 ophiuroids, 2 holothurians, 2 gastropods, and 1 “irregular” echinoid. None would be considered as highly mobile.

Community C occurred at 10 stations at water depths of 799–2039 m, predominantly at southern stations of the central transect (Table 1). In particular it was recorded on the central (nine stations) and western (one station) transects, and spanned depth ranges of 799–1963 m to the south of the Rise (eight stations) and 1491–2039 m to the north (two stations). No sediment data were available. Forty-nine species were present, 13 were endemic, 11 shared with Community A, five with Community B, and 20 with both. Community scores ranged from 49.2 to 0.5. Sixteen species had a score of >20.0, including 5 ophiuroids, 5 asteroids, 3 echinoids, 2 gastropods, and 1 scaphopod. Highly mobile species were absent.

## DISCUSSION

In recent years there have been a considerable number of studies on communities of the continental slope and deeper oceanic regions, sufficient for the production of several comprehensive compilations (e.g., Menzies et al. 1973; Rowe 1983; Gage & Tyler 1991). Studies outside New Zealand have progressed to the recognition of large-scale, widespread communities, the detailed examination of bathymetric distributions, and the spatial distribution of individual species. There is, however, a dearth of comparable studies for the New Zealand continental margins. The present contribution must be viewed as a preliminary study and there is as yet little scope to correlate these results with those from elsewhere. To date there has been no widespread survey of epifaunal communities of the New Zealand slope, nor even those of the shelf. The only extensive study of local benthic communities is that of McKnight (1969), which relates primarily to infaunal shelf communities. Nevertheless, from a comparative viewpoint, this is of some relevance here. Some taxa that are probably shallow

burrowers were commonly collected by the sledge-trawl used in the present study and were included in the analysis. Thus in characterising the communities there may be some overlap between the “infaunal” and “epifaunal” elements, although for some species these designations are somewhat arbitrary.

Community A is essentially the *Serolis bromleyana*–*Spatangus multispinus* community described by Hurley (1961) from the Chatham Rise at depths of 403–604 m. Of the 38 species listed by Hurley (1961), 11 are represented in the characteristic species of the present data; only one species is represented in Community B, and two in Community C. More species would have been represented in both lists if dead molluscs were listed for the present data. Characteristic species, besides the two nominate species, were *Campylonotus rathbunae*, *Nassarius ephamillus*, *Micantapex paregonius*, *Falsilunatia powelli*, *Fusitriton retiolus*, *Cominella alertae*, *Columbarium mariae*, *Neilo australis?*, *Ophiura irrorata*, and probably, *Hyalinoecia tubicola*.

None of these species appears restricted to the Chatham Rise, but the extent of their respective distributions is mostly poorly known. Some of the species encountered in this study are known to be widely distributed at bathyal depths around New Zealand, although Dawson (1982) does identify some biogeographic differences of major bathyal regions of New Zealand. Other species taken in this study occur worldwide. For instance, *Ophiomusium lymani*, a highly characteristic species of Community C, and *Hyalinoecia tubicola* are both global continental margin species (Menzies et al. 1973).

Dawson (1984), in discussing observations from underwater television, noted the mosaic nature of the larger visible fauna of the Chatham Rise. The spatial scale of the present study is much coarser and finer detail is not resolvable. Certainly trawl samples could contain elements of dissimilar infaunal and epifaunal communities, though this is not apparent in our data, which are generally to the west of the area studied by Dawson. Dell (1956: 198) also remarked on the patchiness of benthos from the Chatham Rise at depths of up to 600 m.

### Depth ranges of communities

Of interest are the depth ranges of Communities B and C on either side of the Rise. Community B is present to the north at 625–1693 m, and to the south at 462–665 m. Community C is present at

1491–2039 m to the north and 799–1963 m to the south. Heath (1968) reported on hydrological stations, sampled in April, in the region of the present study. North of the Rise at 42°20'S (Stations D611, D612), interpolated mean temperatures at 500 and 1000 m were 9.5 and 6.1°C respectively, whereas south of the Rise at 45° S (Station D605), the corresponding temperatures were 6.6 and 3.9°C. The core of the Antarctic Intermediate Water is at a depth of c. 750 m south of the Rise and at c. 1000 m to the north (Ridgway et al. 1979). Gradual submergence of bathyal and abyssal faunas at lower latitudes is well known (Ekman 1953; Menzies et al. 1973). At the Chatham Rise, however, there appears to be an abrupt displacement possibly because of the coincidence of the Subtropical Convergence and bathymetry. Carney et al. (1983) indicated that globally a distinction between bathyal and abyssal zones commonly occurs at about the 4°C isotherm. This would roughly correspond to the division of Communities B and C on the north and south sides of the Rise. Community C may represent an abyssal fauna, given the inclusion of such echinoderms as *Ophiomusium lymani*, *Porcellanaster*, and *Urechinus* (Menzies et al. 1973), but further sampling and/or analysis of archived data will be necessary to describe zonation of deep-sea faunas in the New Zealand region.

The causes of vertical zonation of the deep-sea benthos are, however, still largely unclear (Carney et al. 1983). It has often been assumed that temperature is paramount (e.g., Ekman 1953), but such a relationship is largely untested. At least some of the observed zonation may reflect changes in trophic strategies along a gradient in food and hydrodynamic energy. On the Atlantic continental margin of the United States there is a mixture of suspension and carnivore/scavengers on the upper slope, carnivores/scavengers alone on the middle slope, and a mixture of suspension and deposit feeders on the lower slope (Gage & Tyler 1991). From the percentage occurrence of major taxonomic groups in historic data, Dawson (1984) estimated the representation of trophic groups on the Chatham Rise (bathymetric range not specified) to be: carnivores 46.9%, deposit feeders 34.6%, and suspension feeders 18.5%. Although the present data do not permit a thorough analysis of trophic groups, they indicate that the communities may differ in this regard. Prominent among the characteristic species of Community A are mobile crustaceans, likely to be scavengers or carnivores, whereas the two deeper communities lack highly

mobile characteristic species, but include more obvious deposit-feeding components. The coarser sediments on the crest of the Rise indicate that Community A is subject to more energetic hydrodynamic conditions than the deeper communities where the finer sediment would be expected to favour deposit feeding. Distribution of infaunal biomass indicates that organic flux to the benthos is higher on the south side of the Rise than on the north side (Probert & McKnight 1993). If zonation of epifaunal communities was strongly influenced by organic flux, one might expect a higher flux to the south to enable Community B to penetrate to a greater depth there, rather than the reverse being the case.

#### ACKNOWLEDGMENTS

For identifications we thank Dame Patricia Bergquist, University of Auckland, and Chris Battershill, NIWA (Porifera); Geoff Read, NIWA (Polychaeta); Des Hurley (Amphipoda and Isopoda). We also thank Ken Grange, NIWA, for assistance with identification of Mollusca, and Dick Singleton and Megan Oliver, NIWA, for data analysis. Valuable comments on the manuscript were made by NIWA colleagues and anonymous reviewers. The study was funded by the New Zealand Foundation for Research, Science & Technology (FoRST) Contract CRB 402.

#### REFERENCES

- Carney, R. S.; Haedrich, R. L.; Rowe, G. T. 1983: Zonation of fauna in the deep sea. Pp. 367–398 in: *The sea*, vol. 8, Deep-sea biology. Rowe, G. T. ed. New York, Wiley-Interscience.
- Carter, L. 1980: New Zealand region bathymetry 1:6 000 000. *New Zealand Oceanographic Institute chart, miscellaneous series 15*.
- Cullen, D. J. 1987: The submarine phosphate resource on central Chatham Rise. *DMFS reports 2*: 1–22.
- Dawson, E. W. 1982: The bathyal marine fauna of the New Zealand Exclusive Economic Zone. Pp. 43–50 in: *Fauna and hydrobiology of the shelf zones of the Pacific Ocean*. Kussakin, O. G.; Kafanov, A. I. ed. Vladivostok, Far East Science Centre, Academy of Sciences of the USSR. (*Proceedings of the XIV Pacific Science Congress, issue 4*) (In Russian with English summary.)
- Dawson, E. W. 1984: The benthic fauna of the Chatham Rise: an assessment relative to possible effects of phosphorite mining. *Geologisches Jahrbuch D 65*: 213–235.
- Dell, R. K. 1956: The archibenthal Mollusca of New Zealand. *Dominion Museum bulletin 18*: 1–235.
- Ekman, S. 1953: Zoogeography of the sea. London, Sidgwick & Jackson.
- Fell, H. B. 1960: Biological results of the Chatham Islands 1954 expedition. Part 2. Archibenthal and littoral echinoderms. *New Zealand Oceanographic Institute memoir 5*: 54–75.
- Fenaughty, J. M.; Uozumi, Y. 1989: A survey of demersal fish stocks on the Chatham Rise, New Zealand, March 1983. *New Zealand Ministry of Agriculture and Fisheries technical report 12*: 1–42.
- Field, J. G.; Clarke, K. R.; Warwick, R. M. 1982: A practical strategy for analysing multispecies distribution patterns. *Marine ecology progress series 8*: 37–52.
- Francis, R. C.; Fisher, K. A. 1979: Assessment of the deep-water fish resource of the New Zealand area. *New Zealand Ministry of Agriculture and Fisheries occasional publication 21*: 1–29.
- Gage, J. D.; Tyler, P. A. 1991: Deep-sea biology: a natural history of organisms at the deep-sea floor. Cambridge, Cambridge University Press. 504 p.
- Grange, K. R. 1979: Soft-bottom macrobenthic communities of Manukau Harbour, New Zealand. *New Zealand journal of marine and freshwater research 13*: 315–329.
- Heath, R. A. 1968: Geostrophic currents derived from oceanic density measurements north and south of the Subtropical Convergence, east of New Zealand. *New Zealand journal of marine and freshwater research 2*: 659–677.
- Heath, R. A. 1985: A review of the physical oceanography of the seas around New Zealand. *New Zealand journal of marine and freshwater research 19*: 79–124.
- Hurley, D. E. 1961: The distribution of the isopod crustacean *Serolis bromleyana* Suhm with a discussion of an associated deepwater community. *New Zealand Department of Scientific and Industrial Research bulletin 139*: 225–233.
- Knox, G. A. 1960: Biological results of the Chatham Islands 1954 expedition. Part 3. Polychaeta Errantia. *New Zealand Oceanographic Institute memoir 6*: 76–140.
- Lance, G. N.; Williams, W. T. 1967: A general theory of classificatory sorting strategies. 1. Hierarchical systems. *Computer journal 9*: 373–380.
- McDougall, J. C. 1973: Modified equipment for sediment collection and rock sampling. *NZOI records 1*: 167–170.
- McDougall, J. C. 1982: Bounty sediments. *New Zealand Oceanographic Institute chart, oceanic series, 1:1 000 000*.



- McKnight, D. G. 1969: Infaunal benthic communities of the New Zealand continental shelf. *New Zealand journal of marine and freshwater research* 3: 409–444.
- Menzies, R. J.; George, R. Y.; Rowe, G. T. 1973: Abyssal environment and ecology of the world oceans. New York, John Wiley. 488 p.
- Mitchell, J. S.; Carter, L.; McDougall, J. C. 1989: New Zealand region sediments, 1:6 000 000. *New Zealand Oceanographic Institute chart, miscellaneous series 67*.
- Mitchell, J. S.; Cullen, D. J. 1984: Central Chatham Rise bathymetry (2 sheets), 1:50 000. *New Zealand Oceanographic Institute chart, miscellaneous series 60*.
- Probert, P. K.; Grove, S. L.; McKnight, D. G.; Read, G. B. 1996: Polychaete distribution on the Chatham Rise, Southwest Pacific. *International Revue der gesamten Hydrobiologie* 81: 577–588.
- Probert, P. K.; McKnight, D. G. 1993: Biomass of bathyal macrobenthos in the region of the Subtropical Convergence, Chatham Rise, New Zealand. *Deep-sea research* 40: 1003–1007.
- Probert, P. K.; McKnight, D. G.; Grove, S. L. 1997: Benthic invertebrate bycatch from a deep-water trawl fishery, Chatham Rise, New Zealand. *Aquatic conservation: marine and freshwater ecosystems* 7: 27–40.
- Ridgway, N. M.; Heath, R. A.; Marott, C. 1979: South West Pacific Ocean: depth of the core of Antarctic Intermediate Water. *New Zealand Oceanographic Institute chart, miscellaneous series 30*.
- Rowe, G. T. ed. 1983: The sea, vol. 8, Deep-sea biology. New York, Wiley-Interscience. 560 p.
- Thompson, R. M. C. 1991: Gazetteer of seafloor features in the New Zealand region. *New Zealand Oceanographic Institute miscellaneous publication 104*: 1–64.