



An updated checklist of decapod crustaceans (infraorders Astacidea, Thalassinidea, Polychelida, Palinura, and Anomura) from the northern and northeastern Brazilian coast

PETRÔNIO ALVES COELHO^{1,3,4}; ALEXANDRE OLIVEIRA DE ALMEIDA^{1,2}; LUIS ERNESTO ARRUDA BEZERRA¹ & JESSER FIDELIS DE SOUZA-FILHO¹

¹Universidade Federal de Pernambuco, Programa de Pós-Graduação em Oceanografia, Departamento de Oceanografia. Av. Arquitetura, s/n, Cidade Universitária. 50670-901 Recife, PE, Brasil.

²Universidade Estadual de Santa Cruz, Departamento de Ciências Biológicas. Rodovia Ilhéus-Itabuna, km 16. 45662-000 Ilhéus, BA, Brasil.

³Bolsista do CNPq, E-mail: petronio.coelho@pesquisador.cnpq.br

⁴Corresponding author

Abstract

A checklist of the decapod crustacean species from the infraorders Astacidea, Thalassinidea, Polychelida, Palinura, and Anomura from the northern and northeastern (N/NE) Brazilian coast based on literature and material deposited in the carcinological collection of the Universidade Federal de Pernambuco, Recife, Brazil, is provided. The list includes marine and estuarine species reported at least once to each of the studied area, including the oceanic islands and banks along the N/NE Brazil. A total of 146 species is reported, corresponding to an increase of 32.7% when compared to the data published in Paulo Young's Catalogue (1998). The most representative infraorder concerning number of species is Anomura, represented in N/NE Brazil by 90 species and 10 families, followed by Thalassinidea, with 36 species and 6 families, Palinura, with 14 species and 4 families and, finally, Astacidea, which comprises 6 species and 2 families. Families with highest species richness were Porcellanidae (20), Diogenidae (19), Paguridae (18) and Galatheidae (15), all of them included in the infraorder Anomura. Zoogeographic affinities regarding the species are briefly discussed.

Key words: species richness; geographic distribution; crustaceans; northern and northeastern Brazil

Introduction

Reliable lists of species occurring in distinct geographic regions of the world are of multiple uses. In addition to providing comparative figures for biodiversity studies, they serve as an important tool in defining the extension of protected areas, inferring potential impact of anthropogenic activity and complexity of communities, and estimating availability of life resources (Hendrickx 1995).

The studies regarding the marine and estuarine decapod crustaceans from the northern and northeastern (N/NE) Brazilian coast started during the 19th century with the contributions of White (1847) and Smith (1869), among others. In the late 19th century, the Brazilian coast was visited by oceanographic expeditions of high historical importance such as the *Challenger* (Miers 1886; Bate 1888; Henderson 1888) and *Branner-Agassiz* (Rathbun 1900). In the 20th century, the R/V *Calypso* carried out samplings between 1961 and 1962 (Haig 1966; Forest & de Saint-Laurent 1967; Rodrigues da Costa 1968; Christoffersen 1979). Collections by the ships *Canopus* and *Akaroa*, between 1967 and 1969, and the *Almirante Saldanha*, between 1967 and 1987 were the basis for studies of Coelho & Koenig (1972), Coelho & Ramos (1972), Coelho *et al.* (1980). Additional information about the Brazilian oceanographic expeditions can be found in Coelho *et al.* (2004).

More recently, in the 1990's, a large number of crustaceans were collected during the activities of the Project "Recursos Vivos da Zona Econômica Exclusiva Brasileira" (REVIZEE) (Ramos-Porto *et al.* 2000a, 2000b; Cabral *et al.* 2000; Silva *et al.* 2002a, 2002b; Cardoso & Serejo 2003; Ramos-Porto *et al.* 2003; Komai 2004; Cardoso & Young 2005; Coelho Filho 2006). With the publication of data from the REVIZEE project and the registration, in the last years, of a number of new species, families, and superfamilies, we consider this an opportune moment for the organization of an updated list of species, incorporating most recent data and attempting a comparison between these data and those in the Young's Catalogue (1998).

The goal of this study is to provide an updated list of the marine and estuarine species concerning the decapod infraorders Astacidea, Thalassinidea, Polychelida, Palinura, and Anomura from N/NE Brazilian coast. Zoogeographic affinities regarding the species are briefly discussed.

Material and methods

Study area

Study area ranges from Cape Orange, northern Amapá State (04°17'N, 51°32'W) to Abrolhos Archipelago, southern Bahia State (18°19'S, 39°40'W), and towards the east to the St. Paul's Rocks (0°55'N, 29°20'W), which, along the Rocas Atoll (3°45'S and 3°56'S, 33°37'W and 33°56'W), seamounts of the North chains (01°00'S and 04°00'S, 37°00'W and 39°00'W), and Fernando de Noronha (03°00'S and 4°30'S, 32°00'W and 37°00'W), were considered within a single category denominated oceanic islands and banks. This area is included in the N/NE Brazilian Political Regions (Fig. 1).

The region undergoes direct influence of two main currents. The South Equatorial Current splits into two branches near 10° S, and continues towards the northwest into the North Brazilian Current. Another branch turns southwards as the beginning of the Brazil Current (Stramma *et al.* 1990). The northeast part of the continent has a semi-arid climate resulting in reduced supply of terrigenous material to the shelf. This leads to stable conditions of salinity, temperature, and water transparency, favorable to the growth of vegetal life, as calcareous algae, and hermatypical corals, to depths of almost 100 m (Mabesoone *et al.* 1972).

The northern South America has been divided by several authors in zoogeographic provinces (e.g., Dana 1853; Balech 1951; Coelho 1969; Coelho & Ramos 1972; Briggs 1974; Coelho *et al.* 1978, 1980; Boschi 2000). In the present report, we have adopted the classification proposal by Coelho & Ramos (1972), which divided the region in two provinces: Guyanas and Brazilian. The former extends from the Orinoco River delta, Venezuela, to Maranhão State, Brazil, and is mainly characterized by muddy and sandy bottoms and by the influence of equatorial rivers (e.g., Orinoco, Amazon, and Tocantins). The latter extends from Maranhão to Rio de Janeiro State, Brazil, being characterized, in a major part, by calcareous algae bottom and the river discharge does not present an important influence for coastal habitats (Coelho 1969; Coelho & Ramos 1972).

Method procedure

Records in the present contribution were derived from the review of literature dealing with the decapod crustaceans from the north and northeast Brazil (e.g., Smith 1869; Bate 1888; Henderson 1888; Pocock 1890; Rathbun 1900; Forest & de Saint Laurent 1967; Coelho & Ramos 1972; Young 1998; Melo 1999) and from specimens deposited in the crustacean collection at the Departamento de Oceanografia da Universidade Federal de Pernambuco (DOCEAN). A data bank stores the entire acquired information including all the consulted references from the last decades to the present moment. The list of species is presented in a table. References to the first record and locality, and distribution by state from northern and northeastern Brazilian regions are also provided.

Classification of the species in families and higher taxa are based on Martin & Davis' (2001) proposal. However, this classification differs in some aspects from the one in Catalogue of Crustacea of Brazil (Young

1998). To make it viable to compare the numerical data from 1998 and the current, we adapted the classification of the Catalogue to the proposal in Martin & Davis (2001).

The following abbreviations are used: States of North Region: Amapá (AP) and Pará (PA); States of Northeast Region: Maranhão (MA), Piauí (PI), Ceará (CE), Rio Grande do Norte (RN), Paraíba (PB), Pernambuco (PE), Alagoas (AL), Sergipe (SE), Bahia (BA), oceanic islands and banks (IB).



FIGURE 1. The study area. Abbreviations: Amapá (AP), Pará (PA), Maranhão (MA), Piauí (PI), Ceará (CE), Rio Grande do Norte (RN), Paraíba (PB), Pernambuco (PE), Alagoas (AL), Sergipe (SE), and Bahia (BA).

Results

A list of reported species in the study area, including the species distribution by state, is shown in Tables 1–4.

The largest infraorder concerning number of species is Anomura, represented in N/NE Brazil by 90 species and 10 families, followed by Thalassinidea, with 36 species and 6 families, Palinura, with 14 species and 4 families, Astacidea, which comprises 6 species and 2 families, and, finally, Polychelida, with 2 species and 1 family. Families with highest species richness were Porcellanidae (20), Diogenidae (19), Paguridae (18) and Galatheidae (15), all of them included in the infraorder Anomura.

Occurrence of the infraorder Astacidea in the N/NE Brazil was not mentioned prior to 1998 (Tavares 1998). Recently, Tavares & Young (2002) and Silva *et al.* (2003) have reported five species of the family Nephropidae. The family Enoplometopidae, although known to northeast Brazil since the contributions made by Fausto Filho (1970) and Coelho & Ramos (1972), was not cited by Tavares (1998) among the marine Brazilian astacideans.

TABLE 1. Species of the infraorder Astacidea Latreille, 1802 from north and northeast Brazilian coast. Abbreviations: Amapá (AP), Pará (PA), Maranhão (MA), Piauí (PI), Ceará (CE), Rio Grande do Norte (RN), Paraíba (PB), Pernambuco (PE), Alagoas (AL), Sergipe (SE), Bahia (BA), Oceanic Islands and Banks (IB). (X) first record, (+) further record (s).

Taxa	First record	AP	PA	MA	PI	CE	RN	PB	PE	AL	SE	BA	IB
Superfamily Enoplometopidea de Saint Laurent, 1988													
Family Enoplometopidae de Saint Laurent, 1988													
<i>Enoplometopus antillensis</i> Lütken, 1865	Fausto Filho (1970)	-	-	-	-	+	X	-	+	-	-	-	+
Superfamily Nephropoidea Dana, 1852													
Family Nephropidae Dana, 1852													
Subfamily Neophoberinae Glaessner, 1969													
<i>Acanthacaris caeca</i> (A. Milne Edwards, 1881)	Tavares & Young (2002)	+	-	-	-	-	-	-	-	-	-	X	-
<i>Nephropsis aculeata</i> Smith, 1881	Silva <i>et al.</i> (2003)	-	X	-	-	-	-	-	-	-	-	-	-
<i>Nephropsis agassizii</i> A. Milne Edwards, 1880	Tavares & Young (2002)	-	-	-	-	-	-	-	-	-	-	X	-
<i>Nephropsis neglecta</i> Holthuis, 1975	Tavares & Young (2002)	-	-	-	-	-	-	-	-	-	-	X	-
<i>Nephropsis rosea</i> Bate, 1888	Silva <i>et al.</i> (2003)	X	-	-	-	-	-	-	-	-	-	-	-

Regarding the infraorder Thalassinidea, 13 species have been recorded since the work carried out by Rodrigues & Shimizu (1998). These authors have recorded six species of the family Callianassidae from N/NE Brazil, while in the present checklist the number of callianassid species is increased to 12. According to Coelho, P.A. and Ramos-Porto, M. (2006, pers. com.), *Calastacus angulatus* Coelho, 1973 and *C. spinosus* Coelho, 1973 should be known as *Acanthaxius angulata* (Coelho, 1973) and *Calaxius spinosus* (Coelho, 1973), respectively, both assigned to the family Axiidae.

With respect to the superfamily Palinuroidea, among the 13 species recorded from Brazil by Coelho & Ramos-Porto (1998), 12 have been reported in N/NE Brazil. Since then, no additional records of palinuroids have been reported from the study area. Records of the infraorder Polychelida are recent and restricted to the present moment to Pará State (Ramos-Porto *et al.* 2000b; Silva *et al.* 2003). This material was collected during the activities of the REVIZEE program in the northern Brazilian coast.

Regarding the Anomura, a remarkable increase of 17 species has been registered since 1998 (Calado 1998; Melo-Filho 1998; Rieger 1998; Veloso 1998). The family Lithodidae is currently represented by 4 species in the Brazilian N/NE (Melo 1999; Serejo *et al.* 2006).

TABLE 2. Species of the infraorder Thalassinidea Latreille, 1831 from north and northeast Brazilian coast. Abbreviations: Amapá (AP), Pará (PA), Maranhão (MA), Piauí (PI), Ceará (CE), Rio Grande do Norte (RN), Paraíba (PB), Pernambuco (PE), Alagoas (AL), Sergipe (SE), Bahia (BA), Oceanic Islands and Banks (IB). (X) First record, (+) further record (s).

Taxa	First record	AP	PA	MA	PI	CE	RN	PB	PE	AL	SE	BA	IB
Superfamily Callianassoidea Dana, 1852													
Family Callianassidae Dana, 1852													
Subfamily Callianassinae Dana, 1852													
<i>Biffarius biformis</i> (Biffar, 1971)	Coelho (1997)	-	-	-	-	-	-	-	X	-	-	-	-
<i>Biffarius fragilis</i> (Biffar, 1971)	Coelho (1997)	-	-	-	-	-	-	-	X	-	-	-	-
Subfamily Callichirinae Manning & Felder, 1991													
<i>Callichirus major</i> (Say, 1818)	Coelho & Coelho-Santos (1993)	-	-	-	-	+	-	-	X	+	-	-	-
<i>Corallianassa hartmeyeri</i> (Schmitt, 1935)	Coelho (1997)	-	-	-	-	-	-	-	-	X	-	-	-
<i>Corallianassa longiventris</i> (A. Milne Edwards, 1870)	Coelho (1997)	-	X	-	-	-	-	-	-	-	-	-	X
<i>Lepidophthalmus siriboia</i> Felder & Rodrigues, 1993	Rodrigues (1971), as <i>Callianassa</i> (<i>Callichirus</i>) <i>jamaicensis</i>	-	X	+	-	-	-	+	+	+	-	-	X
<i>Neocallichirus grandimana</i> (Gibbes, 1850)	Rathbun (1900), as <i>Glypturus branneri</i>	-	-	-	-	+	-	X	+	-	-	-	-
<i>Neocallichirus maryae</i> Karasawa, 2004	Botter-Carvalho et al. (1995), as <i>N. rathbunae</i>	-	-	-	-	-	-	-	X	-	-	-	-
<i>Sergio guara</i> (Rodrigues, 1971)	Coelho (1969), as <i>Callianassa guara</i>	-	+	X	-	+	-	-	+	+	-	-	-
<i>Sergio guassatinga</i> (Rodrigues, 1971)	Rodrigues (1971), as <i>Callianassa</i> (<i>Callichirus</i>) <i>guassatinga</i>	-	-	-	-	-	-	+	+	-	+	-	X
<i>Sergio mirim</i> (Rodrigues, 1971)	Rodrigues (1971), as <i>Callianassa</i> (<i>Callichirus</i>) <i>mirim</i>	-	-	-	-	-	-	-	-	-	-	-	X
Subfamily Cheraminae Manning & Felder, 1991													
<i>Cheramus marginatus</i> (Rathbun, 1901)	Coelho & Ramos (1972)	+	-	X	+	+	-	-	-	+	-	-	-
Family Ctenochelidae Manning & Felder, 1991													
Subfamily Ctenochelinae Manning & Felder, 1991													
<i>Ctenocheles holthuisi</i> Rodrigues, 1978	Coelho & Ramos (1972), as <i>Ctenocheles</i> sp.	-	-	-	-	-	-	-	-	X	-	-	-
Subfamily Eucalliinae Manning & Felder, 1991													
<i>Eucalliax cearensis</i> Rodrigues & Manning, 1992	Rodrigues & Manning (1992)	-	-	-	-	X	-	-	-	-	-	-	-
Subfamily Gourretinae Sakai, 1999													
<i>Dawsonius latispina</i> (Dawson, 1967)	Coelho (1997), as <i>Callianassa latispina</i>	X	X	-	-	-	-	-	-	X	-	-	-
Family Laomedidae Borradaile, 1903													
<i>Axianassa australis</i> Rodrigues & Shimizu, 1992	Rodrigues & Shimizu (1992)	-	-	-	-	-	-	-	+	-	-	-	X

to be continued.

TABLE 2. (continued)

Taxa	First record	AP	PA	MA	PI	CE	RN	PB	PE	AL	SE	BA	IB
Family Upogebidae Borradaile, 1903													
<i>Pomatogebia operculata</i> (Schmitt, 1924)	Coelho (1969), as <i>Upogebia (Calliadne) operculata</i>	-	-	-	-	X	+	-	+	-	-	+	-
<i>Upogebia acanthura</i> Coelho, 1973	Coelho & Ramos-Porto (1987)	-	X	-	-	+	-	-	X	-	-	-	+
<i>Upogebia brasiliensis</i> Holthuis, 1956	Gomes Corrêa (1968a)	-	X	X	-	-	+	+	+	-	-	+	-
<i>Upogebia careospina</i> Williams, 1993	Williams (1993)	-	-	-	-	X	-	-	-	-	-	-	-
<i>Upogebia marina</i> Coelho, 1973	Coelho (1973)	-	+	-	+	+	X	+	+	+	+	-	-
<i>Upogebia noronhensis</i> Fausto Filho, 1969	Pocock (1890), as <i>Gebia spinigera</i>	-	-	+	-	-	+	-	+	+	-	-	X
<i>Upogebia omissa</i> Corrêa, 1968	Coelho (1966a), as <i>Upogebia</i> sp.	-	-	+	-	+	+	+	X	+	+	+	-
<i>Upogebia omissago</i> Williams, 1993	Williams (1993)	-	-	-	X	-	-	-	-	-	-	-	-
<i>Upogebia paraffinis</i> Williams, 1993	Rathbun (1900), as <i>Upogebia affinis</i>	-	-	-	-	+	-	X	-	X	-	-	-
<i>Upogebia vasquezii</i> Ngoc-Ho, 1989	Williams (1993)	-	-	+	-	+	X	X	-	-	-	X	-
Superfamily Axioidae Huxley, 1879													
Family Axiidae Huxley, 1879													
Subfamily Axinae Huxley, 1879													
<i>Axiopsis brasiliensis</i> Coelho & Ramos-Porto, 1985	Coelho & Ramos (1972), as <i>A. (Axiopsis)</i> sp.	-	-	-	-	X	X	-	+	+	-	-	+
<i>Axiopsis serratifrons</i> (A. Milne Edwards, 1873)	Rodrigues & Kensley (1991)	-	-	-	-	-	-	X	-	-	-	X	-
<i>Axiorygma nethertoni</i> Kensley & Simmons Jr., 1988	Coelho & Ramos-Porto (1985), as <i>Axiopsis (Axiopsis)</i> sp.	X	-	-	-	-	-	-	-	-	-	-	-
<i>Paraxiopsis defensa</i> (Rathbun, 1901)	Coelho & Ramos-Porto (1985), as <i>Axiopsis (Paraxiopsis) defensa</i>	-	-	-	-	-	-	-	X	-	-	-	-
<i>Paraxiopsis vicina</i> (Coelho & Ramos-Porto, 1985)	Coelho & Ramos-Porto (1985), as <i>Axiopsis (Paraxiopsis)</i> sp. A	-	-	-	-	X	-	-	-	-	-	-	-
Subfamily Coralaxiinae Sakai, 1989													
<i>Coralaxius nodulosus</i> (Meinert, 1987)	Coelho & Ramos-Porto (1985), as <i>C. abelei</i>	-	-	-	-	-	-	-	-	-	-	X	X
Family Calocardiidae Ortman, 1891													
<i>Acanthaxius angulatus</i> Coelho, 1973	Coelho (1973), as <i>Calastacus angulatus</i>	X	-	-	-	-	-	-	-	-	-	-	-
<i>Calaxius spinosus</i> Coelho, 1973	Coelho (1969), as <i>Axiopsis</i> sp. C	X	-	-	-	-	-	-	-	-	-	-	-
Family Micheleidae Sakai, 1992													
<i>Marcusitaxius lemoscastroi</i> Rodrigues & Carvalho, 1972	Rodrigues & Carvalho (1972)	X	+	-	-	-	-	-	-	-	-	-	-
<i>Marcusitaxius minutus</i> (Coelho, 1973)	Coelho (1973), as <i>Metriconaxius minutus</i>	X	-	+	-	-	-	-	-	-	-	-	-

TABLE 3. Species of the infraorders Polychelida de Haan 1841 and Palinura Latreille, 1802 from north and northeast Brazilian coast. Abbreviations: Amapá (AP), Pará (PA), Maranhão (MA), Piauí (PI), Ceará (CE), Rio Grande do Norte (RN), Paraíba (PB), Pernambuco (PE), Alagoas (AL), Sergipe (SE), Bahia (BA), Oceanic Islands and Banks (IB). (X) First record, (+) further record (s).

Taxa	First record	AP	PA	MA	PI	CE	RN	PB	PE	AL	SE	BA	IB
Infraorder Polychelida de Haan 1841													
Superfamily Eryonoidea de Haan, 1841													
Family Polychelidae Wood-Mason, 1875													
<i>Polycheles typhlops</i> Heller, 1862	Silva <i>et al.</i> (2003)	-	X	-	-	-	-	-	-	-	-	-	-
<i>Polycheles sculptus</i> Smith, 1880	Ramos-Porto <i>et al.</i> (2000), as <i>Stereomastis sculpta</i>	-	X	-	-	-	-	-	-	-	-	-	-
Infraorder Palinura Latreille, 1802													
Superfamily Palinuroidea Latreille, 1802													
Family Palinuroidea Latreille, 1802													
Family Palinuridae Latreille, 1803													
<i>Jusitii longimana</i> (Milne Edwards, 1837)	Coelho & Vasconcelos (1993)	-	-	-	-	-	X	-	-	-	-	-	+
<i>Palinustus truncatus</i> (Milne Edwards, 1880)	Fausto Filho (1977)	+	X	-	-	-	-	-	-	-	-	-	-
<i>Panulirus argus</i> (Latreille, 1804)	White (1847)	-	+	+	+	+	+	+	+	+	+	X	+
<i>Panulirus echinatus</i> (Smith, 1869)	Smith (1869)	-	-	-	-	X	+	+	+	+	-	+	+
<i>Panulirus laevicauda</i> (Latreille, 1817)	Pocock (1890), as <i>P. ornatus</i>	-	-	+	-	+	+	+	+	-	+	+	X
Family Scyllaridae Latreille, 1825													
Subfamily Ibacinæ Holthuis, 1985													
<i>Parribacus antarcticus</i> (Lund, 1793)	Matthews (1926)	+	+	-	-	+	+	-	X	+	-	+	-
Subfamily Scyllarinæ Latreille, 1825													
<i>Scyllarides brasiliensis</i> Rathbun, 1906	Smith (1869), as <i>Scyllarus aequinoxialis</i>	-	-	-	-	+	+	-	-	-	+	-	-
<i>Scyllarides delfosi</i> Holthuis, 1960	Fausto Filho <i>et al.</i> (1966)	+	+	-	-	X	-	-	-	-	-	-	-
<i>Scyllarus americanus</i> (Smith, 1869)	Bullis & Thompson (1965)	+	X	-	-	-	-	+	+	-	-	-	-
<i>Scyllarus chacei</i> Holthuis, 1960	Coelho (1966b)	-	+	-	-	+	+	+	+	X	X	+	-
<i>Scyllarus depressus</i> (Smith, 1881)	Coelho & Ramos-Porto (1998)	-	X	-	-	-	-	-	-	-	-	-	-
Family Synaxidae Bate, 1881													
<i>Palinurellus gundlachi</i> von Martens, 1878	Coelho (1969)	-	-	-	-	+	-	-	X	-	-	-	-

TABLE 4. Species of the infraorder Anomura MacLeay, 1838 from north and northeast Brazilian coast. Abbreviations: Amapá (AP), Pará (PA), Maranhão (MA), Piauí (PI), Ceará (CE), Rio Grande do Norte (RN), Paraíba (PB), Pernambuco (PE), Alagoas (AL), Sergipe (SE), Bahia (BA), Oceanic Islands and Banks (IB). (X) First record, (+) further record (s).

Taxa	First record	AP	PA	MA	PI	CE	RN	PB	PE	AL	SE	BA	IB
Superfamily Galatheoidea Samouelle, 1819													
Family Chirostyliidae Ortmann, 1892													
<i>Uroptychus minutus</i> Benedict, 1902	Coelho (1969)	X	-	-	-	-	-	-	-	-	-	-	-
<i>Uroptychus nitidus</i> (A. Milne Edwards, 1880)	Melo (1999)	-	-	-	-	-	-	-	X	-	-	-	-
<i>Uroptychus unceifer</i> (A. Milne Edwards, 1880)	Bullis & Thompson (1965)	-	X	-	-	-	-	-	-	-	-	-	-
Family Galatheidae Samouelle, 1819													
<i>Munida angulata</i> Benedict, 1902	Coelho & Ramos-Porto (1995)	-	-	+	-	+	-	-	X	-	-	-	+
<i>Munida atlantica</i> Melo-Filho & Melo, 1994	Melo-Filho & Melo (1994)	-	-	-	-	+	-	-	-	-	-	-	-
<i>Munida constricta</i> A. Milne Edwards, 1880	Henderson (1888), as <i>M. miles</i> (in part)	-	-	-	-	-	-	-	-	X	-	-	-
<i>Munida flinti</i> Benedict, 1902	Henderson (1888), as <i>M. stimpsoni</i>	-	-	-	-	-	-	-	-	X	-	-	-
<i>Munida forceps</i> A. Milne Edwards, 1880	Henderson (1888), as <i>M. miles</i> (in part)	-	-	-	-	-	-	-	-	X	-	-	-
<i>Munida iris</i> A. Milne Edwards, 1880	Coelho <i>et al.</i> (1983)	+	+	+	-	-	-	-	-	X	-	-	-
<i>Munida irrasa</i> A. Milne Edwards, 1880	Melo-Filho & Melo (2001)	X	X	X	-	-	-	-	-	-	-	-	-
<i>Munida petronioi</i> Melo-Filho & Melo, 1994	Melo-Filho & Melo (1994)	-	-	-	-	-	+	-	-	-	-	-	-
<i>Munida pusilla</i> Benedict, 1902	Melo-Filho & Melo (2001)	X	-	-	-	+	-	-	-	-	-	-	-
<i>Munida sanctipauli</i> Henderson, 1885	Henderson (1888)	-	-	-	-	-	-	-	-	-	-	-	X
<i>Munida simplex</i> Benedict, 1903	Calado (1996)	-	-	X	-	-	-	-	-	-	-	-	-
<i>Munida spinifrons</i> Henderson, 1885	Henderson (1888)	+	-	-	-	+	-	-	+	-	-	-	X
<i>Munida valida</i> Smith, 1883	Henderson (1888), as <i>M. miles</i> (in part)	+	-	-	-	-	-	-	-	X	-	-	-
<i>Munidopsis barbarae</i> (Boone, 1927)	Coelho & Ramos-Porto (1995)	-	-	-	-	-	-	-	X	-	-	-	-
<i>Munidopsis erinacea</i> (A. Milne Edwards, 1880)	Henderson (1888)	-	-	-	-	-	-	-	+	X	-	-	-
Family Porcellanidae Haworth, 1825													
<i>Megalobrachium mortenseni</i> Haig, 1962	Coelho (1969)	-	X	-	X	-	+	+	+	+	-	-	+
<i>Megalobrachium poeyi</i> (Guérin, 1855)	Calado (1996)	-	-	-	-	-	-	-	X	-	-	-	-
<i>Megalobrachium roseum</i> (Rathbun, 1900)	Rathbun (1900), as <i>Porcellana rosea</i>	-	-	+	-	+	+	X	X	+	-	+	-
<i>Megalobrachium soriatum</i> (Say, 1818)	Haig (1966)	-	-	-	-	+	-	+	+	X	-	+	-
<i>Minyocerus angustus</i> (Dana, 1852)	Rathbun (1900)	-	+	+	-	+	+	X	+	+	+	+	-
<i>Pachycheles ackleyanus</i> A. Milne Edwards, 1880	Coelho (1964a)	-	-	+	-	+	+	+	X	+	+	+	-
<i>Pachycheles chacei</i> Haig, 1956	Melo (1999)	-	-	-	-	-	-	-	-	X	-	-	-
<i>Pachycheles greeleyi</i> (Rathbun, 1900)	Rathbun (1900), as <i>Pisosoma riisei</i>	-	+	+	+	+	+	+	+	X	-	+	-
<i>Pachycheles laevidactylus</i> Ortmann, 1892	Rodrigues da Costa (1960), as <i>P. haigae</i>	-	-	+	-	-	-	-	X	-	-	+	-
<i>Pachycheles monilifer</i> (Dana, 1852)	Cano (1889), as <i>P. moniliferus</i>	-	-	-	-	-	+	+	X	+	-	+	-
<i>Pachycheles riisei</i> (Stimpson, 1858)	Rathbun (1900), as <i>Pisosoma riisei</i>	-	-	-	-	-	-	+	+	X	-	+	-

to be continued..

TABLE 4. (continued)

Taxa	First record	AP	PA	MA	PI	CE	RN	PB	PE	AL	SE	BA	IB
<i>Pachycheles rugimanus</i> A. Milne Edwards, 1880	Coelho (1964a)	+	-	-	-	-	-	+	X	-	-	-	-
<i>Petrolisthes amoenus</i> (Guérin, 1855)	Henderson (1888), as <i>P. serratus</i>	-	-	-	-	+	+	+	+	+	-	X	+
<i>Petrolisthes armatus</i> (Gibbes, 1850)	Pocock (1890), as <i>P. marginatus</i>	-	+	-	-	+	+	+	+	+	-	+	X
<i>Petrolisthes costai</i> Haig, 1968	Haig (1968) in Rodrigues da Costa, 1968	-	-	-	-	-	-	-	-	-	-	X	-
<i>Petrolisthes galathinus</i> (Bosc, 1802)	Cano (1889), as <i>P. brasiliensis</i>	-	+	+	+	+	+	+	X	+	+	+	-
<i>Petrolisthes marginatus</i> Stimpson, 1859	Pocock (1890)	-	-	+	-	-	-	-	-	-	-	-	X
<i>Petrolisthes rosariensis</i> Werding, 1978	Coelho <i>et al.</i> (1990)	-	-	-	-	-	-	+	+	X	-	+	-
<i>Pisidia brasiliensis</i> Haig, 1968	Coelho (1964a), as <i>Megalobrachium poeyi</i>	-	+	+	-	+	+	X	+	-	+	+	-
<i>Polyonyx gibbesi</i> Haig, 1956	Veloso & Melo (1993)	-	-	-	-	X	-	-	X	-	-	X	-
<i>Porcellana sayana</i> (Leach, 1820)	Coelho (1964a)	+	+	-	-	+	+	+	X	+	-	+	-
<i>Porcellana sigsbeiana</i> A. Milne Edwards, 1880	Veloso & Melo (1993)	-	X	-	-	-	-	-	-	-	-	-	-
Superfamily Hippoidea Latreille, 1825													
Family Alboneidae Stimpson, 1858													
<i>Albonea gibbesii</i> Stimpson, 1859	Coelho & Ramos (1972)	-	-	-	-	-	-	-	+	X	-	-	-
<i>Albonea paretii</i> Guérin-Méneville, 1853	Coelho (1969)	+	X	X	+	+	+	+	-	-	-	+	-
<i>Lepidopa distincta</i> Gomes Corrêa, 1968	Fausto Filho (1968); Gomes Corrêa (1968b)	-	-	-	X	-	-	-	+	X	-	-	-
<i>Lepidopa richmondi</i> Benedict, 1903	Fausto Filho (1979)	-	-	-	-	X	+	+	-	-	-	-	-
<i>Lepidopa venusta</i> Stimpson, 1860	Melo (1999)	-	-	-	-	-	-	-	-	-	-	X	-
<i>Zygopa michaelis</i> Holthuis, 1960	Calado <i>et al.</i> (1990)	-	-	X	-	X	-	-	-	-	-	-	-
Family Hippidae Latreille, 1825													
<i>Emerita portoricensis</i> Schmitt, 1935	Rodrigues da Costa (1962)	-	-	+	-	X	+	+	+	-	+	+	-
<i>Hippa testudinaria</i> (Herbst, 1791)	Pocock (1890), as <i>Remipes scutellatus</i>	-	-	-	-	+	+	+	+	-	-	+	X
Superfamily Paguroidea Latreille, 1802													
Family Diogenidae Ortmann, 1892													
<i>Calcinus tibicen</i> (Herbst, 1791)	Smith (1869), as <i>C. sulcatus</i>	-	-	-	-	+	+	+	+	+	-	X	+
<i>Cancelanus ornatus</i> Benedict, 1901	Forest & de Saint Laurent (1967)	-	-	-	-	-	-	-	+	-	-	X	-
<i>Cibbanarius antillensis</i> Stimpson, 1859	Smith (1869)	-	-	-	+	+	+	+	+	+	-	X	+
<i>Cibbanarius foresti</i> Holthuis, 1959	Coelho (1969)	-	-	-	-	-	-	-	-	-	-	-	-
<i>Cibbanarius scopetarius</i> (Herbst, 1796)	Smith (1869)	-	-	-	+	+	+	+	+	-	-	X	-
<i>Cibbanarius tricolor</i> (Gibbes, 1850)	Coelho (1964b)	-	-	-	-	-	-	-	-	-	-	-	X
<i>Cibbanarius vitatus</i> (Bosc, 1802)	Smith (1869)	-	-	+	-	+	+	+	+	+	+	X	-
<i>Dardanus fucosus</i> Biffar & Provenzano, 1972	Coelho (1969), as <i>Dardanus</i> sp.	-	+	+	-	+	+	+	+	+	-	-	-
<i>Dardanus venosus</i> (Milne Edwards, 1848)	Rathbun (1900), as <i>Petrochirus insignis</i>	-	+	+	+	+	+	+	X	X	+	+	+
<i>Isocheles sawayai</i> Forest & de Saint Laurent, 1967	Fausto Filho (1966), as <i>I. wurdemanni</i>	-	-	-	-	X	-	-	+	+	-	-	-
<i>Paguristes angustitheca</i> McLaughlin & Provenzano, 1974	McLaughlin & Provenzano (1974)	X	-	-	-	-	-	-	-	-	-	-	-
<i>Paguristes callioptis</i> Forest & de Saint Laurent, 1967	Forest & de Saint Laurent (1967)	-	-	-	-	-	-	-	-	-	-	-	X

to be continued.

TABLE 4. (continued)

Taxa	First record	AP	PA	MA	PI	CE	RN	PB	PE	AL	SE	BA	IB
<i>Paguristes erythrops</i> A. Milne Edwards, 1880	Forest & de Saint Laurent (1967)	-	-	+	+	-	+	+	X	X	X	X	-
<i>Paguristes oxyphthalms</i> Holthuis, 1959	Calado (1996)	-	X	-	-	-	-	-	-	-	-	-	-
<i>Paguristes perplexus</i> McLaughlin & Provenzano, 1974	Forest & de Saint Laurent (1967), as <i>P. tortugae</i>	-	-	-	-	+	-	-	X	-	-	X	-
<i>Paguristes spinipes</i> A. Milne Edwards, 1880	Henderson (1888), as <i>P. visor</i>	-	-	+	+	+	+	+	-	X	-	-	-
<i>Paguristes tortugae</i> Schmitt, 1933	McLaughlin & Provenzano (1974)	-	X	-	-	+	-	-	-	-	-	-	+
<i>Paguristes triangulopsis</i> Forest & de Saint Laurent, 1967	Forest & de Saint Laurent (1967)	-	-	-	-	-	-	-	-	-	X	X	-
<i>Petrochirus diogenes</i> (Linnaeus, 1758)	Smith (1869), as <i>P. granulatus</i>	+	-	+	-	+	+	+	+	+	+	+	X
Family Lithodidae Samouelle, 1819													
<i>Lithodes mamingi</i> Macpherson, 1988	Young & Serejo (2002)	-	-	-	-	-	-	-	-	-	-	-	X
<i>Neolithodes agassizii</i> (Smith, 1882)	Young & Serejo (2002)	-	-	-	-	-	-	-	-	-	-	-	X
<i>Paralomis cubensis</i> Chace, 1939	Melo (1999)	-	X	-	-	-	-	-	-	-	-	-	-
<i>Paralomis formosa</i> Henderson, 1888	Young & Serejo (2002)	-	-	-	-	-	-	-	-	-	-	-	X
Family Paguridae Latreille, 1803													
<i>Agaricochirus gibbosimanus</i> (A. Milne Edwards, 1880)	Bullis & Thompson (1965)	-	X	-	-	-	-	-	-	-	-	-	-
<i>Anisopagurus bartlettii</i> (A. Milne Edwards, 1880)	Bullis & Thompson (1965)	-	X	-	-	-	-	-	-	-	-	-	-
<i>Iridopagurus dispar</i> (Stimpson, 1859)	Coelho <i>et al.</i> (1983)	-	-	-	-	-	-	-	-	X	-	-	-
<i>Iridopagurus iris</i> (A. Milne Edwards, 1880)	Coelho & Ramos-Porto (1986)	X	-	-	-	-	-	-	-	-	-	-	-
<i>Iridopagurus reticulatus</i> Garcia-Goméz, 1983	Coelho <i>et al.</i> (1990)	-	-	-	-	-	-	-	-	X	-	-	-
<i>Iridopagurus violaceus</i> de Saint Laurent, 1966	Forest & de Saint Laurent (1967)	-	+	+	+	+	+	+	X	+	X	X	X
<i>Nematopaguroides fagei</i> Forest & de Saint Laurent, 1967	Forest & de Saint Laurent (1967)	-	-	-	-	-	-	-	X	-	-	+	-
<i>Nematopaguroides? pusillus</i> Forest & de S. Laurent, 1967	Forest & de Saint Laurent (1967)	-	-	-	-	-	-	-	X	-	-	-	-
<i>Pagurus brevidactylus</i> (Stimpson, 1859)	Coelho (1964b)	-	-	-	-	-	-	-	X	+	+	+	+
<i>Pagurus criniticornis</i> (Dana, 1852)	Forest & de Saint Laurent (1967)	-	-	-	-	+	+	+	+	+	+	X	+
<i>Pagurus leptonyx</i> Forest & de Saint Laurent, 1967	Fausto Filho (1970)	-	-	-	-	X	-	-	-	+	+	-	-
<i>Pagurus limatulus</i> Fausto Filho, 1970	Fausto Filho (1970)	-	-	-	-	X	+	-	-	-	-	-	-
<i>Pagurus longimanus</i> Wass, 1963	Coelho (1969)	X	-	-	-	-	-	-	-	-	-	-	-
<i>Pagurus provenzanoi</i> Forest & de Saint Laurent, 1967	Forest & de Saint Laurent (1967)	+	-	-	-	+	+	+	X	+	+	X	X
<i>Phimochirus holthuisi</i> (Provenzano, 1961)	Coelho & Ramos (1972)	-	X	-	-	-	-	-	+	-	-	+	-
<i>Phimochirus ocellus</i> (Henderson, 1888)	Henderson (1888)	-	-	-	-	-	-	-	X	-	-	-	-
<i>Pylopagurus discoidalis</i> (A. Milne Edwards, 1880)	Bullis & Thompson (1965)	+	X	-	-	-	-	-	-	-	-	-	-
<i>Tomopagurus wassi</i> McLaughlin, 1981	McLaughlin (1981)	-	-	X	-	-	-	-	-	-	-	-	-
Family Parapaguridae Smith, 1882													
<i>Oncopagurus bicristatus</i> (A. Milne Edwards, 1880)	Lemaitre (1989), as <i>Sympagurus bicristatus</i>	X	-	X	-	-	-	-	-	-	-	-	-
<i>Oncopagurus gracilis</i> (Henderson, 1888)	Henderson (1888), as <i>Parapagurus gracilis</i>	-	-	+	-	-	-	-	X	-	-	-	-
Family Pylochelidae Bate, 1888													
<i>Mixtopagurus paradoxus</i> A. Milne Edwards, 1880	Bullis & Thompson (1965)	+	X	-	-	-	-	-	-	-	-	-	-

Discussion

In Paulo S. Young's Catalogue of Crustacea of Brazil (Young 1998), a total of 140 species belonging to the studied groups are cited from Brazil, with 110 assigned to the northern and/or northeastern Brazilian coast. In the present contribution we report a total of 146 species, which amounts to an increase of 32.7% since 1998. Regarding the higher taxa, the number of families and superfamilies in this portion of the Brazilian coast rose from 6 to 9 and from 18 to 23, respectively, between 1998 and 2006. The bulk of new records for the study area can be attributed to the material collected by the REVIZEE project (Tavares & Young 2002, Ramos-Porto *et al.* 2000b, Serejo *et al.* 2006; Silva *et al.* 2003).

Boschi (2000), studying the decapod crustacean distribution in the South American marine zoogeographic provinces, recorded a total of 153 species between the Orinoco River delta (Venezuela, 08°56'N, 60°47'W) and Cape Frio (Brazil, 22°53'S, 42°02'W). This zoogeographic area is said to comprise the Brazilian Province, a more extensive area than the N/NE Brazilian coast, but less representative, since 146 species are reported in this contribution. Campos *et al.* (2003) mentioned 172 species from the Caribbean coast of Colombia, an area with a high diversity of coastal habitats, but is shorter than the N/NE Brazil coast. With these data, it is possible to conclude that the number of species from this part of the Brazilian littoral zone is still poorly known.

Analysis of the currently known distribution of the species reported herein, based on Melo (1985), allows identification of four patterns of longitudinal distribution: western Atlantic species, Amphi-Atlantic species, Amphi-American species, and circum-tropical species.

The first group comprises species endemic to the western Atlantic, with 126 species. Most of the species have a wide latitudinal distribution, occurring along merely the entire coast of the Americas, spanning more than one zoogeographic province, including tropical and temperate provinces. However, some species range from North America to the area under influence of the Amazon's Rivers (e.g., *Axiorygma nethertoni*, *Palinurus truncatus*, *Dardanus fucosus*, and *Paguristes angustithecica*). Yet, other species seem to have a disjunct distribution, occurring from North America to southern South America, but not in the area under influence of the Amazon's Rivers (e.g., *Albunea gibbesi*, *Calcinus tibicen*, *Pagurus brevidactylus*, and *Emerita portoricensis*). However, it is important to keep in mind that some of these species could be 'absent' due to the lack of collection effort.

Among the 126 species, 28 species have a distribution restricted to the Guyanas and/or Brazilian provinces. *Acanthaxius angulata*, *Calaxius spinosus*, *Marcusiarius minutus*, *Clibanarius foresti*, *Paguristes oxyphthalmus*, and *Pagurus longimanus* are endemic to the Guyanas province, an area characterized by soft bottoms and influenced by the Amazon's Rivers discharge. Another group is known to be endemic of the Brazilian province, *Ctenocheles holthuisi*, *Eucalliix cearensis*, *Axianassa australis*, *Upogebia careospina*, *U. noronhesis*, *U. omissago*, *U. paraffinis*, *Axiopsis brasiliensis*, *Paraxiopsis vicina*, *Munida atlantica*, *M. petronioi*, *Paguristes triangulopsis*, *Nematopaguroides fagei*, *N. pusillus*, and *Pagurus limatulus*. Among them, *A. australis*, *U. paraffinis*, and *N. fagei* ranges to the southeast Brazilian coast, which is considered by some authors as a region of peculiar hydrological features and a zoogeographical transition zone between tropical and temperate faunas (Coelho *et al.* 1978; Melo 1990; Melo-Filho 2006), recognized by several authors as the Paulista province (Coelho & Ramos 1972; Coelho *et al.* 1978; Palacio 1982). Finally, the species *Sergio guara*, *Scyllarides delfosi*, *Pachycheles greeleyi*, *Pisidia brasiliensis*, *Paguristes calliopsis*, *P. erythropis*, and *Iridopagurus violaceus* are common to the Guyanas and Brazilian provinces, although the distribution could be extended to the Caribbean or Paulista provinces.

The second group is formed by the astacid *Enoplometopus antillensis*, the palinurans *Panulirus argus* and *P. echinatus*, and the anomurans, *Munida iris*, *M. sanctipauli*, *Petrolisthes marginatus*, *Albunea paretii*, *Onco-pagurus bicristatus*, and *O. gracilis*, which are represented in both western and eastern Atlantic. The third group comprises the thalassinid *Neocallichirus grandimana* and the anomurans *Megalobrachium mortenseni*,

M. roseum, *M. soriatum*, *Pachycheles ackleianus*, *P. chacei*, and *P. monilifer* which are represented in both sides of South America.

Finally, the last group is composed of species with circum-tropical distribution, which include the thalassinid *Axiopsis serratifrons*, the palinuran *Parribacus antarcticus*, and the anomurans *Petrolisthes armatus* and *P. galathinus*. These porcellanids are widely distributed, but are absent from the Indo-West Pacific (Melo 1999). Both are morphologically variable *taxa*, believed to represent species complexes (Werding *et al.* 2003; Rodriguez *et al.* 2005).

Acknowledgements

P.A. Coelho would like to thank CNPq for a research productivity scholarship during the period of this study. J.F. Souza-Filho was supported by a master's scholarship (CNPq). L.E.A. Bezerra thanks PROPESQ/UFPE for the provision of a Ph.D scholarship. A.O. Almeida thanks FAPESB (Fundação de Amparo à Pesquisa do Estado da Bahia) for the provision of a research productivity scholarship (PP3). Special thanks to M.Sc. Paula C. Jimenez for revising the written English language.

References

- Balech, E. (1951) División zoogeográfica del litoral sudamericano. *Revista de Biología Marina*, 4, 184–195.
- Bate, C.S. (1888) Report of the Crustacea Macrura collected by the H.M.S. "Challenger" during the years 1873–1876. *Report on the Scientific Results of the Voyage of H. M. S. "Challenger"*, *Zoology*, 24, 1–942.
- Boschi, E.E. (2000) Species of decapod crustaceans and their distribution in the american marine zoogeographic provinces. *Revista de Investigación y Desarrollo Pesquero*, 13, 7–136.
- Botter-Carvalho, M., Carvalho, P.V.V.D.B.C & Ramos-Porto, M. (1995) Registro de *Neocallichirus rathbunae* (Schmitt, 1935) em águas do litoral brasileiro (Decapoda: Callianassidae). *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 23, 199–202. [Dated 1994/95, published 1995].
- Briggs, J.C. (1974) *Marine Zoogeography*. McGraw-Hill, New York, 475 pp.
- Bullis Jr., H.R. & Thompson, J.R. (1965) Collections by the exploratory fishing vessels "Oregon", "Silver Bay", "Combat", and "Pelican" made during 1956 to 1960 in the southwestern north Atlantic. *Special Scientific Report of the United States Fisheries Fish and Wildlife Service*, 510, 1–130.
- Cabral, E., Ramos-Porto, M., Santos, M.C.F., Acioli, F.D., Torres, M.F.A. & Viana, G.F.S. (2000) Shrimps collected in the Northeast of Brazil during the REVIZEE Program (Decapoda, Caridea). *Nauplius*, 8 (2), 245–248.
- Calado, T.C.S. (1996) Crustacea Decapoda Anomura e Stomatopoda da Comissão "Pavasas I" na costa Norte e Nordeste do Brasil. *Boletim de Estudos de Ciências do Mar*, 9, 149–174.
- Calado, T.C.S. (1998) Malacostraca. Eucarida. Hippoidea. In: Young, P.S. (Ed.), *Catalogue of Crustacea of Brazil*. Museu Nacional, Rio de Janeiro, pp. 407–411.
- Calado, T.C.S., Coelho, P.A. & Ramos-Porto, M. (1990). Crustáceos decápodos da superfamília Hippoidea na costa brasileira. *Arquivos de Biologia e Tecnologia*, 33 (4), 743–757.
- Campos, N.H., Lemaitre, R. & Navas, G.R. (2003) La fauna de crustáceos decápodos de la costa Caribe colombiana: un aporte al conocimiento de la biodiversidad en Colombia. In: G. Montañez Gómez (Ed.), *Mundo Marino en Colombia: Investigación y Desarrollo de Territorios Olvidados*. Red de estudios del mundo marino REMAR, Universidad Nacional de Colombia, Bogotá, 174–184.
- Cano, G. (1889) Crostacei brachiuri ed anomuri raccolti nel viaggio dela "Vettor Pisani" in torno al globo. *Bollettino della Societa dei Naturalisti in Napoli*, 1 (3), 79–268.
- Cardoso, I.A. & Serejo, C.S. (2003) Sergestidae (Crustacea, Dendrobranchiata) from the southwestern Atlantic, collected by the Revizee program. *Boletim do Museu Nacional, Nova Série Zoologia*, 512, 1–15.
- Cardoso, I.A. & Young, P. (2005) Deep-sea Oplophoridae (Crustacea Caridea) from the southwestern Brazil. *Zootaxa*, 1031, 1–76.
- Christoffersen, M.L. (1979) Decapod Crustacea: Alpheoidea. Campagne de la Calypso au large des côtes atlantiques de l'Amérique du Sud (1961–1962). I. 36, *Annales de l'Institut Océanographique, Monaco*, (Suppl.), 55, 297–377.
- Coelho, P.A. (1964a) Lista dos Porcellanidae (Crustacea, Decapoda, Anomura) do litoral de Pernambuco e dos Estados vizinhos. *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 5/6, 51–68. [Dated 1963/64, pub-

- lished 1964].
- Coelho, P.A. (1964b) Alguns crustáceos decápodos novos para Pernambuco e Estados vizinhos na Coleção Carcinológica do Instituto Oceanográfico da Universidade do Recife. *Ciência e Cultura*, 16 (2), 225–226.
- Coelho, P.A. (1966a) Os crustáceos decápodos de alguns manguezais pernambucanos. *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 7/8, 71–89. [Dated 1965/66, published 1966].
- Coelho, P.A. (1966b). Alguns crustáceos decápodos novos para Pernambuco e Estados vizinhos na Coleção Carcinológica do Instituto Oceanográfico da Universidade Federal de Pernambuco. Segunda lista. *Ciência e Cultura*, 18 (2), 139–140.
- Coelho, P.A. (1969) A distribuição dos crustáceos decápodos reptantes do Norte do Brasil. *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 9/11, 223–238. [Dated 1967/69, published 1969].
- Coelho, P.A. (1973) Descrição preliminar de *Calastacus angulatus*, n. sp., e de *C. spinosus*, n. sp., do norte do Brasil (Crustacea, Decapoda, Callianassidae). *Ciência e Cultura*, 25 (6) (Supl.), 344–345.
- Coelho, P.A. (1997) Revisão das espécies de Thalassinidea encontradas em Pernambuco, Brasil (Crustacea, Decapoda). *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 25, 137–161.
- Coelho, P.A. & Coelho-Santos, M.A. (1993) A família Callianassidae no litoral do Estado de Pernambuco (Crustacea, Decapoda, Thalassinidea). *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 22, 243–257. [Dated 1991/93, published 1993].
- Coelho, P.A. & Koenig, M.L. (1972) A distribuição dos crustáceos pertencentes às ordens Stomatopoda, Tanaidacea e Isopoda no Norte e Nordeste do Brasil. *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 13, 245–260.
- Coelho P.A. & Ramos, M. (1972) A constituição e a distribuição da fauna de decápodos do litoral leste da América do Sul entre as latitudes de 5° N e 39° S. *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 13, 133–236.
- Coelho, P.A. & Ramos-Porto, M. (1985) Sinopse dos crustáceos decápodos brasileiros (famílias Scyllaridae, Palinuridae, Nephropidae, Parastacidae e Axiidae). *Anais da Universidade Federal Rural de Pernambuco, Ciências Biológicas*, 8/10, 47–88. [Dated 1983/85, published 1985].
- Coelho, P.A. & Ramos-Porto, M. (1986) Sinopse dos crustáceos decápodos brasileiros (famílias Callianassidae, Callianideidae, Upogebiidae, Parapaguridae, Paguridae, Diogenidae). *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 19, 27–53. [Dated 1985/86, published 1986].
- Coelho, P.A. & Ramos-Porto, M. (1995) Crustáceos da região de Tamandaré, Estado de Pernambuco, Brasil. *Boletim Técnico Científico do CEPENE*, 3(1), 57–80.
- Coelho, P.A. & Ramos-Porto, M. (1998) Malacostraca. Eucarida. Palinuridea. In: P.S. Young (Ed.), *Catalogue of Crustacea of Brazil*. Museu Nacional, Rio de Janeiro, pp. 387–392.
- Coelho, P.A., Ramos-Porto, M. & Calado, T.C.S. (1983) Litoral de Alagoas e Sergipe: Decapoda. *Anais da Sociedade Nordestina de Zoologia*, 1 (1), 133–155.
- Coelho, P.A., Ramos-Porto, M. and Koenig, M.L. (1978) Crustáceos marinhos do Brasil, do Uruguai e da Argentina (ao norte de Mar del Plata), considerações biogeográficas. *Anais da Universidade Federal Rural de Pernambuco, Ciências Biológicas*, 2/3, 227–256. [Dated 1977/78, published 1978].
- Coelho, P.A., Ramos-Porto, M. & Koenig, M.L. (1980) Biogeografia e bionomia dos crustáceos do litoral equatorial brasileiro. *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 15, 7–138.
- Coelho, P.A., Ramos-Porto, M. & Melo, G.A.S. (1990) Crustáceos decápodos do Estado de Alagoas. *Anais da Sociedade Nordestina de Zoologia*, 3, 21–34.
- Coelho, P.A. & Rattacaso, M.C.A. (1988) Revisão das espécies de *Upogebia* encontradas em Pernambuco, Brasil (Crustacea, Decapoda, Thalassinidea). *Revista Brasileira de Zoologia*, 5 (3), 381–392.
- Coelho, P.A., Tenório, D.O., Ramos-Porto, M. & Mello, R.L.S. (2004) A fauna bêntica do Estado de Pernambuco. In: E. Eskinazi-Leça, Neumann-Leitão, S. & Costa, M.F. (Eds), *Oceanografia: um cenário tropical*. Bagaço, Recife, pp. 477–527.
- Coelho, P.A. & Vasconcelos, E.M.S. (1993) Ocorrência de *Justitia longimanus* (Milne Edwards, 1837), (Crustacea, Decapoda, Palinuridae), no litoral do Estado do Rio Grande do Norte. *Boletim Técnico Científico do CEPENE*, 1 (1), 25–28.
- Coelho Filho, P.A. (2006) Checklist of the Decapods (Crustacea) from the outer continental shelf and seamounts from Northeast of Brazil - REVIZEE Program (NE III). *Zootaxa*, 1184, 1–27.
- Dana, J.B. (1853) An isothermic oceanic chart, illustrating the geographical distribution of marine animal. *American Journal of Science*, 16, 153–167, 314–327.
- Fausto Filho, J. (1966) Primeira contribuição ao inventário dos crustáceos decápodos marinhos do Nordeste Brasileiro. *Arquivos da Estação de Biologia Marinha da Universidade Federal do Ceará*, 6 (1), 31–37.
- Fausto Filho, J. (1968) Terceira contribuição ao inventário dos crustáceos decápodos marinhos do Nordeste Brasileiro. *Arquivos de Ciências do Mar*, 8 (1), 43–45.
- Fausto Filho, J. (1970) Quarta contribuição ao inventário dos crustáceos decápodos marinhos do Nordeste Brasileiro.

- Arquivos de Ciências do Mar*, 10 (1), 55–60.
- Fausto Filho, J. (1977) Sobre a ocorrência de *Palinustus truncatus* (A. Milne Edwards, 1880), no litoral brasileiro e de *Panulirus echinatus* Smith 1869, no litoral do Estado do Ceará, Brasil (Crustacea, Decapoda, Palinuridae). *Arquivos de Ciências do Mar*, 17 (1), 75–76.
- Fausto Filho, J. (1979) Crustáceos estomatópodos e decápodos dos substratos de areia do nordeste brasileiro. *Arquivos de Ciências do Mar*, 19 (1/2), 45–56.
- Fausto Filho, J., Matthews, H.R. & Lima, H.H. (1966) Nota preliminar sobre a fauna dos bancos de lagostas no Ceará. *Arquivos da Estação de Biologia Marinha da Universidade Federal do Ceará*, 6 (2), 127–130.
- Forest, J. & de Saint Laurent, M. (1967) Campagne de la “Calypso” au large des côtes atlantiques de l’Amérique du Sud (1961–1962). 6. Crustacés Décapodes: Pagurides. *Annales de l’Institut Océanographique*, 45 (2), 47–171.
- Gomes Corrêa, M.M. (1968a) Sobre as espécies de *Upogebia* Leach do litoral brasileiro, com descrição de uma espécie nova. *Revista Brasileira de Biologia*, 28 (2), 97–109.
- Gomes Corrêa, M.M. (1968b) Descrição de uma nova espécie do gênero *Lepidopa* Stimpson e sua ocorrência no litoral brasileiro (Decapoda, Albuneidae). *Revista Brasileira de Biologia*, 28 (1), 77–86.
- Haig, J. (1966) Campagne de la “Calypso” au large des côtes atlantiques de l’Amérique du Sud (1961–1962). 1. Porcellanid crabs (Crustacea Anomura). *Annales de l’Institut Océanographique*, 44, 351–358.
- Henderson, J.R. (1888) Report of the Crustacea Anomura collected by the H.M.S. “Challenger” during the years 1873–1876. *Report on the Scientific Results of the Voyage of H. M. S. “Challenger”*, *Zoology*, 27 (1), 1–221.
- Hendrickx, M.E. (1995) Checklist of brachyuran crabs (Crustacea: Decapoda) from the eastern tropical Pacific. *Bulletin de l’Institut Royal des Sciences Naturelles de Belgique*, 65, 125–150.
- Komai, T. (2004) Deep-sea shrimps of the genus *Glyphocrangon* A. Milne-Edwards (Crustacea, Decapoda, Caridea, Glyphocrangonidae) from off southeastern coast of Brazil collected during the Revizee program. *Arquivos do Museu Nacional*, 62 (1), 31–44.
- Lemaitre, R. (1989) Revision of the genus *Parapagurus* (Anomura: Paguroidea: Parapaguridae), including redescription of the western Atlantic species. *Zoologische Verhandlungen*, 253, 1–106.
- Mabesoone, J.M., Kempf, M. & Coutinho, P.N. (1972) Characterization of surface sediments on the northern and eastern Brazilian shelf. *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 13, 41–48.
- Martin, J.W. & Davis, G.E. (2001) An updated classification of the recent Crustacea. *Natural History Museum of Los Angeles County, Science Series*, 39, 1–124.
- Matthews, L.H.A. (1926) A fauna dos recifes de Pernambuco, Brasil. *Boletim do Museu Nacional*, 2 (2), 1–20.
- McLaughlin, P.A. (1981) Revision of *Pylopagurus* and *Tomopagurus* (Crustacea: Decapoda: Paguridae), with the descriptions of new genera and species. *Bulletin of Marine Science*, 31 (1), 1–30.
- McLaughlin, P.A. & Provenzano Jr., A.J. (1974) Hermit crabs of the genus *Paguristes* (Crustacea: Decapoda: Diogenidae) from the western Atlantic. Part 1. The *Paguristes tortugae* complex, with notes on variation. *Bulletin of Marine Science*, 24 (1), 165–234.
- Melo, G.A.S. (1985) Taxonomia e padrões distribucionais e ecológicos dos Brachyura (Crustacea, Decapoda) do litoral sudeste do Brasil. Ph.D. Thesis, Universidade de São Paulo, 215pp.
- Melo, G.A.S. (1990) A presença, no litoral sudeste brasileiro, de espécies de Brachyura (Crustacea: Decapoda) originárias das regiões biogeográficas Magelânica e Argentina no Atlântico Sul. *Atlântica*, 12 (2), 71–83.
- Melo, G.A.S. (1999) *Manual de identificação dos Crustacea Decapoda do litoral brasileiro: Anomura, Thalassinidea, Palinuridea, Astacidea*. Editora Plêiade, São Paulo, 551 pp.
- Melo-Filho, G.A.S. (1998). Malacostraca. Eucarida. Chirostylidae and Galatheidae. In: P.S. Young (Ed.), *Catalogue of Crustacea of Brazil*. Museu Nacional, Rio de Janeiro, 393–397.
- Melo-Filho, G.A.S. (2006) Reports on the results of the N. Oc. “Prof. W. Besnard” expeditions to the southern coast of Brazil under the Revizee Program: Chirostylidae and Galatheidae (Crustacea: Decapoda: Anomura). *Zootaxa*, 1238, 1–22.
- Melo-Filho, G.A.S. & Melo, G.A.S. (1994) Three new species of *Munida* (Crustacea: Anomura: Galatheidae) from the Brazilian coast. *Papéis Avulsos de Zoologia*, 39 (5), 49–60.
- Melo-Filho, G.A.S. & Melo, G.A.S. (2001) Espécies do gênero *Munida* Leach (Crustacea, Decapoda, Galatheidae) distribuídas na costa do Brasil. *Revista Brasileira de Zoologia*, 18 (4), 1135–1175.
- Miers, E.J. (1886) Report on the Brachyura dredged by H. M. S. “Challenger” during the years 1873–75. *Report on the Scientific results of the Voyage of H. M. S. Challenger during the Years 1873–76*, *Zoology*, 17 (49), 1–362.
- Palacio, F.J. (1982) Revisión zoogeográfica marina del sur del Brasil. *Boletim do Instituto Oceanográfico*, 31 (1), 69–92.
- Pocock, R.L. (1890) Crustacea. In: H.N. Ridley (Ed.), Notes on the Zoology of Fernando Noronha. *Journal of the Linnean Society of London, Zoology*, 20, 506–526.
- Ramos-Porto, M., Muniz, A.P.M., Silva, K.C.A., Cintra, I.H.A. & Viana, G.F.S. (2003) Camarões da subordem Pleocyemata Burkenroad, 1963 capturados durante pescarias experimentais para o Programa REVIZEE/NORTE (Crustacea, Decapoda). *Boletim Técnico Científico do CEPNOR*, 3 (1), 77–106.
- Ramos-Porto, M., Silva, K.C.A., Viana, G.F.S. & Cintra, I.H.A. (2000a) Camarões de profundidade coletados no Norte

- do Brasil (Crustacea: Penaeidea e Caridea). *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 28 (1), 71–85.
- Ramos-Porto, M., Viana, G.F.S., Silva, K.C.A., Cintra, I.H.A. & Coelho, P.A. (2000b) *Stereomastis sculpta* (Smith, 1880) (Decapoda: Polycheloidea: Polychelidae) in Brazilian waters. *Nauplius*, 8 (2), 249–251.
- Rathbun, M.J. (1900) Results of the Branner-Agassiz Expedition to Brazil. I. The Decapod and Stomatopod Crustacea. *Proceedings of the Washington Academy of Sciences*, 2, 133–156.
- Rieger, P.J. (1998) Malacostraca. Eucarida. Paguroidea. In: P.S. Young (Ed.), *Catalogue of Crustacea of Brazil*. Museu Nacional, Rio de Janeiro, pp. 413–429.
- Rodrigues, S.A. (1971) Mud shrimps of the genus *Callianassa* Leach from the Brazilian coast (Crustacea, Decapoda). *Arquivos de Zoologia*, 20 (3), 191–223.
- Rodrigues, S.A. & de Carvalho, H.A. (1972) *Marcusiarius lemoscatroi*, g. n., sp. n., primeira ocorrência da família Axiidae (Crustacea, Decapoda, Thalassinidea) no Brasil. *Ciência e Cultura*, 24 (6) (Suppl.), 337.
- Rodrigues, S.A. & Kensley, B. (1991) *Eutrichocheles pindatyba*, a new axiid shrimp (Crustacea: Decapoda: Thalassinidea) from Brazil. *Proceedings of the Biological Society of Washington*, 104 (3), 556–560.
- Rodrigues, S.A. & Manning, R.B. (1992) Two new callianassid shrimps from Brazil (Crustacea: Decapoda: Thalassinidea). *Proceedings of the Biological Society of Washington*, 105 (2), 324–330.
- Rodrigues S.A. & Shimizu, R.M. (1992) Description of a new *Axianassa* (Crustacea: Decapoda: Thalassinidea) from Brazil, and its first larval stage. *Proceedings of the Biological Society of Washington*, 105 (2), 317–323.
- Rodrigues, S.A. & Shimizu, R.M. (1998) Malacostraca. Eucarida. Thalassinidea. In: P.S. Young (Ed.), *Catalogue of Crustacea of Brazil*. Museu Nacional, Rio de Janeiro, pp. 379–385.
- Rodrigues da Costa, H. (1960) *Pachycheles haigae*, nueva especie de la familia Porcellanidae (Crustacea, Anomura). *Neotropica*, 6, 21–24.
- Rodrigues da Costa, H. (1962) Notas sobre os Hippidea da costa brasileira. *Centro de Estudos Zoológicos da Universidade do Brasil, Avulso*, 14, 1–12.
- Rodrigues da Costa, H. (1968) Crustacea Brachyura récoltés par les dragages de la “Calypso” sur les côtes brésiliennes (1962). *Recueil des Travaux de la Station Marine d’Endoume*, 43 (59), 333–343.
- Rodriguez, I.T., Hernandez, G. & Felder, D.L. (2005) Review of the Western Atlantic Porcellanidae (Crustacea: Decapoda: Anomura) with new records, systematic observations and comments on biogeography. *Caribbean Journal of Science*, 41 (3), 544–582.
- Serejo, C.S., Young, P.S., Cardoso, I.A., Tavares, C.R. & Abreu Junior, C.R. (2006) Abundância, diversidade e zonação de Crustacea no talude da costa central do Brasil (11° 22° S) coletado pelo Programa REVIZEE Pesca Score Central. In: Costa, P.A.S., Martins, A.S., Olavo, G. (Org.), Pesca e potenciais de exploração de recursos vivos da região central da Zona Econômica Exclusiva brasileira. Parte 2. Rio de Janeiro: Museu Nacional Universidade Federal do Rio de Janeiro, *in press*.
- Silva, K.C.A., Cintra, I.H.A., Ramos-Porto, M. & Viana, G.F.S. (2003) Lagostas capturadas durante pescarias experimentais para o Programa REVIZEE/NORTE (Crustacea, Nephropoidea, Eryonoidea, Palinuroidea). *Boletim Técnico Científico do CEPNOR*, 3 (1), 21–35.
- Silva, K.C.A., Muniz, A.P.M., Ramos-Porto, M., Viana, G.F.S. & Cintra, I.H.A. (2002a) Camarões da superfamília Penaeoidea Rafinesque, 1815, capturados durante pescarias experimentais para o programa REVIZEE/Norte (Crustacea: Decapoda). *Boletim Técnico Científico do CEPNOR*, 2 (1), 9–40.
- Silva, K.C.A., Ramos-Porto, M., Cintra, I.H.A., Muniz, A.P.M. & Silva, M.C.N. (2002b) Crustáceos capturados durante o programa REVIZEE na costa Norte Brasileira. *Boletim Técnico Científico do CEPNOR*, 2 (1), 97–108.
- Smith, S.I. (1869) Notice of the Crustacea collected by Prof. C. F. Hartt on the coast of Brazil in 1867. *Transactions of the Connecticut Academy of Arts and Sciences*, 2, 1–41.
- Stramma, L., Ikeda, Y. & Peterson, R.G. (1990) Geostrophic transport in the Brazil Current region north of 20° S. *Deep-Sea Research*, 37 (12), 1875–1886.
- Tavares, C.R. & Young, P.S. (2002) Nephropidae (Crustacea, Decapoda) collected by the Revizee Score-Central program from off Bahia to Rio de Janeiro states, Brazil. *Arquivos do Museu Nacional*, 60 (2), 79–88.
- Tavares, M. (1998) Malacostraca. Eucarida. Nephropidae. In: P.S. Young (Ed.), *Catalogue of Crustacea of Brazil*. Museu Nacional, Rio de Janeiro, pp. 377–378.
- Veloso, V.G. (1998) Malacostraca. Eucarida. Porcellanidae. In: P.S. Young (Ed.), *Catalogue of Crustacea of Brazil*. Museu Nacional, Rio de Janeiro, pp. 399–405.
- Veloso, V.G. & Melo, G.A.S. (1993) Taxonomia e distribuição da Família Porcellanidae (Crustacea, Decapoda, Anomura) no litoral brasileiro. *Iheringia, Zoologia*, 75, 171–186.
- Werding, B., Hiller, A. & Lemaitre, R. (2003) Geographic and depth distributional patterns of western Atlantic Porcellanidae (Crustacea: Decapoda: Anomura), with and updated list of species. *Memoirs of the Museum of Victoria*, 60 (1), 79–85.
- White, A. (1847) *List of the specimens of Crustacea in the collection of the British Museum*. British Museum, London, pp. i-viii, 1-143.

- Williams, A.B. (1993) Mud shrimps, Upogebiidae, from the western Atlantic (Crustacea: Decapoda: Thalassinidea). *Smithsonian Contributions to Zoology*, 544, 1–77.
- Young, P.S. (Ed). (1998) *Catalogue of Crustacea of Brazil*, Museu Nacional, Rio de Janeiro, 717 pp.

Note: Recently, the species *Munida coltroi* was described by Melo-Filho & Melo (2006) from Ceará State coast, adding a new species to the study area.

- Melo-Filho, G.A.S. & Melo, G.A.S. (2006) A new species of the genus *Munida* Leach (Crustacea: Decapoda: Galatheididae) from the northeast coast of Brazil, *Nauplius*, 14, 49–54.