

# A new western Atlantic species of *Eurysquilla* (Crustacea: Stomatopoda: Eurysquilloidea) with a key to the world species

DÉBORA LUCATELLI<sup>1</sup>, SHANE T. AHYONG<sup>2</sup>, LUIS ERNESTO ARRUDA BEZERRA<sup>3</sup>  
AND PAULO JORGE PARREIRA DOS SANTOS<sup>1</sup>

<sup>1</sup>Universidade Federal de Pernambuco, Programa de Pós-Graduação em Oceanografia, Departamento de Oceanografia, Avenida Arquitetura s/n, Cidade Universitária, CEP 50740-550, Recife, PE, Brazil, <sup>2</sup>Australian Museum, 6 College Street, Sydney, NSW 2010, Australia, and School of Biological, Earth and Environmental Sciences, University of New South Wales, NSW 2052, Australia, <sup>3</sup>Universidade Federal Rural do Semi-Árido (UFERSA), Departamento de Ciências Animais, Avenida Francisco Mota, 572, Mossoró, Rio Grande do Norte, Brazil, CEP: 99625-900 and Programa de Pós-Graduação em Ciências Marinhas Tropicais, Instituto de Ciências do Mar (LABOMAR), Fortaleza, Ceará, Brazil

*A new species of eurysquilloid stomatopod, Eurysquilla petronioi sp. nov., is described from the tropical western Atlantic. Eurysquilla petronioi sp. nov. is the fourteenth species of the genus to be recognized worldwide, the fifth species from the western Atlantic and the second from Brazilian waters. It is most closely related to E. maiaquesensis from Puerto Rico, but differs chiefly in having an unarmed versus apically spinous rostral plate and 6 or 7 rather than 8–10 teeth on dactylus of raptorial claw. A key to the species of the genus is provided.*

**Keywords:** Hoplocarida, Eurysquillidae, South Atlantic, western Atlantic, Barbados, north-east Brazil, taxonomy

Submitted 10 August 2012; accepted 21 August 2012; first published online 29 October 2012

## INTRODUCTION

The stomatopod crustacean superfamily, Eurysquilloidea Manning, 1977, is supported as monophyletic by the ventrally originating outer intermediate and lateral denticles of the telson (Ahyong & Harling, 2000). Eurysquilloidea includes a single family Eurysquillidae Manning, 1977, with 31 living species and one fossil, distributed in six genera worldwide (Ahyong, 2001, 2010; De Angeli & Beschin, 2006), with only *Eurysquilla* represented in the Americas. Preliminary studies of the phylogeny of *Eurysquilla* (Cappola & Manning, 1999; Ahyong & Harling, 2000) found the genus to be paraphyletic and requiring further research. A comprehensive phylogenetic analysis of the Eurysquilloidea is currently in preparation by the second author.

Prior to the present study, *Eurysquilla* Manning, 1963 included thirteen species worldwide. Four species are known from the Indo-Pacific: *E. crosnieri* Moosa, 1991; *E. foresti* Moosa, 1986; *E. pacifica* Manning, 1975; and *E. sewelli* (Chopra, 1939). Three species are known from the Eastern Pacific: *E. pumae* Hendrickx & Salgado-Barragán, 1987; *E. veleronis* (Schmitt, 1940); and *E. solari* Manning, 1970. Two species, *E. galathea* Manning, 1977 and *E. leloeuffi* Manning, 1977 occur in the eastern Atlantic and four species are known from the western Atlantic Ocean: *E. chacei* Manning, 1969; *E. holthuisi* Manning, 1969; *E. maiaquesensis* (Bigelow, 1901); and *E. plumata* (Bigelow, 1901). A new western Atlantic species is described herein.

## MATERIALS AND METHODS


Specimens studied herein are deposited in the carcinological collection of the Museu de Oceanografia Petrônio Alves Coelho of Universidade Federal de Pernambuco, Recife, Brazil (MOUFPE) and the National Museum of Natural History, Washington, DC (USNM). The Brazilian specimen was collected by N/RB 'Astro Garoupa' of PETROBRAS (Petróleo do Brasil S/A), as part of the Project 'Monitoramento da Bacia Potiguar/Rio Grande do Norte/ BR' (BPot). The sampling design consisted of 69 stations, collected between 2002 and 2004 with a 70-l rectangular dredge pulled for 5 minutes at a speed of 2 knots. The specimen studied herein was preserved in 70% ethanol. Total length (TL) is measured along the dorsal midline from the apex of the rostral plate to the apices of the submedian teeth of the telson. Carapace length (CL) is measured along the dorsal midline. The propodal index (PI) of the raptorial claw is given as 100CL/propodus length. Morphological terminology and the systematic classification follow those of Ahyong *et al.* (2008) and Ahyong (2012).

## RESULTS

### SYSTEMATICS

Order STOMATOPODA Latreille, 1817  
Suborder UNIPELTATA Latreille, 1825  
Superfamily EURYSQUILLOIDEA Manning, 1977  
Family EURYSQUILLIDAE Manning, 1977  
*Eurysquilla petronioi* sp. nov.  
(Figure 1)

**Corresponding author:**  
D. Lucatelli  
Email: dla.lucatelli@gmail.com



**Fig. 1.** *Eurysquilla petronioi* sp. nov.: (A–M) female holotype, total length (TL) 20 mm, USNM 111111; (N) male paratype, TL 11 mm, MOUFPE 14201. (A) Carapace and anterior appendages; (B) right eye; (C) right raptorial claw; (D) right lateral processes of thoracic somites 5–8; (E) thoracic somite 5, right lateral view; (F–H) right pereopods 1–3, posterior view; (I) thoracic somite 8 sternal keel; (J) posterior abdominal somites, telson and right uropod; (K) right uropod, ventral view; (L) telson, right lateral view; (M) telson, ventral view; (N) right pleopod 1 endopod, anterior view. Scale bars: A–H, J–M = 1.0 mm; I = 0.5 mm; N = 0.3 mm.

*Eurysquilla maiaguesensis*.—Manning, 1969: 254, 257 (Barbados specimen only; not *Lysiosquilla maiaguesensis* Bigelow, 1901).

#### TYPE MATERIAL

Holotype: USNM 111111, female (TL 20 mm), Barbados (~13°00'N 59°30'W), Caribbean Sea, Station NR1-3, 91.5–366 m (originally recorded as 50–200 fathoms), coll. J.B. Lewis, summer, 1960. Paratype: MOUFPE 14201, male (TL 11.25 mm), Rio Grande do Norte, Brazil, 04°36.64'S 36°44.89'W, 368 m, BPot II No. 33, dredge, 20 May 2003.

#### COMPARATIVE MATERIAL

*Eurysquilla maiaguesensis* (Bigelow, 1901): USNM 64824, male lectotype (TL 23 mm), Mayaguez Harbor, 294.6–314.8 m (originally recorded as 161–172 fathoms), sand and mud, beam trawl, U.S.F.C. Steamer 'Fish Hawk', Station 6066, 20 January 1899; USNM 110939, 1 male paralectotype (TL 20 mm), 1 female paralectotype (broken), USNM 119137, male (TL 20.5 mm), Atlantic Ocean, north of Puerto Rico (18°31'30"N 66°14'55"W), 219.4 m (originally recorded as 120 fathoms), Johnson–Smithsonian Deep-Sea Expedition, Station 105, coll. P.S. Bartsch, 8 March 1933.

#### DIAGNOSIS

Cornea slightly broadened, subglobular, with 6 rows of midband ommatidia. Antennal protopod without papillae. Rostral plate slightly wider than long; apex rounded. Raptorial claw dactylus with 6 or 7 teeth on occlusal margin. Mandibular palp absent. Maxillipeds 1–4 each with epipod. Thoracic somites 5–8 and abdominal somites 1–5 posterolaterally unarmed. Male pleopod 1 endopod posterior endite without lateral lobe. Telson dorsal surface with posteriorly spined median and anterior intermediate carinae, surface otherwise smooth. Uropodal protopod broad, flattened, terminating in 2 primary spines; protopod inner margin lined with spines. Endopod length 4 or more times breadth.

#### DESCRIPTION

Dorsal integument smooth. Eye extending slightly beyond antennal peduncle segment 2; cornea slightly broadened, subglobular, set obliquely on stalk, with 6 rows of midband ommatidia. Ophthalmic somite anterior margin unarmed. Ocular scales broad, triangular, anteriorly convex, apices inclined directed laterally.

Antennular peduncle length 0.81–0.83 CL; segment 1 with small rounded lateral proximal lobe. Antennular somite dorsal processes spiniform, directed anteriorly. Antennal protopod without papillae; dorsally unarmed. Antennal scale length 3.17–3.50 times width, 0.37–0.49 CL; setose on anterior distal three-quarters and entire posterior margin.

Rostral plate slightly wider than long; subtriangular apex rounded, unarmed; lacking dorsal or ventral carina.

Carapace anterolateral angles broadly rounded; marginal carina faintly indicated posterolaterally.

Raptorial claw dactylus with 6 or 7 teeth on occlusal margin; outer margin broadly curved, uninflated, without proximal notch. Propodus longer than carapace length, when folded, distal margin extending posteriorly beyond ischium; occlusal margin fully pectinate, with 3 moveable spines proximally; PI 75 (male), 69 (female). Carpus unarmed dorsally; merus outer inferodistal angle rounded.

Mandibular palp absent. Maxillipeds 1–4 each with epipod. Maxilliped 5 basal segment unarmed; epipod absent.

Body dorsoventrally flattened, somites loosely articulated.

Thoracic somites 5–8 without dorsal carinae. Thoracic somite 5 lateral process obsolete, truncate, ventral lobe blunt. Thoracic somites 6–8 lateral process rounded, posterolaterally unarmed. Thoracic somite 8 sternal keel narrow, blunt, directed ventrally.

Pereopods 1–3 basal segment each with outer, ventrally directed spine; endopods elongate, spatulate, 1-segmented, shortest on pereopod 1, longest on pereopod 3, setose on outer margin and margin of inner distal half.

Abdominal somites 1–5 pleura unarmed posterolaterally, with low marginal carina; somite 5 with indistinct intermediate swelling. Abdominal somite 6 smooth, without submedian carina but pair of submedian spines on posterior margin; intermediate and lateral carina with posterior spine; ventrolateral margin anterior to uropodal articulation unarmed; sternum posterior margin unarmed.

Male pleopod 1 endopod with tube process distinctly longer than hook process; posterior endite with outer margin non-setose, without lateral lobe.

Telson wider than long, subtriangular. Submedian teeth moveable, spiniform, separated by narrow fissure; submedian denticles absent. Intermediate teeth slender, apices slightly deflected dorsally, extending posteriorly almost to level of base of submedian teeth; with rounded, upright dorsal lobe; with 2 spiniform intermediate denticles, outer arising ventrally, inner on rounded lobe. Lateral teeth slender; apices slightly deflected dorsally, with rounded, mesially inclined dorsal lobe and 1 spiniform lateral denticle arising ventrally. Telson dorsal surface with median and anterior intermediate carinae, surface otherwise smooth. Median carina low, smooth, uninterrupted; apex spiniform. Anterior intermediate carinae slightly convergent posteriorly, smooth, uninterrupted; apex spiniform. Marginal carina indistinct. Telson ventral surface lacking postanal carina.

Uropodal protopod broad, flattened, distinctly wider than endopod, terminating in 2 primary spines; inner primary spine about twice as long as outer, with small, low spine or lobe at midlength of outer margin; protopod unarmed dorsally except for small dorsal spine above proximal exopod articulation; protopod inner margin lined with 13–20 spines.

Uropodal exopod proximal segment unarmed dorsally; with distoventral spine; inner margin broadly convex; outer margin with 4 or 5 graded, moveable spines, distal spine longest, arcuate, reaching to distal quarter of distal segment. Exopod distal segment ovate, unarmed, longer than proximal segment. Endopod length 4.0–4.5 times breadth, unarmed, entire margin setose.

#### COLOUR IN LIFE

Not known. In preservation completely faded.

#### MEASUREMENTS

Male (N = 1) TL 11 mm, female (N = 1) TL 20 mm. Other measurements of holotype: CL 3.5 mm, antennular peduncle length 2.9 mm, antennal scale length 1.7 mm, raptorial claw propodus length 5.1 mm.

**ETYMOLOGY**

The specific epithet is in honour of Dr Petrônio Alves Coelho, who recently passed away, for his contribution to the field of carcinology in Brazil.

**HABITAT**

The Brazilian specimen was found at 368 m depth, with salinity 35.23‰ and temperature 11.1°C. Sediment was stratified, with a light-coloured top layer of silt, fine sand and bioclasts. The base was dark-coloured clay. It has been classified as sandy loam (Freire *et al.*, 1997).

**DISTRIBUTION**

Presently known only from Barbados and Rio Grande do Norte, Brazil; 91.5–368 m (Figure 2).

**REMARKS**

*Eurysquilla petronioi* sp. nov. is the fourteenth species of the genus to be recognized. It is readily separated from *E. foresti*, *E. galatheae*, *E. lelouffi*, *E. plumata*, *E. pumae*, *E. sewelli*, *E. solari* and *E. veleronis* by having a broad, mesially multispinous uropodal protopod, rather than a slender Y-shaped protopod that is armed at most with a single spine on the inner margin. Like *E. petronioi*, the remaining species of *Eurysquilla* have a broad uropodal protopod with a multi-spinous inner margin. *Eurysquilla petronioi* is separable from *E. chacei*, *E. holthuisi*, *E. crosnieri* and *E. pacifica* by its broadened and subglobular, rather than strongly bilobed cornea.

The new species most closely resembles *E. maiaguesensis* from Puerto Rico, sharing a similar eye, uropod and telson structure. The two species differ in the form of the rostral plate (distally rounded and slightly wider than long in *E. petronioi*; distally spinous and as long as or longer than wide in *E. maiaguesensis*), the armature of the dactylus of the raptorial claw (6 or 7 in *E. petronioi*; 8–10, usually 9, in *E. maiaguesensis*) and slenderness of the uropodal endopod (4–4.5 times as long as wide in *E. petronioi*; more than 6 times as long as wide in *E. maiaguesensis*). Other differences are in the number of outer uropodal exopod spines (4 or 5 in *E. petronioi*; 6 in *E. maiaguesensis*) and the length of the distal uropodal exopod spine, which reaches the distal quarter of the distal exopod segment in *E. petronioi*, rather than the end in *E. maiaguesensis*.

The holotype and paratype agree in most respects. At TL 11 mm, however, the young male paratype is approximately half the size of the female holotype (TL 20 mm) and exhibits several size related differences: the rostral plate is slightly more triangular in shape, the eyes are more elongate, 6 instead of 7 teeth are present on the dactylus of the raptorial claw, the telson carinae are less pronounced, the dorsal lobes on the intermediate and lateral primary teeth of the telson are less upright and the inner uropodal spines are fewer in number (13–16 versus 18–20). Similar allometric changes are evident in other species of *Eurysquilla* including *E. maiaguesensis*.

*Eurysquilla petronioi* sp. nov. is the fifth western Atlantic species of the genus and the second from Brazilian waters, alongside *E. plumata*. A key to the species of the genus is given below.



Fig. 2. Distribution of *Eurysquilla petronioi* sp. nov., in the western Atlantic: holotype (◆) and paratype (●).

**KEY TO THE SPECIES OF EURYSQUILLA MANNING, 1963**

1. Uropodal protopod distally Y-shaped, slender, about as wide as endopod; inner margin with at most one lobe or spine . . . . . 2
  - Uropodal protopod broad, flattened, distinctly wider than endopod; inner margin lined with row of five or more slender spines . . . . . 9
2. Uropodal protopod with spine on inner margin . . . . . 3
  - Uropodal protopod without spine on inner margin . . . . . 4
3. Cornea subglobular . . . . . *E. pumae*
  - Cornea elongate, distinctly widened . . . . . *E. veleronis*
4. Telson without carina or tubercles flanking median carina. Maxilliped 5 without epipod . . . . . *E. lelouffi*
  - Telson with irregular carina or row of tubercles flanking median carina Maxilliped 5 with epipod . . . . . 5
5. Rostral plate without apical spine. Cornea subglobular . . . . . *E. plumata*
  - Rostral plate with apical spine. Cornea distinctly widened . . . . . 6
6. Abdominal somite 4 posterior margin with spine above posterolateral spine . . . . . *E. sewelli*
  - Abdominal somite 4 posterior margin with posterolateral spine only, otherwise unarmed . . . . . 7

- 7. Abdominal somite 5 posterior margin with 2 pairs of spines (intermediate, lateral) . . . . . *E. foresti*  
 — Abdominal somite 5 posterior margin with 3 pairs of spines (intermediate, lateral, marginal) . . . . . 8
- 8. Rostral plate with carina of distal spine extending onto anterior half of surface. Intermediate carinae of thoracic somites 6—8 distinct . . . . . *E. galathea*  
 — Rostral plate with carina of distal spine extending onto anterior one-fifth of surface. Intermediate carinae of thoracic somites 6—8 indistinct, faintly indicated . . . . .  
 . . . . . *E. solari*
- 9. Uropodal protopod with rounded lobe between terminal primary spines . . . . . 10  
 — Uropodal protopod without rounded lobe between terminal primary spines . . . . . 11
- 10. Rostral plate distinctly longer than wide. Telson surface without tubercles between median carina and anterior intermediate carina . . . . . *E. holthuisi*  
 — Rostral plate as long as wide. Telson surface with tubercles between median carina and anterior intermediate carina . . . . . *E. chacei*
- 11. Cornea broad, set transversely on stalk. Rostral plate distinctly wider than long . . . . . 12  
 — Cornea little broadened, subglobular, set obliquely on stalk. Rostral plate longer than wide to slightly wider than long . . . . . 13
- 12. Abdominal somite 6 with 8 posterior spines . . . . .  
 . . . . . *E. pacifica*  
 — Abdominal somite 6 with 10 posterior spines . . . . .  
 . . . . . *E. crosnieri*
- 13. Rostral plate slightly wider than long, distally rounded, without spine . . . . . *E. petronioi* sp. nov.  
 — Rostral plate as long as or longer than wide, with apical spine . . . . . *E. maiaguesensis*

ACKNOWLEDGEMENTS

The authors are grateful to PETROBRAS for donating the paratype specimen and providing collection data. D.L. thanks CAPES for providing a doctoral scholarship and P.J.P.S. acknowledges a CNPq research fellowship. S.T.A thanks Rafael Lemaitre (USNM) and Karen Reed (USNM) for the loan of material and assistance in Washington, DC in 2000. Special thanks to Dr Jessor F. Souza-Filho, MSc Catarina L. Araújo-Silva and MSc Marina S.L.C. Araújo for comments on the manuscript.

REFERENCES

Ahyong S.T. (2001) Revision of Australian stomatopod Crustacea. *Records of the Australian Museum* Supplement 26, 1–326.

Ahyong S.T. (2010) Stomatopoda from the Philippines collected by the AURORA 2007 Expedition (Euryssquilloidea, Lysiosquilloidea and Gonodactyloidea). In Fransen C.H.J.M, De Grave S. and Ng P.K.L. (eds) *Studies on Malacostraca: Lipke Bijdeley Holthuis Memorial Volume. Crustaceana Monographs* 14. Leiden, The Netherlands: E.J. Brill, pp. 101–113.

Ahyong S.T. (2012) The marine fauna of New Zealand: mantis shrimps (Crustacea: Stomatopoda). *NIWA Biodiversity Memoir* 125, 1–112.

Ahyong S.T., Chan T.-Y. and Liao Y.-C. (2008) *A catalog of the mantis shrimps (Stomatopoda) of Taiwan*. 1st edition. Taipei: National Science Council, Taiwan, R.O.C.

Ahyong S.T. and Harling C. (2000) The phylogeny of the stomatopod Crustacea. *Australian Journal of Zoology* 48, 607–642.

Bigelow R.P. (1901) The Stomatopoda of Porto Rico. *Bulletin of the United States Fish Commission* 20, 149–160.

Chopra B. (1939) Stomatopoda. *The John Murray Expedition, 1933–34, Scientific Reports* 6, 137–181.

Cappola V. and Manning R.B. (1999) Relationships within the stomatopod genus *Eurysquilla*. In Schram F.R. and von Vaupel Klein J.C. (eds) *Proceedings of the Fourth International Crustacean Congress, 1998. Volume 1*. Leiden, The Netherlands: E.J. Brill, pp. 383–393.

De Angeli A. and Beschin C. (2006) Stomatopodi Terziari del Veneto (Italia Settentrionale). *Studi e Ricerche—Associazione Amici del Museo—Museo Civico 'G. Zannato' Montecchio Maggiore (Vicenza)* 13, 25–34.

Freire G.S.S., Cavalcanti V.M.M., Maia L.P. and Lima S.F. (1997) Classificação dos sedimentos da plataforma continental do Estado do Ceará. In *Anais do Simpósio de Geologia do Nordeste, Fortaleza-CE*, pp. 209–211.

Hendrickx M.E. and Salgado-Barragán J. (1987) A new species of stomatopod, *Eurysquilla pumae* (Crustacea: Stomatopoda: Eurysquilloidea), from the Gulf of California, Mexico. *Proceedings of the Biological Society of Washington* 100, 529–531.

Manning R.B. (1963) Preliminary revision of the genera *Pseudosquilla* and *Lysiosquilla* with descriptions of six new genera. *Bulletin of Marine Science of the Gulf and Caribbean* 13, 308–328.

Manning R.B. (1969) Stomatopod Crustacea of the Western Atlantic. *Studies in Tropical Oceanography* 8, i–viii, 1–380.

Manning R.B. (1970) Nine new American stomatopod crustaceans. *Proceedings of the Biological Society of Washington* 83, 99–114.

Manning R.B. (1975) *Eurysquilla pacifica*, a new stomatopod crustacean from New Britain. *Proceedings of the Biological Society of Washington* 88, 249–252.

Manning R.B. (1977) A monograph of the west African stomatopod Crustacea. *Atlantide Report* 12, 25–181.

Moosa M.K. (1986) Stomatopoda Crustacea. In Crosnier A. (ed.) *Résultats du Campagnes MUSORSTOM I & II Philippines 2. Mémoires du Muséum National d'Histoire Naturelle, Paris, series A, Zoologie*. 133, pp. 367–414, pl 1. [Dated 1985, published 1986].

Moosa M.K. (1991) The Stomatopoda of New Caledonia and Chesterfield Islands. In Richer de Forges B. (ed.) *Le benthos de fonds meubles des lagons de Nouvelle-Calédonie* 1. Paris: Editions de l'ORSTOM, pp. 149–219.

and

Schmitt W.L. (1940) The stomatopods of the west coast of America based on collections made by the Allan Hancock Expedition (1933–1938). *Allan Hancock Pacific Expedition* 5, 129–225, figures 1–33.

Correspondence should be addressed to:

D. Lucatelli  
 Universidade Federal de Pernambuco  
 Programa de Pós-Graduação em Oceanografia  
 Departamento de Oceanografia  
 Avenida Arquitetura s/n, Cidade Universitária, CEP 50740-550, Recife, PE, Brazil  
 email: dla.lucatelli@gmail.com